Mallalieu, Joseph ORCID:

https://orcid.org/0000-0002-1988-8594, Carrivick, Jonathan, Quincey, Duncan and Raby, Cassandra (2021) Ice-marginal lakes associated with enhanced recession of the Greenland Ice Sheet. Global and Planetary Change, 202. pp. 1-12.

Downloaded from: http://ray.yorksj.ac.uk/id/eprint/5240/

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version: https://www.journals.elsevier.com/global-and-planetary-change

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. Institutional Repository Policy Statement



Research at the University of York St John For more information please contact RaY at <u>ray@yorksj.ac.uk</u>

1 2 3	© 2021 J. Mallalieu, J.L. Carrivick, D.J. Quincey and C.L. Raby. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <u>http://creativecommons.org/licenses/by-nc-nd/4.0/</u>
4	
5	Ice-marginal lakes associated with enhanced recession of the Greenland Ice
6	Sheet
7	
8	Joseph Mallalieu ^{a,b} *, Jonathan L. Carrivick ^b , Duncan J. Quincey ^b , Cassandra L. Raby ^c
9	^a School of Humanities, York St John University, York, YO31 7EX, UK
10	^b School of Geography and water@leeds, University of Leeds, Leeds, LS2 9JT, UK
11	^c Institute of Integrative Biology, University of Liverpool, Liverpool, L69 7ZB, UK
12	*Corresponding author: j.mallalieu@yorksj.ac.uk
13	
14	HIGHLIGHTS
15	Prevalent and accelerating ice-margin recession in south-west Greenland from 1992
16	Contrasting ice-marginal environments demonstrate a heterogeneous response to warming
17	• Lacustrine ice-margins recede faster than terrestrial, but slower than marine margins
18	Lacustrine recession rates progressively outpaced terrestrial rates between 1987-2015
19	Significant correlations between lake parameters and recession rates are identified
20	
21	ABSTRACT
22	There has been a progressive increase in the number and area of ice-marginal lakes situated along the
23	south-western margin of the Greenland Ice Sheet (GrIS) since the 1980s. The increased prevalence of
24	ice-marginal lakes is notable because of their capacity to enhance mass loss and ice-margin recession
25	through a number of thermo-mechanical controls. Although such effects have been extensively
26	documented at alpine glaciers, an understanding of how ice-marginal lakes impact the dynamics of

27 the GrIS has been limited by a sparsity of observational records. This study employs the Landsat 28 archive to conduct a multi-decadal, regional-scale statistical analysis of ice-margin advance and 29 recession along a ~5000 km length of the south-western margin of the GrIS, incorporating its 30 terrestrial, lacustrine and marine ice-margins. We reveal an extended and accelerating phase of ice-31 margin recession in south-west Greenland from 1992 onwards, irrespective of margin type, but also 32 observe considerable heterogeneity in the behaviour of the different ice-marginal environments. Marine ice-margins exhibited the greatest magnitude and variability in ice-margin change, however 33 34 lacustrine termini were notable for a progressive increase in ice-margin recession rates from 1987 to 35 2015, which increasingly outpaced those measured at terrestrial ice-margins. Furthermore, significant 36 correlations were identified between lake parameters and rates of lacustrine ice-margin recession, 37 including lake area, latitude, altitude and the length of the lake – ice-margin interface. These results 38 suggest that ice-marginal lakes have become increasingly important drivers of ice-margin recession 39 and thus mass loss at the GrIS, however further research is needed to better parameterise the causal 40 connections between ice-marginal lake evolution and enhanced ice-margin recession. More widely, a detailed understanding of the impacts of ice-marginal lakes on ice-margin dynamics across Greenland 41 42 is increasingly necessary to accurately forecast the response of the ice sheet to enhanced ice-marginal 43 lake prevalence and thus refine projections of recession, mass loss and sea level rise.

44

45 **KEYWORDS**: Greenland Ice Sheet; ice-marginal lake; proglacial lake; glacier dynamics; meltwater

46

47 **1. INTRODUCTION**

Since a period of near equilibrium mass balance in the 1980s, rates of mass loss at the Greenland Ice Sheet (GrIS) have generally accelerated in response to increased atmospheric and oceanic warming (Hanna et al., 2013; Shepherd et al., 2020). Over the same time period, enhanced rates of meltwater runoff (Hanna et al., 2008; Trusel et al., 2018) have coincided with a progressive increase in the number and area of ice-marginal lakes situated along the south-western margin of the GrIS (Carrivick 53 and Quincey, 2014; How et al. 2021). The presence of ice-marginal lakes is significant because of their 54 capacity to regulate ice-margin dynamics through a number of thermo-mechanical controls, including the onset and promotion of calving (Carrivick and Tweed, 2013). In particular, ice-marginal lake 55 formation and expansion is typically associated with enhanced rates of mass loss and ice-margin 56 57 recession (e.g. Kirkbride, 1993; Boyce et al., 2007; Schomacker, 2010; Basnett et al., 2013; Brun et al., 58 2019; King et al., 2019; Tsutaki et al., 2019; Liu et al., 2020; Sutherland et al., 2020). However, whilst 59 the effects of ice-marginal lakes on alpine glacier dynamics have been increasingly well-documented, 60 knowledge of their effects on the dynamics of ice sheets is presently limited by a sparsity of 61 observational records (Mallalieu et al., 2017, 2020). A detailed understanding of the impacts of ice-62 marginal lakes on ice-margin dynamics across Greenland is therefore increasingly necessary to 63 accurately forecast the response of the ice sheet to enhanced ice-marginal lake prevalence and thus 64 further refine projections of mass loss and sea level rise.

65

66 An analysis of outlet glacier extent by Warren (1991) revealed significant variability in the behaviour of the terrestrial, lacustrine and marine outlets of the GrIS throughout the mid-20th century, despite 67 68 having undergone comparable climatic forcing. Both lacustrine and marine outlets were found to 69 exhibit much greater variability in frontal behaviour than their terrestrial counterparts due to their 70 partial decoupling from climatic forcing and the increased role of topographic and bathymetric 71 controls on terminus advance and recession. However, subsequent analyses of ice-margin behaviour 72 and extent in Greenland have omitted measurements from lacustrine ice-margins, instead focusing 73 on changes at the major marine-terminating outlets (e.g. Howat et al., 2008; Howat and Eddy, 2011; 74 Catania et al., 2018), and the terrestrial termini of peripheral glaciers and ice caps (PGICs) (e.g. Citterio 75 et al., 2009; Leclercq et al., 2012; Rastner et al., 2012; Bjørk et al., 2018). In addition, the few studies 76 that have incorporated measurements of ice-margin change from terrestrial outlets of the main ice sheet typically include a sparse number of terrestrial data points (e.g. Moon and Joughin, 2008; Carr 77 78 et al., 2013; Mouginot et al., 2019), or concern a relatively limited footprint in south-east Greenland (e.g. Kargel et al., 2012; Mernild et al., 2012). As a consequence, the relative magnitude of recent
changes at the terrestrial, lacustrine and marine margins of the GrIS remain unknown.

81

82 The long temporal record of the Landsat image archive, now extending into its fourth decade with the 83 launch of Landsat 8 (Roy et al., 2014), provides a unique opportunity to perform a multi-decadal, regional-scale analysis of ice-margin extent for the disparate ice-marginal environments of the GrIS 84 85 between the 1980s and the present day. South-west Greenland is the optimal site for such an analysis 86 because: (i) it has the greatest regional concentration of land-terminating, and thus lacustrine, 87 margins of the GrIS (Figure 1); (ii) the region has experienced some of the highest increases in mean 88 annual air temperatures recorded in the Arctic since the 1990s (Carr et al., 2013; Ding et al., 2014); 89 and (iii) the region is forecast to undergo some of the greatest rates of ice-margin recession and 90 reductions in ice-cover over the next millennium (Aschwanden et al., 2019).

91



- 93 Figure 1. Study location in south-west Greenland. The spatial extent of the analysis is illustrated with
- 94 false-colour Landsat scenes from 2015 (see Table 1).

95 [SINGLE COLUMN WIDTH]

96

Table 1. Attributes of Landsat scenes used in this study.

Epoch	Sensor	Scene ID	Date of acquisition	Path	Row
	OLI	LC80010172015227LGN00	15/08/2015	1	17
	OLI	LC80020172015234LGN00	22/08/2015	2	17
	OLI	LC80040172015216LGN00	04/08/2015	4	17
	OLI	LC80050162015255LGN00	12/09/2015	5	16
	OLI	LC80060152015214LGN00	02/08/2015	6	15
2015	OLI	LC80060162015214LGN01	02/08/2015	6	16
	OLI	LC80070132015237LGN00	25/08/2015	7	13
	OLI	LC80070142015237LGN00	25/08/2015	7	14
	OLI	LC80080122015196LGN00	15/07/2015	8	12
	OLI	LC80100102015210LGN00	29/07/2015	10	10
	OLI	LC80100112015210LGN00	29/07/2015	10	11
	ETM+	LE70020172011231EDC00	19/08/2011	2	17
	ETM+	LE70040162009207EDC00	26/07/2009	4	16
	ETM+	LE70040172009207EDC00	26/07/2009	4	17
	ETM+	LE70060152011211ASN00	30/07/2011	6	15
2010	ETM+	LE70070132010231EDC00	19/08/2010	7	13
	ETM+	LE70070142011234EDC00	22/08/2011	7	14
	ETM+	LE70090112009210EDC00	29/07/2009	9	11
	ETM+	LE70090122010229EDC00	17/08/2010	9	12
	ETM+	LE70100102009217ASN00	05/08/2009	10	10
	ETM+	LE70020172004244ASN01	31/08/2004	2	17
	ETM+	LE70040162007202EDC00	21/07/2007	4	16
	ETM+	LE70040172007202EDC00	21/07/2007	4	17
	ETM+	LE70060142007216EDC00	04/08/2007	6	14
2005	ETM+	LE70060152006245EDC00	02/09/2006	6	15
	ETM+	LE70070132005217EDC00	05/08/2005	7	13
	ETM+	LE70090112007221EDC00	09/08/2007	9	11
	ETM+	LE70090122007221EDC00	09/08/2007	9	12
	ETM+	LE70110102005229EDC00	17/08/2005	11	10
	ETM+	LE70020172000217AGS00	04/08/2000	2	17
	ETM+	LE70040161999212EDC01	31/07/1999	4	16
	ETM+	LE70040171999212EDC01	31/07/1999	4	17
	ETM+	LE70060152001215AGS00	03/08/2001	6	15
2000	ETM+	LE70070132001190EDC00	09/07/2001	7	13
2000	ETM+	LE70070142001190EDC00	09/07/2001	7	14
	ETM+	LE70090112001188EDC00	07/07/2001	9	11
	ETM+	LE70090122001188EDC00	07/07/2001	9	12
	ETM+	LE70100102000257SGS00	13/09/2000	10	10
	ETM+	LE70110102000168EDC00	16/06/2000	11	10
	TM	LT50020171992219PAC00	06/08/1992	2	17
	TM	LT50040161992217PAC00	04/08/1992	4	16
	TM	LT50040171992217PAC00	04/08/1992	4	17
	TM	LT50050161993242PAC00	30/08/1993	5	16
1992	TM	LT50060141992263PAC00	19/09/1992	6	14
	TM	LT50060151992263PAC00	19/09/1992	6	15
	TM	LT50080121994170KIS00	19/06/1994	8	12
	TM	LT50080131994170PAC00	19/06/1994	8	13
	TM	LT40090111992212XXX02	30/07/1992	9	11
	TM	LT50050151987242XXX03	30/08/1987	5	15
	TM	LT50050161987258XXX01	15/09/1987	5	16
	TM	LT50060141987201XXX08	20/07/1987	6	14
	TM	LT50060151987201XXX08	20/07/1987	6	15
1987	TM	LT50070131987176XXX01	25/06/1987	7	13
	TM	LT40080121988146XXX01	25/05/1988	8	12
	TM	LT50090111985248KIS00	05/09/1985	9	11
	TM	LT40090121988169XXX01	17/06/1988	9	12
	TM	LT50110101987236KIS00	24/08/1987	11	10

97 [SINGLE COLUMN WIDTH]

This study therefore aims to quantify changes in ice-margin extent at the terrestrial, lacustrine and marine margins of the GrIS in south-west Greenland, and to investigate how the properties of icemarginal lakes relate to rates of lacustrine ice-margin change. The objectives comprise: (i) the generation of an ice-marginal lake inventory and delineation of the ice sheet margin for 6 epochs at approximately 5-year intervals between 1987 and 2015; (ii) the quantification of ice-margin advance and recession at terrestrial, lacustrine and marine ice-margins between successive epochs; and (iii) a statistical analysis of ice-marginal lake parameters and rates of change at lacustrine margins.

106

107 2. DATA AND METHODS

108 2.1 LANDSAT SCENE SELECTION

109 A total of 58 Landsat Thematic Mapper (TM), Enhanced Thematic Mapper Plus (ETM+) and Operational 110 Land Imager (OLI) scenes were downloaded from the USGS Global Visualisation Viewer to encompass 111 the predominantly terrestrial margins of the GrIS in south-west Greenland between the mid-1980s 112 and 2015 (Figure 1, Table 1). All scenes were Level 1TP (radiometrically calibrated and orthorectified) 113 products and possessed a horizontal ground resolution of 30 m. The scenes were selected to coincide 114 with the melt season (late May to early September) in order to minimise seasonal variability and to 115 also reduce the incidence of frozen lakes and snow cover along the ice-margin. Extensive cloud and/or 116 persistent snow cover in some years necessitated a flexible sampling interval for the acquisition of 117 scenes throughout the study period. Therefore, following the method of Carrivick and Quincey (2014), 118 scenes were assigned to one of 6 epochs (1987, 1992, 2000, 2005, 2010 and 2015), with 86 % of scenes 119 acquired within ± 1 year of their respective epoch, and the remaining scenes acquired within ± 2 years 120 (Table 1). Scenes in the 2005 and 2010 epochs were also selected to mitigate the effects of the failed 121 ETM+ Scan Line Corrector (SLC) by utilising the considerable scene overlap within the study area. 122 Where SLC failure induced stripes were unavoidable, gaps were filled via mosaicing with an unaffected scene from the closest viable time period. Processing of the Landsat scenes was conducted in software
 ENVI v.5.2 and Esri ArcMap v.10.3.1.

125

126 2.2 ICE-MARGINAL LAKE INVENTORY

127 The ice-marginal lake inventory used in this study was derived by refining the 1987-2010 lake dataset 128 mapped in Carrivick and Quincey (2014) and extending the duration of the survey to incorporate 129 Landsat scenes from 2015. Details of the scene processing are fully described and evaluated in 130 Carrivick and Quincey (2014), hence a synopsis is provided here. Scenes were classified by applying 131 the Normalised Difference Water Index (NDWI) (McFeeters, 1996) to the near infrared (NIR) and blue 132 bands of the respective TM, ETM+ and OLI spectral channels, where NDWI = $((B_{NIR} - B_{Blue})/(B_{NIR} + B_{Blue}))$ 133 and B is the spectral channel. The blue, rather than the more established green, spectral channel was 134 employed because of its improved ability to discriminate water from snow and ice in cold 135 environments (Huggel et al., 2002). An upper NDWI threshold of -0.5 was used to automatically detect 136 lakes and a median filter (3×3 kernel) was used to reduce noise and remove isolated pixels. Classified 137 lakes were exported as polygons for quality assurance in ArcMap, with misclassified areas of cloud 138 and shadow manually corrected through comparison with scenes from adjacent epochs. Manual 139 digitisation was used to delineate several frozen lakes, accounting for ~0.5 % of the total lake dataset. 140 The analysis here was subsequently restricted to lakes that: (i) retained contact with the ice-margin; 141 (ii) were endorheic (with no visible outflow); and (iii) were greater than 25,000 m² in area. The ice-142 contact and endorheic conditions were included to specifically consider the effect of meltwater 143 retention on ice-margin change.

144

145 In order to establish a dataset of lake parameters, each lake was assigned a consistent identifier 146 throughout the study period by calculating the centroid of the total lake extent (the maximum outline 147 of a given lake across all epochs) (Figure 2). Lake areas were subsequently calculated within each 148 epoch, but lakes that lost ice-contact via drainage or ice-margin recession were discounted from the

dataset for the respective epoch(s). In the event of a partial lake drainage, only the lake basin that maintained ice-contact was retained in the analysis (e.g. Figure 3). In addition, each lake was assigned a persistence score (from 1-6) to indicate its permanence across the 6 epochs. The length of the interface between individual lakes and the ice-margin was measured by calculating the geometric intersection of lake polygons and the delineated ice-margin to within a tolerance of 30 m (Figure 3). Finally, the latitude and altitude of each lake centroid was extracted from a Digital Elevation Model (DEM) of the GrIS generated from 1985 aerial photography, with a ground resolution of 25 m and horizontal and vertical accuracies of ±10 m ±6 m respectively (Korsgaard et al., 2016). The delineation of lake extent was assumed to be accurate to within ±1 pixel (30 m) of the true lake perimeter. Consequently, the absolute error associated with each area measurement was dependent on lake size and planform, and thus resulted in a declining power law relationship whereby the greatest errors were associated with the smallest lakes. For example, lakes measuring 0.5 km² had an area uncertainty of ~9 %, whilst lakes measuring > 5 km² had an uncertainty of < 3 %.

- -/0



Figure 2. Example of dataset, comprising terrestrial, lacustrine and marine ice-margins, and the respective fixed points/centroids used for measurements of ice-margin change. Small circles on marine and lacustrine ice-margins represent the vertices over which distance measurements are averaged. Total lake extent represents the maximum outline of each lake across all epochs. Basemap: 2015 false-colour Landsat OLI scene.

- 183 [DOUBLE COLUMN WIDTH]
- 184
- 185
- 186
- 187
- 188





Figure 3. Example of temporal variation in lake area and lake – ice-margin intersect over the survey period. In particular note the partial lake drainage between 2000 and 2005, and subsequent refilling. [DOUBLE COLUMN WIDTH]

197 The ice sheet margin in south-west Greenland was delineated by using the green and shortwave 198 infrared (SWIR) bands of the respective TM, ETM+ and OLI spectral channels to classify scenes with 199 the Normalised Difference Snow Index (NDSI) (Hall et al., 1995), where NDSI = 200 $((B_{Green} - B_{SWIR})/(B_{Green} + B_{SWIR}))$. An NDSI threshold of 0.45 ± 0.1 was used to classify areas of snow and 201 ice on a scene by scene basis with the aim of minimising subsequent manual post-processing. Due to 202 the similar spectral properties of snow, ice and water in the green and SWIR spectral bands, an 203 additional threshold of 0.45 was applied in the respective NIR band to mask out water bodies in the ice-marginal environment and thus improve the accuracy of ice-margin delineation. A median filter 204 205 (3x3 kernel) was also applied to reduce noise and remove small snow patches. Manual editing was 206 subsequently employed to refine the delineated ice-margins in isolated areas affected by shadow, 207 debris cover and late-lying snow. Consistent mapping of the ice-margin was achieved in these regions 208 through consultation with scenes from neighbouring epochs and high resolution DigitalGlobe imagery 209 in Google Earth. Finally, delineated ice-margins in adjacent scenes were merged to generate a single 210 ice-margin for south-west Greenland within each epoch (Figures 1, 3), which were subsequently used 211 to derive the measurements of ice-margin advance and recession detailed in Section 2.4. 212 Measurements of total ice-margin length, comparable to the length of the lake – ice-margin intersects 213 detailed in Section 2.2, were generated by smoothing the delineated ice-margins to a tolerance of 214 30 m. Given that all results presented here are regionally aggregated, it is assumed that any over-215 estimation of the ice-margin position is cancelled out by an equal and opposite under-estimation, and 216 uncertainty in the ice-margin positions is therefore not specifically assessed for these bulk figures.

217

218 2.4 MEASUREMENTS OF ICE-MARGIN CHANGE

Rates of ice-margin advance and recession between successive epochs were calculated by measuring
changes in ice-margin position relative to a series of fixed reference points across the study period.
Existing techniques for measuring changes in glacier extent have been primarily developed to quantify
changes in the position of glacier termini occupying troughs (e.g. Lea et al., 2014), and are thus

223 unsuited to analysing changes at lacustrine margins which typically occupy a greater diversity of ice-224 marginal environments, particularly the lateral margins of outlet glaciers (cf. Figure 2). For example, 225 techniques that measure change along the centre-line of the glacier (e.g. Bevan et al., 2012; Mernild 226 et al., 2012) are not applicable at the majority of lacustrine margins. In addition, the highly dynamic 227 nature of many lake – ice-margin interfaces (cf. Figure 3) prevents the use of fixed boxes to calculate 228 area averaged advance or recession between successive epochs (e.g. Howat and Eddy, 2011; Hill et 229 al., 2018). Therefore, changes in the extent of lacustrine and marine ice-margins here were measured 230 using the bow method outlined in Bjørk et al. (2012). The centroids of the total lake extents were used 231 as fixed reference points from which to measure distances to the respective lake – ice-margin intersect 232 within each epoch, with a series of points established on the vertices of each intersect to permit the 233 calculation of a mean lake centroid – intersect distance (Figure 2). Rates of ice-margin advance or 234 recession were then calculated by differencing the mean distance values in successive epochs and 235 dividing by the interval duration. Consequently, rates of change at lacustrine margins were only 236 generated when a lake was present in two or more successive epochs. In the rare instances in which 237 a lake - ice-margin intersect was manifest in multiple sections (e.g. Figure 3) the loss/addition of 238 intersect sections between successive epochs resulted in small under- and over-estimates of ice-239 margin recession respectively, which collectively had a negligible effect on the aggregated measures 240 of lacustrine margin change. Changes in the extent of marine margins were measured in the same 241 manner as lacustrine margins by establishing fixed marine points in front of each terminus and 242 calculating changes in mean distance between the marine points and respective marine margin 243 vertices in successive epochs (Figure 2). Changes at terrestrial margins were calculated by creating a series of fixed points at 1 km intervals along a 250 m buffer of the delineated 1992 ice-margin. 244 245 Distances between the fixed terrestrial points and the proximal point on the terrestrial margin were 246 measured using proximity analysis and subsequently differenced to calculate rates of advance and 247 recession between successive epochs.

248

249 2.5 STATISTICAL ANALYSES

250 Data were analysed using multivariate regression methods in R v.3.6.0 (R Core Team, 2019) to: (i) 251 investigate differences in rates of change at terrestrial, lacustrine and marine ice-margins; and (ii) 252 assess the influence of lake parameters on rates of change at lacustrine margins. Two linear mixed-253 effects models (LMMs) were fitted (Bates et al., 2015) using the rate of ice-margin change as the 254 dependent variable in both models. One data point from Jakobshavn Isbrae was omitted due to its 255 extreme outlying status (> 2 km recession between the 2000 and 2005 epochs). The repeated sampling of the same sites across epochs was accounted for by including location as a random effect. 256 257 LMM 1 compared rates of change at the disparate margin types, and included: ice-margin type 258 (lacustrine, marine, terrestrial); epoch; and latitude as independent variables (with latitudinal data 259 included to control for the spatial clustering of particular margin types along the ice-margin). LMM 2 assessed rates of change at lacustrine margins, and included the independent variables: latitude; 260 261 altitude; lake area; intersect length; persistence; and epoch. All independent variables were tested for 262 multicollinearity prior to model fitting; however, lake area and intersect length failed to meet this 263 assumption (|r| > 0.7; Dormann et al., 2013). Consequently, two alternate versions of LMM 2 were 264 fitted to accommodate lake area and intersect length respectively.

265

266 **3. RESULTS**

267 3.1 MODEL FIT

Testing of the fitted LMMs for normality and heteroscedasticity revealed that the distribution of the residuals was heavy-tailed. Consequently, additional Robust LMMs were constructed to assess the impact of outliers on model fit (Koller, 2006). The resultant similarity of the respective LMM and Robust LMM coefficients (Table 2) indicated that the outliers had a limited effect on the fit of the models, therefore the outputs of the initial LMMs are presented henceforth.

273

LMM 1* LMM 2(a) LMM 2(b) Robust Robust Robust Ind. variables Ind. variables LMM Ind. variables LMM LMM LMM LMM LMM -0.120 -0.081 0.051 0.068 Epoch Latitude 0.065 Latitude 0.049 Altitude Type:Marine -2.637 -0.612 0.132 0.074 Altitude 0.132 0.073 Type:Terrestrial 0.215 0.100 -0.191 -0.074 Intersect length -0.223 -0.095 Lake area 0.012 Latitude 0.020 Epoch -0.157 -0.142 Epoch -0.153 -0.142 Persistence 0.030 0.021 Persistence 0.094 0.049

276 Table 2. LMM and Robust LMM model variables and coefficients.

277 * Note reference values for categorical variables 'Type' in LMM 1 are Lacustrine.

278 [DOUBLE COLUMN WIDTH]

279

275

280 3.2 ICE-MARGIN CHANGE AT TERRESTRIAL, LACUSTRINE AND MARINE MARGINS

The ice-margins mapped in this study delineate a ~5000 km length of the south-western margin of the 281 282 GrIS. Cumulative totals of ice-margin type remained broadly consistent between 1987 and 2015, with 283 ~89 % of the ice-margin in the study area terminating in a terrestrial setting, ~8 % in a lacustrine setting and ~3 % in a marine setting (Table 3). The number of measurements of ice-margin change between 284 285 successive epochs was substantial throughout the study period, with each period incorporating 286 measurements from between 22 to 35 marine margins, 353 to 439 lacustrine margins and 2469 to 287 3325 terrestrial margins (Table 4). From 1992 onwards, mean change at all margin types was negative, 288 signifying an extended duration of ice-margin recession in south-west Greenland. However, positive 289 values of mean change between 1987 and 1992 reveal an earlier period of ice-margin advance at both 290 terrestrial and marine margins, although mean change at lacustrine margins remained negative 291 (Table 4).

292

293

294

295

297 Table 3. Summary statistics of ice-margin composition in south-west Greenland throughout the study

298	period.	

Epoch	Total i	Total ice-margin length		Terrestrial ice- margin length		Lacustrine ice- margin length		Marine ice-margin length	
	km	% of total	km	% of total	km	% of total	km	% of total	
1987	3722*	100.00	3306	88.82	319	8.56	98	2.63	
1992	5029	100.00	4500	89.47	398	7.91	132	2.62	
2000	5019	100.00	4466	88.98	421	8.38	132	2.64	
2005	4916	100.00	4377	89.04	405	8.24	133	2.71	
2010	4966	100.00	4429	89.19	402	8.10	135	2.71	
2015	4932	100.00	4345	88.10	434	8.80	153	3.10	

* The reduced length of the 1987 ice-margin is due to the unavailability of Landsat TM scenes from the southern end of the study

300 area in the years 1985-1988.

301 [DOUBLE COLUMN WIDTH]

302

Table 4. Summary statistics of ice-margin change in south-west Greenland throughout the study

304 period. Note positive and negative values represent ice-margin advance and recession respectively.

	Terr	estrial ice-ma	Irgins Lacustrine ice-margins			argins	Marine ice-margins			
Period	n	Mean change (m)	Mean annual change (m)	n	Mean change (m)	Mean annual change (m)	n	Mean change (m)	Mean annual change (m)	
1987-1992	2469	5.8	1.2	353	-5.3	-1.1	22	96.3	19.3	
1992-2000	3325	-1.4	-0.2	439	-13.0	-1.6	35	-250.6	-31.3	
2000-2005	3325	-24.0	-4.8	414	-28.3	-5.7	35	-640.3	-69.7	
2005-2010	3325	-15.3	-3.1	401	-32.2	-6.4	35	-197.6	-39.5	
2010-2015	3325	-13.8	-2.8	374	-57.3	-11.5	35	-417.8	-83.6	

305 [DOUBLE COLUMN WIDTH]

306

LMM 1 identified a significant negative correlation between rate of ice-margin change and epoch (p < 0.001), signifying increasing rates of ice-margin recession in south-west Greenland between 1987 and 2015, irrespective of margin type (Table 5). The model also identified significant differences between rates of change at lacustrine and marine margins (p < 0.001), and lacustrine and terrestrial margins (p < 0.001) (Table 5). Marine margins exhibited both the greatest mean rates of ice-margin recession and the greatest variability in frontal behaviour throughout the study period, with rates of advance and recession at several termini exceeding 100 m per year (Table 4, Figure 4a). The magnitude and variability of changes at terrestrial and lacustrine margins were more comparable, although changes at lacustrine margins were less clustered around the median and typically more negative than their terrestrial counterparts (Table 4, Figure 4b). Notably, although rates of recession increased at both terrestrial and lacustrine ice-margins between 1987 and 2015, recession at lacustrine margins increasingly outpaced that of terrestrial margins throughout the survey period (Figure 5).

319

320	Table 5. LMM results.	Significant relation	ationships are	highlighted in bold.
-----	-----------------------	----------------------	----------------	----------------------

LMM	Ind. Variables	Estimate	Std. Error	t value	95 % Confidence intervals		p value
110.					lower	upper	
	(Intercept)	-0.167	0.024	-6.975	-0.214	-0.120	< 0.001
	Epoch	-0.120	0.007	-17.360	-0.134	-0.107	< 0.001
1	Type:Marine*	-2.637	0.088	-29.907	-2.809	-2.464	< 0.001
	Type:Terrestrial*	0.215	0.025	8.466	0.165	0.265	< 0.001
	Latitude	0.020	0.008	2.453	0.004	0.036	0.014
	(Intercept)	-0.039	0.040	-0.986	-0.117	0.039	0.324
	Latitude	0.065	0.026	2.438	0.013	0.116	0.015
2(0)	Altitude	0.132	0.025	5.241	0.083	0.181	< 0.001
2(a)	Area	-0.191	0.022	-8.687	-0.234	-0.148	< 0.001
	Epoch	-0.157	0.022	-7.215	-0.200	-0.114	< 0.001
	Persistence	0.030	0.103	0.293	-0.171	0.231	0.770
	(Intercept)	-0.072	0.040	-1.796	-0.150	0.006	0.073
	Latitude	0.068	0.026	2.576	0.016	0.119	0.010
2 (b)	Altitude	0.132	0.025	5.299	0.084	0.181	< 0.001
2(D)	Intersect length	-0.223	0.022	-10.026	-0.267	-0.180	< 0.001
	Epoch	-0.153	0.022	-7.085	-0.196	-0.111	< 0.001
	Persistence	0.094	0.103	0.917	-0.107	0.295	0.359

321 * Note reference values for categorical variables 'Type' are lacustrine.

322 [DOUBLE COLUMN WIDTH]

323

324

326

325

327



Figure 4. Box plots of ice-margin change throughout the study period at: (a) terrestrial, lacustrine and marine margins; and (b) terrestrial and lacustrine margins only. Note positive and negative values of annual change represent ice-margin advance and recession respectively. To improve clarity, 7 and 21 outlying data points have been cropped from panels (a) and (b) respectively.

334 [DOUBLE COLUMN WIDTH]



Figure 5. Linear regression of annual change and year, showing trends in ice-margin recession at terrestrial and lacustrine margins. The final year of each epoch has been used to plot the linear relationship. Grey shading represents the 95 % confidence interval. Individual data points have been removed to improve clarity (terrestrial n = 15769; lacustrine n = 1981).

348 [DOUBLE COLUMN WIDTH]

349

350 3.3 CONTROLS ON LACUSTRINE ICE-MARGIN CHANGE

Both variants of LMM 2 identified a number of significant relationships between lake parameters and 351 352 rates of change at lacustrine margins (Table 5). The significant positive correlation between latitude 353 and rate of change (p < 0.05) indicates that ice-margin recession was accentuated at lower latitudes 354 (Figure 6a). Altitude was also found to act as a control on ice-margin change (p < 0.001), with increased 355 rates of recession at lower altitudes (Figure 6b). Both lake area (p < 0.001) and intersect length 356 (p < 0.001) possessed a similar significant negative correlation with rate of change, demonstrating that 357 increased rates of ice-margin recession are associated with larger lakes and longer lake – ice-margin interfaces (Figures 6c, 6d). Finally, there was a significant negative correlation between epoch and 358 359 rate of change (p < 0.001), signifying that rates of recession at lacustrine margins increased

360 throughout the duration of the study (Figure 6e). No significant correlation was found between



362



Figure 6. Linear regressions of annual change at lacustrine margins and lake parameters, comprising: (a) latitude; (b) altitude; (c) lake area; (d) intersect length; and (e) year. Note the final year of each epoch has been used to plot the linear relationship in (e). Grey shading represents the 95 % confidence interval. Individual data points have been removed to improve clarity (n = 1981).

368 [DOUBLE COLUMN WIDTH]

369

370 **4. DISCUSSION**

371 4.1 ICE-MARGIN CHANGE IN SOUTH-WEST GREENLAND

372 The temporal patterns of ice-margin advance and recession that we record in south-west Greenland 373 broadly reflect previously-documented changes in ice sheet mass balance. Notably, in contrast to the 374 mean ice-margin recession observed at all margin types post-1992, a mean advance at terrestrial and 375 marine margins, and concurrent minima of lacustrine margin recession, is evident between 1987 and 376 1992 (Table 4). Although this distinction could be accentuated by the unavailability of satellite imagery 377 from the southern reaches of the survey area in 1987 (Table 3), this pattern coincides with existing 378 records of outlet glacier advance and general ice sheet expansion in the southern and western GrIS 379 during the 1980s (e.g. Weidick, 1991; Zwally, 1989; Van Tatenhove et al., 1995; Knight et al., 2000) 380 following a period of net mass gain in the preceding decade (Mouginot et al., 2019). Similarly, the 381 period of ice-margin recession recorded post-1992 occurs following a transition to net mass loss at 382 the GrIS in the 1980s (Mouginot et al., 2019), and is sustained through a further fivefold increase in 383 the rate of mass loss between the 1990s and 2010 (Shepherd et al., 2020). However, the statistically 384 significant differences between rates of change at the disparate margin types (Table 5) and 385 considerable differences in variability of frontal behaviour (Figure 4) are indicative of heterogeneous 386 responses at the respective terrestrial, lacustrine and marine margins of the GrIS, despite undergoing 387 comparable climatic forcing over the survey duration.

388

The relative magnitude and variability of the changes recorded at the disparate ice-marginal environments in this study are similar to those observed at the western margins of the GrIS in the mid-20th century by Warren (1991), in which marine margins were found to exhibit the greatest magnitude and variability in frontal behaviour, and terrestrial margins the least. Because terrestrial termini lack oceanic or lacustrine forcing, changes in ice-margin extent are typically a delayed response to regional climatic forcing, with inter-glacier variability arising from glacier-specific factors, including: glacier geometry; hypsometry; debris-cover; and local climatic conditions (e.g. Pelto and Hedlund, 2001; Scherler et al., 2011; Davies et al., 2012; Sakai and Fujita, 2017; Lovell et al., 2019). Consequently, the
relatively limited variability in frontal behaviour at the terrestrial margins of the GrIS, in comparison
to its marine margins, was expected (Figure 4a), and has been similarly observed in analyses of outlet
glacier and PGIC extent in south-eastern Greenland over the same period (e.g. Mernild et al., 2012).
Furthermore, the low magnitude of the changes observed at terrestrial margins compares favourably
with existing records of terrestrial frontal behaviour in western Greenland in the 1990s and 2000s (e.g.
Moon and Joughin, 2008; Carr et al., 2013).

403

404 In comparison to terrestrial margins, marine termini demonstrated considerably greater magnitude 405 and variability in frontal behaviour over the duration of the study, with a mean annual advance of 19 ma⁻¹ between 1987 and 1992, succeeded by mean annual recessions exceeding 31 ma⁻¹ in all 406 remaining periods (Table 4). In addition, the mean changes masked considerable complexity in the 407 408 behaviour of individual marine terminating glaciers, with advances and recessions in the order of 10s 409 and 100s of ma⁻¹ respectively becoming increasingly prevalent from 2000-2005 onwards (Figure 4a). Isolating the exact drivers of change at marine ice-margins is challenging due to the complexities and 410 411 interactions of both atmospheric and oceanic forcings, as well as glacier-specific controls including 412 terminus geometry and bathymetry (McFadden et al., 2011; Porter et al., 2018). However, oceanic 413 forcing is increasingly recognised as a key control on the dynamics of the marine outlets of the GrIS 414 (Seale et al., 2011; Straneo and Heimbach, 2013). Accordingly, the observed transition from mean 415 terminus advance to mean terminus recession at marine margins in 1992-2000 coincides with 416 recorded increases in subsurface ocean temperatures along the west coast of Greenland in the mid-417 1990s (Myers et al., 2007; Holland et al., 2008), which are hypothesised to have triggered the collapse 418 of several floating termini and a subsequent phase of regional marine ice-margin recession in response 419 to debuttressing (e.g. Joughin et al., 2012). A further increase in marine ice-margin recession observed 420 in this study in the early 2000s also concurs with similar observations from the same time period at

421 marine terminating outlets in south-eastern Greenland (Mernild et al., 2012) and across the wider ice
422 sheet (Moon and Joughin, 2008; Howat and Eddy, 2011).

423

424 Despite mean changes in marine ice-margin extent being approximately an order of magnitude 425 greater than those recorded at terrestrial and lacustrine ice-margins between 1987 and 2015, caution 426 is necessary when interpreting and comparing the frontal behaviour of the disparate ice-marginal 427 environments of the GrIS. In particular, marine ice-margins constitute by far the smallest component 428 of the overall dataset (n = 22-35), and typically exhibit seasonal variations in terminus advance and 429 recession that can be challenging to control for using multi-annual snapshots of terminus position 430 (Schild and Hamilton, 2013). Furthermore, the relatively similar magnitudes of change observed at 431 terrestrial and lacustrine termini conceal a notable divergence in the behaviour of the respective margin types over the duration of the study. In particular, the persistently negative values of mean 432 433 changes at lacustrine margins, coupled with a progressive increase in their magnitude and their 434 increased outpacing of change at terrestrial margins (Table 4, Figure 5), could be indicative of 435 amplified lacustrine forcing and mass loss at the lake terminating margins of the GrIS between 1987 436 and 2015.

437

438 4.2 LACUSTRINE ICE-MARGIN RECESSION

439 The observed dissimilarities in the frontal behaviour of the terrestrial and lake terminating margins of 440 the GrIS over the course of the study can be explained by the impact of lacustrine forcing on ice-margin 441 dynamics. In particular, lake formation has significant implications for processes and rates of mass loss 442 at ice-margins through the onset of both calving (Kirkbride, 1993; Motyka et al., 2003) and subaqueous 443 melt (Eijpen et al., 2003; Haresign and Warren, 2005; Truffer and Motyka, 2016). Furthermore, ice-444 marginal lake formation can destabilise and perturb wider ice-margin dynamics through the initiation 445 of a positive feedback whereby enhanced rates of mass loss increase local ice-surface gradients, thus 446 promoting acceleration, thinning and fracture of the ice-margin, which in turn creates favourable

conditions for amplified calving losses (Benn et al., 2007; Carrivick and Tweed, 2013). This feedback
has been invoked as the cause of the rapid ice-margin recession observed at an increasing number of
alpine glaciers (Naruse and Skvarca, 2000; Boyce et al., 2007; Basnett et al., 2013; Trussel et al., 2013;
King et al., 2018; Liu et al., 2020).

451

452 The correlations between rate of ice-margin change and lake area and intersect length respectively 453 (Figure 6c-d, Table 5), suggest that lake size exerts a control on rates of mass loss at lacustrine margins. 454 Although the augmented rates of ice-margin recession at larger lakes can be hypothesised to arise 455 from the combined effects of calving and subaqueous melt occurring over a greater length of the ice-456 margin, it is likely that the greater water depths typically associated with larger lakes (e.g. Huggel et 457 al., 2002; Cook and Quincey, 2015) are also a key driver of ice-margin recession. In particular, several 458 empirical relationships have linked increased calving rates to greater lake depths (Warren et al., 1995; 459 Warren and Kirkbride, 2003), and accelerated rates of lacustrine ice-margin recession have been 460 observed following the retreat of termini into glacial overdeepenings (e.g. Kirkbride, 1993; Boyce et 461 al., 2007; Larsen et al., 2015). In addition, the increased buoyancy and reduced effective pressure 462 apparent at ice-margins terminating in deeper water favours the positive feedback between mass loss 463 and terminus recession.

464

465 Increases in lake area and depth could also explain the progressive growth in mean annual lacustrine margin recession rates by an order of magnitude throughout the study, from 1.1 ma⁻¹ between 1987 466 and 1992 to 11.5 ma⁻¹ between 2010 and 2015 (Table 4, Figure 6e). A behavioural analysis of the lake 467 468 dataset in Carrivick and Quincey (2014) revealed that ~45 % of all ice-marginal lakes in south-west 469 Greenland formed or increased in size between 1987 and 2010, in contrast to only ~30 % of lakes 470 decreasing in size or draining over the same period. Furthermore, the inverse bed slope along much 471 of the ice sheet margin in south-west Greenland creates favourable conditions for ongoing lake 472 expansion in response to ice-margin recession (Carrivick et al., 2017a; Morlighem et al., 2017). Lake

473 persistence was the only independent variable in the LMMs that did not significantly correlate with 474 the rate of ice-margin change at lacustrine termini (Table 5), which may be indicative of a multifaceted 475 relationship between ice-margin recession and lake stability. For example, although the most 476 persistent lakes may be associated with greater rates of ice-margin recession due to their extended 477 prevalence, lakes in contact with rapidly retreating ice-margins may also be inherently less stable due 478 to increased opportunities for lake drainage through failure of the ice-dam or rapid changes to lake 479 morphometry (e.g. Russell et al., 2011; Carrivick et al., 2017b; Carrivick and Tweed, 2019).

480

481 An additional cause of the enhanced rates of recession at lacustrine margins observed over the 482 duration of this survey could be the lengthening of the season over which lacustrine processes, 483 including subaqueous melt and calving, were able to promote mass loss. For example, analyses of non 484 ice-contact lakes in the Arctic have identified an earlier break-up of winter ice-cover and an increase 485 in ice-free days in response to atmospheric warming over recent decades (Duguay et al., 2006; 486 Smejkalova et al., 2016; Surdu et al., 2016). Similar changes to the ice-cover regimes of ice-marginal 487 lakes in south-west Greenland could therefore have amplified ice-margin recession through the 488 prolonged operation of lacustrine processes associated with higher rates of mass loss, such as melt-489 undercutting (e.g. Mallalieu et al., 2020). These processes are likely to be further accentuated in lakes 490 with a reduced duration of ice-cover by enhanced lake temperatures arising from the low albedo of 491 open water. Notably, the mean annual lacustrine margin recession rates measured in this study 492 increased following the switch to a negative phase of the North Atlantic Oscillation in the mid-1990s 493 (Table 4), which is typically associated with enhanced summertime warming in west Greenland (Hanna 494 et al., 2008; Bevis et al., 2019). In addition, further evidence of climatic control on rates of lacustrine 495 margin recession is provided by the significant positive correlations between ice-margin change and 496 latitude and altitude respectively (Figure 6a-b, Table 5), which highlight a strong association between high rates of lacustrine recession and the warmer climatic conditions typically associated with lower 497 498 latitudes and altitudes. Consequently, the relationships identified here between latitude, altitude and

rates of lacustrine margin change, could be considered as tentative indicators of the future response
of lacustrine margins to anticipated increases in atmospheric forcing in western Greenland (Bevis et
al., 2019).

502

503 4.3 IMPLICATIONS AND FUTURE RESEARCH

504 In addition to enhancing local rates of ice-margin recession, the presence of lakes at the margin of the 505 GrIS could have profound implications for wider ice sheet dynamics and stability. For example, Price 506 et al. (2008) demonstrated that dynamic changes at the margins of the GrIS can propagate dozens of 507 kilometres up-ice via longitudinal coupling. Therefore enhanced recession at lacustrine margins and 508 resultant increases in surface gradients have significant potential to amplify surface velocities and 509 promote dynamic thinning up-ice of the lacustrine termini, particularly where lakes are large and deep 510 relative to the thickness of the ice-margin. Similar responses to lake formation and growth have been 511 extensively documented in the Himalaya, where lacustrine terminating glaciers account for an 512 increasingly disproportionate share of regional mass loss (Basnett et al., 2013; King et al., 2018; Brun 513 et al., 2019; King et al., 2019). Currently 434 km (~9 %) of the ice sheet margin in south-west Greenland 514 terminates in a lacustrine setting, in contrast to 153 km (~3 %) in a marine setting (Table 3). However, 515 thinning at the margins of the ice-sheet (Krabill et al., 2004; Pritchard et al., 2009), coupled with 516 continued atmospheric warming (Pattyn et al., 2018; Bevis et al., 2019) and the recession of the ice-517 margin over an inverse bed slope (Carrivick et al., 2017a; Morlighem et al., 2017), will create 518 favourable conditions for enhanced ice-marginal lake formation and growth in south-west Greenland 519 in coming decades. Additionally, over longer timescales, the recession of marine termini onto land 520 (e.g. Joughin et al., 2010; Nick et al. 2013) will further increase the potential for lake formation at the 521 ice sheet margin. Consequently, it can be hypothesised that ice-marginal lakes will play an increasingly 522 important role in rates and patterns of deglaciation in Greenland, and that continued lake expansion 523 will amplify future mass loss from the south-western margin of the GrIS. Furthermore, inadequate 524 consideration of the impacts of lacustrine forcing at the margin of the GrIS could lead to increasing

error in projections of the ice sheet's response to climate change, and its contribution to sea level rise.
The inclusion and parameterisation of lake – ice-margin interactions in numerical ice sheet models is
therefore increasingly desirable (Carrivick et al., 2020).

528

529 A more advanced understanding of the impact of lacustrine forcing on the margin of the GrIS could be 530 developed by focusing future research efforts in three main areas. Firstly, sections of the ice-margin 531 susceptible to lake formation and growth could be determined through the development of 532 morphometric and dynamic criteria, similar to those employed to forecast ice-marginal lake formation 533 in the Himalaya (Reynolds, 2000; Quincey et al., 2007), particularly if integrated with recent high-534 resolution mapping of GrIS bed topography (e.g. Morlighem et al., 2017). In addition, knowledge of 535 basal topography and ice thickness can facilitate predictions of lake area and depth, which are 536 significant controls on rates of recession at lacustrine margins (Figure 6c). Secondly, regional-scale 537 analyses of changes in velocity, structure and ice-surface elevation up-ice of the lacustrine termini of 538 the GrIS are necessary to determine the magnitude of the dynamic response of the ice-sheet to lake 539 formation and thus refine estimates of mass loss and sea level rise from lacustrine margins. This 540 objective, and the extent to which the observations and conclusions drawn from this study in south-541 west Greenland may be applicable to the wider ice sheet, will be greatly facilitated by the recent 542 generation of the first Greenland-wide multi-sensor inventory of ice-marginal lakes in How et al. 543 (2021). Finally, local-scale analyses of lacustrine ice-margin dynamics are required to improve 544 knowledge of the mechanisms driving enhanced ice-margin recession. In particular, calving processes 545 and rates of subaqueous melt remain relatively poorly constrained at lacustrine ice-margins (Haresign 546 and Warren, 2005; Trussel et al., 2013; Purdie et al., 2016; Truffer and Motyka, 2016; Mallalieu et al., 2020). 547

548

549 5. CONCLUSIONS

550 This study has presented the first systematic analysis of changes in the extent of the terrestrial, 551 lacustrine and marine margins of the GrIS in south-west Greenland between 1987 and 2015. The 552 analysis revealed an extended and accelerating phase of ice-margin recession in south-west Greenland 553 from 1992 onwards, irrespective of ice-margin type. However, significant differences in rates of ice-554 margin change also indicated a heterogeneous response at the respective ice-marginal environments 555 of the GrIS to comparable climatic forcing over the survey duration. Marine terminating ice-margins 556 exhibited the greatest magnitude and variability in ice-margin change, with rapid ice-margin recession 557 becoming pervasive from 1992. Mean ice-margin recession rates and variability in frontal behaviour 558 were also consistently greater at lacustrine termini than their terrestrial counterparts. In addition, 559 mean ice-margin recession rates at lacustrine termini increased by an order of magnitude over the 560 duration of the survey and progressively outpaced those measured at terrestrial ice-margins. This 561 study has also identified significant correlations between rates of lacustrine ice-margin recession and 562 lake parameters, including lake area, latitude, altitude and the length of the lake - ice-margin 563 interface. The progressive increase in rates of lacustrine ice-margin recession measured over the 564 duration of the survey are theorised to have arisen from increases in lake size and a lengthening of 565 the season in which calving and subaqueous melt processes can promote mass loss at lacustrine 566 termini. These results suggest that ice-marginal lakes have become increasingly significant drivers of 567 ice-margin recession and thus mass loss at the GrIS, and are likely to further increase in importance in 568 response to enhanced ice-marginal lake prevalence in coming decades. Further research is therefore 569 necessary to better parameterise the causal connections between ice-marginal lake evolution and 570 enhanced ice-margin recession in Greenland, and thus refine the contribution of mass loss from the 571 lacustrine margins of the GrIS to sea level rise projections.

572

573 ACKNOWLEDGEMENTS

JM was funded by a Graduate Assistantship from the School of Geography at the University of Leeds.
CR was the recipient of a NERC Case Award (grant ref. NE/K007599/1). The authors would like to thank

- two anonymous reviewers for their valuable comments and insights which helped to enhance this
- 577 manuscript.
- 578

579 DATA STATEMENT

- 580 The dataset of ice-margin change used in this analysis is available from the UK Polar Data Centre:
- 581 <u>https://doi.org/</u>... The R code for the statistical analyses is available from GitHub: <u>https://doi.org/</u>...
- 582 (DOIs will be confirmed prior to publication).
- 583

584 **REFERENCES**

- Aschwanden, A., Fahnestock, M.A., Truffer, M., Brinkerhoff, D.J., Hock, R., Khroulev, C., Mottram, R.
 and Khan, S.A. 2019. Contribution of the Greenland Ice Sheet to sea level over the next millennium. *Science Advances.* 5(6), eaav9396.
- 588 Basnett, S., Kulkarni, A.V. and Bolch, T. 2013. The influence of debris cover and glacial lakes on the 589 recession of glaciers in Sikkim Himalaya, India. *Journal of Glaciology*. **59**(218), pp.1035-1046.
- Bates, D., Machler, M., Bolker, B.M. and Walker, S.C. 2015. Fitting Linear Mixed-Effects Models Using
 Ime4. *Journal of Statistical Software*. 67(1), pp.1-48.
- 592 Benn, D.I., Warren, C.R. and Mottram, R.H. 2007. Calving processes and the dynamics of calving 593 glaciers. *Earth-Science Reviews*. **82**(3-4), pp.143-179.
- Bevan, S.L., Luckman, A.J. and Murray, T. 2012. Glacier dynamics over the last quarter of a century at
 Helheim, Kangerdlugssuaq and 14 other major Greenland outlet glaciers. *The Cryosphere*. 6(5),
 pp.923-937.
- Bevis, M., Harig, C., Khan, S.A., Brown, A., Simons, F.J., Willis, M., Fettweis, X., van den Broeke, M.R.,
 Madsen, F.B., Kendrick, E., Caccamise, D.J., van Dam, T., Knudsen, P. and Nylen, T. 2019. Accelerating
 changes in ice mass within Greenland, and the ice sheet's sensitivity to atmospheric forcing. *Proceedings of the National Academy of Sciences.* 116(6), pp.1934-1939.
- Bjørk, A.A., Aagaard, S., Lutt, A., Khan, S.A., Box, J.E., Kjeldsen, K.K., Larsen, N.K., Korsgaard, N.J.,
 Cappelen, J., Colgan, W.T., Machguth, H., Andresen, C.S., Peings, Y. and Kjaer, K.H. 2018. Changes in
 Greenland's peripheral glaciers linked to the North Atlantic Oscillation. *Nature Climate Change.* 8(1),
 pp.48-52.
- Bjørk, A.A., Kjær, K.H., Korsgaard, N.J., Khan, S.A., Kjeldsen, K.K., Andresen, C.S., Box, J.E., Larsen, N.K.
 and Funder, S. 2012. An aerial view of 80 years of climate-related glacier fluctuations in southeast
 Greenland. *Nature Geoscience*. 5(6), pp.427-432.
- 608 Boyce, E.S., Motyka, R.J. and Truffer, M. 2007. Flotation and retreat of a lake-calving terminus, 609 Mendenhall Glacier, Southeast Alaska, USA. *Journal of Glaciology*. **53**(181), pp.211-224.
- Brun, F., Wagnon, P., Berthier, E., Jomelli, V., Maharjan, S.B., Shrestha, F. and Kraaijenbrink, P.D.A.
- 611 2019. Heterogeneous Influence of Glacier Morphology on the Mass Balance Variability in High
- 612 Mountain Asia. *Journal of Geophysical Research: Earth Surface*. **124**(6), pp.1331-1345.

- 613 Carr, J.R., Stokes, C.R. and Vieli, A. 2013. Recent progress in understanding marine-terminating Arctic
- outlet glacier response to climatic and oceanic forcing: Twenty years of rapid change. *Progress in Physical Geography: Earth and Environment.* **37**(4), pp.436-467.
- 616 Carrivick, J.L. and Quincey, D.J. 2014. Progressive increase in number and volume of ice-marginal lakes 617 on the western margin of the Greenland Ice Sheet. *Global and Planetary Change*. **116**, pp.156-163.
- 618 Carrivick, J.L. and Tweed, F.S. 2013. Proglacial lakes: character, behaviour and geological importance.
 619 *Quaternary Science Reviews.* 78, pp.34-52.
- 620 Carrivick, J.L. and Tweed, F.S. 2019. A review of glacier outburst floods in Iceland and Greenland with 621 a megafloods perspective. *Earth-Science Reviews*. **196**, 102876.
- 622 Carrivick, J.L., Tweed, F.S., Ng, F., Quincey, D.J., Mallalieu, J., Ingeman-Nielsen, T., Mikkelsen, A.B.,
- Palmer, S.J., Yde, J.C., Homer, R. and Russell, A.J. 2017b. Ice-dammed lake drainage evolution at Russell
 Glacier, West Greenland. *Frontiers in Earth Science*. 5, 2017.00100
- 625 Carrivick, J.L., Tweed, F.S., Sutherland, J.L. and Mallalieu, J. 2020. Toward numerical modeling of 626 interactions between ice-marginal proglacial lakes and glaciers. *Frontiers in Earth Science*. **8**, 627 2020.577068.
- 628 Carrivick, J.L., Yde, J., Russell, A.J., Quincey, D.J., Ingeman-Nielsen, T. and Mallalieu, J. 2017a. Ice-629 margin and meltwater dynamics during the mid-Holocene in the Kangerlussuaq area of west 630 Greenland. *Boreas.* **46**(3), pp.369-387.
- 631 Catania, G.A., Stearns, L.A., Sutherland, D.A., Fried, M.J., Bartholomaus, T.C., Morlighem, M., Shroyer,
- E. and Nash, J. 2018. Geometric Controls on Tidewater Glacier Retreat in Central Western Greenland. *Journal of Geophysical Research-Earth Surface.* 123(8), pp.2024-2038.
- 634 Citterio, M., Paul, F., Ahlstrom, A.P., Jepsen, H.F. and Weidick, A. 2009. Remote sensing of glacier
 635 change in West Greenland: accounting for the occurrence of surge-type glaciers. *Annals of Glaciology*.
 636 **50**(53), pp.70-80.
- 637 Cook, S.J. and Quincey, D.J. 2015. Estimating the volume of Alpine glacial lakes. *Earth Surface.*638 *Dynamics*. **3**, pp.559-575.
- Davies, B.J., Carrivick, J.L., Glasser, N.F., Hambrey, M.J. and Smellie, J.L. 2012. Variable glacier response
 to atmospheric warming, northern Antarctic Peninsula, 1988–2009. *The Cryosphere.* 6(5), pp.10311048.
- 642 Ding Q., Wallace J.M., Battisti D.S., Steig E.J., Gallant A.J.E., Kim H. and Geng, L. 2014. Tropical forcing
 643 of the recent rapid Arctic warming in northeastern Canada and Greenland. *Nature*. 509, pp.209-212.
- Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carre, G., Marquez, J.R.G., Gruber, B.,
- Lafourcade, B., Leitao, P.J., Munkemuller, T., McClean, C., Osborne, P.E., Reineking, B., Schroder, B.,
- 546 Skidmore, A.K., Zurell, D. and Lautenbach, S. 2013. Collinearity: a review of methods to deal with it
- 647 and a simulation study evaluating their performance. *Ecography.* **36**(1), pp.27-46.
- Duguay, C.R., Prowse, T.D., Bonsal, B.R., Brown, R.D., Lacroix, M.P. and Menard, P. 2006. Recent trends
 in Canadian lake ice cover. *Hydrological Processes.* 20(4), pp.781-801.
- Eijpen, K.J., Warren, C.R. and Benn, D.I. 2003. Subaqueous melt rates at calving termini: a laboratory
 approach. *Annals of Glaciology.* 36(1), pp.179-183.
- Hall, D.K., Riggs, G.A. and Salomonson, V.V. 1995. Development of methods for mapping global snow
- 653 cover using moderate resolution imaging spectroradiometer data. *Remote Sensing of Environment.*
- 654 **54**(2), pp.127-140.

- Hanna, E., Huybrechts, P., Steffen, K., Cappelen, J., Huff, R., Shuman, C., Irvine-Fynn, T., Wise, S. and
 Griffiths, M. 2008. Increased runoff from melt from the Greenland Ice Sheet: A response to global
 warming. *Journal of Climate.* 21(2), pp.331-341.
- Hanna, E., Navarro, F.J., Pattyn, F., Domingues, C.M., Fettweis, X., Ivins, E.R., Nicholls, R.J., Ritz, C.,
 Smith, B., Tulaczyk, S., Whitehouse, P.L. and Zwally, H.J. 2013. Ice-sheet mass balance and climate
 change. *Nature.* 498(7452), pp.51-59.
- Haresign, E. and Warren, C.R. 2005. Melt rates at calving termini: a study at Glaciar León, Chilean
 Patagonia. *Geological Society, London, Special Publications.* 242(1), pp.99-109.
- Hill, E.A., Carr, J.R., Stokes, C.R. and Gudmundsson, G.H. 2018. Dynamic changes in outlet glaciers in
 northern Greenland from 1948 to 2015. *The Cryosphere*. **12**(10), pp.3243-3263.
- Holland, D.M., Thomas, R.H., de Young, B., Ribergaard, M.H. and Lyberth, B. 2008. Acceleration of
 Jakobshavn Isbræ triggered by warm subsurface ocean waters. *Nature Geoscience*. 1(10), pp.659-664.
- How, P., Messerli, A., Mätzler, E., Santoro, M., Wiesmann, A., Caduff, R., Langley, K., Bojesen, M.H.,
 Paul, F., Kääb, A. and Carrivick, J.L. 2021. Greenland-wide inventory of ice marginal lakes using a multimethod approach. *Scientific Reports.* 11, 4481.
- Howat, I.M. and Eddy, A. 2011. Multi-decadal retreat of Greenland's marine-terminating glaciers. *Journal of Glaciology.* 57(203), pp.389-396.
- Howat, I.M., Joughin, I., Fahnestock, M., Smith, B.E. and Scambos, T.A. 2008. Synchronous retreat and
 acceleration of southeast Greenland outlet glaciers 2000-06: ice dynamics and coupling to climate. *Journal of Glaciology.* 54(187), pp.646-660.
- Huggel, C., Kaab, A., Haeberli, W., Teysseire, P. and Paul, F. 2002. Remote sensing based assessment
 of hazards from glacier lake outbursts: a case study in the Swiss Alps. *Canadian Geotechnical Journal*. **39**(2), pp.316-330.
- Joughin, I., Smith, B.E., Howat, I.M., Floricioiu, D., Alley, R.B., Truffer, M. and Fahnestock, M. 2012.
 Seasonal to decadal scale variations in the surface velocity of Jakobshavn Isbrae, Greenland:
 Observation and model-based analysis. *Journal of Geophysical Research: Earth Surface*. 117, F02030.
- Joughin, I., Smith, B.E., Howat, I.M., Scambos, T. and Moon, T. 2010. Greenland flow variability from
 ice-sheet-wide velocity mapping. *Journal of Glaciology*. 56, pp.415-430.
- Kargel, J.S., Ahlstrøm, A.P., Alley, R.B., Bamber, J.L., Benham, T.J., Box, J.E., Chen, C., Christoffersen, P.,
 Citterio, M., Cogley, J.G., Jiskoot, H., Leonard, G.J., Morin, P., Scambos, T., Sheldon, T. and Willis, I.
 2012. Brief communication: Greenland's shrinking ice cover: "Fast times" but not that fast. *Cryosphere.*6(3), pp.533-537.
- King, O., Bhattacharya, A., Bhambri, R. and Bolch, T. 2019. Glacial lakes exacerbate Himalayan glacier
 mass loss. *Scientific Reports.* 9, 18145.
- King, O., Dehecq, A., Quincey, D. and Carrivick, J. 2018. Contrasting geometric and dynamic evolution
 of lake and land-terminating glaciers in the central Himalaya. *Global and Planetary Change.* 167,
 pp.46-60.
- 692 Kirkbride, M.P. 1993. The temporal significance of transitions from melting to calving termini at 693 glaciers in the central Southern Alps of New Zealand. *The Holocene*. **3**(3), pp.232-240.

Knight, P.G., Waller, R.I., Patterson, C.J., Jones, A.P. and Robinson, Z.P. 2000. Glacier advance, ice marginal lakes and routing of meltwater and sediment: Russell Glacier, Greenland. *Journal of Glaciology*. 46(154), pp.423-426.

Korsgaard, N.J., Nuth, C., Khan, S.A., Kjeldsen, K.K., Bjørk, A.A., Schomacker, A. and Kjær, K.H. 2016.
Digital elevation model and orthophotographs of Greenland based on aerial photographs from 1978– *Scientific Data*. 3(1), 160032.

Krabill, W., Hanna, E., Huybrechts, P., Abdalati, W., Cappelen, J., Csatho, B., Frederick, E., Manizade,
S., Martin, C., Sonntag, J., Swift, R., Thomas, R. and Yungel, J. 2004. Greenland Ice Sheet: Increased
coastal thinning. *Geophysical Research Letters.* **31**(24) L24402.

- Larsen, C.F., Burgess, E., Arendt, A.A., O'Neel, S., Johnson, A.J. and Kienholz, C. 2015. Surface melt dominates Alaska glacier mass balance. *Geophysical Research Letters.* **42**(14), pp.5902-5908.
- Lea, J.M., Mair, D.W.F. and Rea, B.R. 2014. Evaluation of existing and new methods of tracking glacier
 terminus change. *Journal of Glaciology.* 60(220), pp.323-332.
- Leclercq, P.W., Weidick, A., Paul, F., Bolch, T., Citterio, M. and Oerlemans, J. 2012. Brief
 communication "Historical glacier length changes in West Greenland". *Cryosphere.* 6(6), pp.13391343.
- Liu Q., Mayer C., Wang X., Nie Y., Wu K., Wei J. and Liu S. 2020. Interannual flow dynamics driven by
- 711 frontal retreat of a lake-terminating glacier in the Chinese Central Himalaya. *Earth and Planetary*
- 712 Science Letters. **546**, 116450.
- Lovell, A.M., Carr, J.R. and Stokes, C.R. 2019. Spatially Variable Glacier Changes in the Annapurna
 Conservation Area, Nepal, 2000 to 2016. *Remote Sensing.* 11(12), 1452.
- Mallalieu, J., Carrivick, J.L., Quincey, D.J., Smith, M.W. and James, W.H., 2017. An integrated Structurefrom-Motion and time-lapse technique for quantifying ice-margin dynamics. *Journal of Glaciology*.
 63(242), pp.937-949.
- Mallalieu, J., Carrivick, J.L., Quincey, D.J. and Smith, M.W. 2020. Calving Seasonality Associated With
 Melt-Undercutting and Lake Ice Cover. *Geophysical Research Letters.* 47(8), GL086561.
- McFadden, E.M., Howat, I.M., Joughin, I., Smith, B.E. and Ahn, Y. 2011. Changes in the dynamics of
 marine terminating outlet glaciers in west Greenland (2000–2009). *Journal of Geophysical Research: Earth Surface.* 116, F02022.
- 723 McFeeters, S.K. 1996. The use of the Normalized Difference Water Index (NDWI) in the delineation of 724 open water features. *International Journal of Remote Sensing*. **17**(7), pp.1425-1432.
- Mernild, S.H., Malmros, J.K., Yde, J.C. and Knudsen, N.T. 2012. Multi-decadal marine- and landterminating glacier recession in the Ammassalik region, southeast Greenland. *Cryosphere.* **6**(3), pp.625-639.
- Moon, T. and Joughin, I. 2008. Changes in ice front position on Greenland's outlet glaciers from 1992
 to 2007. *Journal of Geophysical Research-Earth Surface*. **113**, F02022.
- 730 Morlighem, M., Williams, C.N., Rignot, E., An, L., Arndt, J.E., Bamber, J.L., Catania, G., Chauché, N.,
- Dowdeswell, J.A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T.M.,
- Kjeldsen, K.K., Millan, R., Mayer, L., Mouginot, J., Noël, B.P.Y., O'Cofaigh, C., Palmer, S., Rysgaard, S.,
 Seroussi, H., Siegert, M.J., Slabon, P., Straneo, F., van den Broeke, M.R., Weinrebe, W., Wood, M. and
- 734 Zinglersen, K.B. 2017. BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of
- 735 Greenland From Multibeam Echo Sounding Combined With Mass Conservation. *Geophysical Research*
- 736 Letters. 44(21), pp.11051-11061.
- 737 Motyka, R.J., O'Neel, S., Connor, C.L. and Echelmeyer, K.A. 2003. Twentieth century thinning of
- 738 Mendenhall Glacier, Alaska, and its relationship to climate, lake calving, and glacier run-off. *Global and*
- 739 *Planetary Change.* **35**(1-2), pp.93-112.

- Mouginot, J., Rignot, E., Bjørk, A.A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl,
 B. and Wood, M. 2019. Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018.
- 742 *Proceedings of the National Academy of Sciences.* **116**(19), pp.9239-9244.
- Myers, P.G., Kulan, N. and Ribergaard, M.H. 2007. Irminger Water variability in the West Greenland
 Current. *Geophysical Research Letters*. 34(17), L17601.
- 745 Nick, F.M., Vieli, A., Andersen, M.L., Joughin, I., Payne, A., Edwards, T.L., Pattyn, F. and van de Wal,
- R.S.W. 2013. Future sea-level rise from Greenland's main outlet glaciers in a warming climate. *Nature*. **497**, pp.235-238.
- Naruse, R. and Skvarca, P. 2000. Dynamic features of thinning and retreating Glaciar Upsala, a
 Lacustrine Calving Glacier in southern Patagonia. *Arctic Antarctic and Alpine Research.* 32(4), pp.485491.
- 751 Pattyn, F., Ritz, C., Hanna, E., Asay-Davis, X., DeConto, R., Durand, G., Favier, L., Fettweis, X., Goelzer,
- H., Golledge, N.R., Kuipers Munneke, P., Lenaerts, J.T.M., Nowicki, S., Payne, A.J., Robinson, A.,
 Seroussi, H., Trusel, L.D. and van den Broeke, M. 2018. The Greenland and Antarctic ice sheets under
 1.5 °C global warming. *Nature Climate Change.* 8(12), pp.1053-1061.
- Pelto, M.S. and Hedlund, C. 2001. Terminus behavior and response time of North Cascade glaciers,
 Washington, U.S.A. *Journal of Glaciology*. 47(158), pp.497-506.
- Porter, D.F., Tinto, K.J., Boghosian, A.L., Csatho, B.M., Bell, R.E. and Cochran, J.R. 2018. Identifying
 Spatial Variability in Greenland's Outlet Glacier Response to Ocean Heat. *Frontiers in Earth Science*.
 6(90), 2018.00090.
- Price, S.F., Payne, A.J., Catania, G.A. and Neumann, T.A. 2008. Seasonal acceleration of inland ice via
 longitudinal coupling to marginal ice. *Journal of Glaciology*. 54(185), pp.213-219.
- Pritchard, H.D., Arthern, R.J., Vaughan, D.G. and Edwards, L.A. 2009. Extensive dynamic thinning on
 the margins of the Greenland and Antarctic ice sheets. *Nature*. 461(7266), pp.971-975.
- Purdie H., Bealing P., Tidey E., Gomez C. and Harrison J. 2016. Bathymetric evolution of Tasman Glacier
- terminal lake, New Zealand, as determined by remote surveying techniques. *Global and Planetary Change*. 147, pp.1-11.
- Quincey, D.J., Richardson, S.D., Luckman, A., Lucas, R.M., Reynolds, J.M., Hambrey, M.J. and Glasser,
 N.F. 2007. Early recognition of glacial lake hazards in the Himalaya using remote sensing datasets. *Global and Planetary Change*. 56(1), pp.137-152.
- Rastner, P., Bolch, T., Molg, N., Machguth, H., Le Bris, R. and Paul, F. 2012. The first complete inventory
 of the local glaciers and ice caps on Greenland. *Cryosphere*. 6(6), pp.1483-1495.
- Reynolds, J.M. 2000. On the formation of supraglacial lakes on debris-covered glaciers. In: Nakawo,
 M., et al. eds. *Debris-Covered Glaciers*. Wallingford: Int Assoc Hydrological Sciences, pp.153-161.
- 774 Roy, D.P., Wulder, M.A., Loveland, T.R., C.E, W., Allen, R.G., Anderson, M.C., Helder, D., Irons, J.R.,
- Johnson, D.M., Kennedy, R., Scambos, T.A., Schaaf, C.B., Schott, J.R., Sheng, Y., Vermote, E.F., Belward,
- A.S., Bindschadler, R., Cohen, W.B., Gao, F., Hipple, J.D., Hostert, P., Huntington, J., Justice, C.O., Kilic,
- A., Kovalskyy, V., Lee, Z.P., Lymburner, L., Masek, J.G., McCorkel, J., Shuai, Y., Trezza, R., Vogelmann,
 J., Wynne, R.H. and Zhu, Z. 2014. Landsat-8: Science and product vision for terrestrial global change
- research. *Remote Sensing of Environment*. **145**, pp.154-172.
- 780 Russell, A.J., Carrivick, J.L., Ingeman-Nielsen, T., Yde, J.C. and Williams, M. 2011. A new cycle of
- jökulhlaups at Russell Glacier, Kangerlussuaq, West Greenland. *Journal of Glaciology*. 57(202), pp.238246.
- Sakai, A. and Fujita, K. 2017. Contrasting glacier responses to recent climate change in high-mountain
 Asia. *Scientific Reports.* 7(1), 13717.

- Scherler, D., Bookhagen, B. and Strecker, M.R. 2011. Spatially variable response of Himalayan glaciers
 to climate change affected by debris cover. *Nature Geoscience*. 4(3), pp.156-159.
- Schild, K.M. and Hamilton, G.S. 2013. Seasonal variations of outlet glacier terminus position in
 Greenland. *Journal of Glaciology*. 59(216), pp.759-770.
- Schomacker, A. 2010. Expansion of ice-marginal lakes at the Vatnajokull ice cap, Iceland, from 1999 to
 2009. *Geomorphology*. **119**(3-4), pp.232-236.
- Seale, A., Christoffersen, P., Mugford, R.I. and O'Leary, M. 2011. Ocean forcing of the Greenland Ice
 Sheet: Calving fronts and patterns of retreat identified by automatic satellite monitoring of eastern
 outlet glaciers. *Journal of Geophysical Research-Earth Surface*. **116**, F03013.
- 794 Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, 795 K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., Geruo, A., Agosta, C., 796 Ahlstrøm, A., Babonis, G., Barletta, V.R., Bjørk, A.A., Blazquez, A., Bonin, J., Colgan, W., Csatho, B., 797 Cullather, R., Engdahl, M.E., Felikson, D., Fettweis, X., Forsberg, R., Hogg, A.E., Gallee, H., Gardner, A., 798 Gilbert, L., Gourmelen, N., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, 799 M., Khan, S., Kjeldsen, K.K., Konrad, H., Langen, P.L., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, 800 M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mottram, R., Mouginot, J., Moyano, G., Muir, A., 801 Nagler, T., Nield, G., Nilsson, J., Noël, B., Otosaka, I., Pattle, M.E., Peltier, W.R., Pie, N., Rietbroek, R., Rott, H., Sørensen, L.S., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W., 802 803 Simonsen, S.B., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., Jan van de Berg, W., van der
- Wal, W., van Wessem, M., Vishwakarma, B.D., Wiese, D., Wilton, D., Wagner, T., Wouters, B., Wuite,
 J. and The Imbie Team. 2020. Mass balance of the Greenland Ice Sheet from 1992 to 2018. *Nature*. **579**, pp.233-239.
- 807 Smejkalova, T., Edwards, M.E. and Dash, J. 2016. Arctic lakes show strong decadal trend in earlier 808 spring ice-out. *Scientific Reports.* **6**, 38449.
- Straneo, F. and Heimbach, P. 2013. North Atlantic warming and the retreat of Greenland's outlet glaciers. *Nature*. **504**(7478), pp.36-43.
- Surdu, C.M., Duguay, C.R. and Prieto, D.F. 2016. Evidence of recent changes in the ice regime of lakes
 in the Canadian High Arctic from spaceborne satellite observations. *Cryosphere.* 10(3), pp.941-960.
- Sutherland, J.L., Carrivick, J.L., Gandy, N., Shulmeister, J., Quincey, D.J. and Cornford, S.L. 2020.
 Proglacial lakes control glacier geometry and behavior during recession. *Geophysical Research Letters*.
 47(19), GL088865.
- Truffer, M. and Motyka, R.J. 2016. Where glaciers meet water: Subaqueous melt and its relevance to glaciers in various settings. *Reviews of Geophysics.* **54**, pp.220-239.
- Trusel, L.D., Das, S.B., Osman, M.B., Evans, M.J., Smith, B.E., Fettweis, X., McConnell, J.R., Noël, B.P.Y.
 and van den Broeke, M.R. 2018. Nonlinear rise in Greenland runoff in response to post-industrial Arctic
 warming. *Nature*. 564(7734), pp.104-108.
- 821 Trussel, B.L., Motyka, R.J., Truffer, M. and Larsen, C.F. 2013. Rapid thinning of lake-calving Yakutat
- Glacier and the collapse of the Yakutat Icefield, southeast Alaska, USA. *Journal of Glaciology*. 59(213),
 pp.149-161.
- Tsutaki S., Fujita K., Nuimura T., Sakai A., Sugiyama S., Komori J. and Tshering P. 2019. Contrasting thinning patterns between lake- and land-terminating glaciers in the Bhutanese Himalaya. *The Cryosphere*. **13**, pp. 2733-2750.
- 827 Van Tatenhove, F.G.M., Roelfsema, C.M., Blommers, G. and Voorden, A.V. 1995. Change in position
- and altitude of a small outlet glacier during the period 1943–92: Leverett Glacier, West Greenland. *Annals of Glaciology*. 21, pp.251-258.

- Warren, C.R. 1991. Terminal environment, topographic control and fluctuations of West Greenlandglaciers. *Boreas.* 20(1), pp.1-15.
- Warren, C.R., Greene, D.R. and Glasser, N.F. 1995. Glaciar Upsala, Patagonia: Rapid calving retreat in
 fresh water. *Annals of Glaciology.* 21, pp.311-316.
- 834 Warren, C.R. and Kirkbride, M.P. 2003. Calving speed and climatic sensitivity of New Zealand lake-835 calving glaciers. *Annals of Glaciology*. **36**, pp.173-178.
- Weidick, A. 1991. Present-day expansion of the southern part of the Inland Ice. *Rapport Gronlands Geologiske Undersogelse*. **152**, pp.73-79.
- Zwally, H.J. 1989. Growth of Greenland ice sheet Interpretation. *Science*. **246**(4937), pp.1589-1591.
- 839