

Est.
1841

YORK
ST JOHN
UNIVERSITY

Bartels, Susan A., Bolarinwa, Obasanjo
ORCID: <https://orcid.org/0000-0002-9208-6408>, Fortune, Effiong,
Aboagye, Richard Gyan ORCID: <https://orcid.org/0000-0002-3498-2909>, Seidu, Abdul-Aziz ORCID: <https://orcid.org/0000-0001-9734-9054>, Olagunju, Olalekan Seun, Nwagbara, Ugochinyere Ijeoma
ORCID: <https://orcid.org/0000-0003-1771-0570>, Ameyaw, Edward
Kwabena ORCID: <https://orcid.org/0000-0002-6617-237X> and
Ahinkorah, Bright Opoku (2021) Health facility delivery among
women of reproductive age in Nigeria: Does age at first birth
matter? PLOS ONE, 16 (11). e0259250.

Downloaded from: <http://ray.yorks.ac.uk/id/eprint/8475/>

The version presented here may differ from the published version or version of record. If
you intend to cite from the work you are advised to consult the publisher's version:
<http://dx.doi.org/10.1371/journal.pone.0259250>

Research at York St John (RaY) is an institutional repository. It supports the principles of
open access by making the research outputs of the University available in digital form.
Copyright of the items stored in RaY reside with the authors and/or other copyright
owners. Users may access full text items free of charge, and may download a copy for
private study or non-commercial research. For further reuse terms, see licence terms
governing individual outputs. [Institutional Repository Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at ray@yorks.ac.uk

RESEARCH ARTICLE

Health facility delivery among women of reproductive age in Nigeria: Does age at first birth matter?

Obasanjo Afolabi Bolarinwa^{1,2*}, Effiong Fortune³, Richard Gyan Aboagye⁴, Abdul-Aziz Seidu^{5,6}, Olalekan Seun Olagunju², Ugochinyere Ijeoma Nwagbara¹, Edward Kwabena Ameyaw⁷, Bright Opoku Ahinkorah⁷

1 Department of Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal, Durban, South Africa, **2** Faculty of Social Sciences, Department of Demography and Social Statistics, Obafemi Awolowo University, Ile-Ife, Nigeria, **3** Department of Medical Laboratory Science, University of Calabar, Calabar, Nigeria, **4** School of Public Health, University of Health and Allied Sciences, Ho, Ghana, **5** College of Public Health, Medical and Veterinary Sciences, James Cook University, Townsville, Australia, **6** Department of Estate Management, Takoradi Technical University, Takoradi, Ghana, **7** School of Public Health University of Technology Sydney, Sydney, Australia

* bolarinwaobasanjo@gmail.com



OPEN ACCESS

Citation: Bolarinwa OA, Fortune E, Aboagye RG, Seidu A-A, Olagunju OS, Nwagbara UI, et al. (2021) Health facility delivery among women of reproductive age in Nigeria: Does age at first birth matter? PLoS ONE 16(11): e0259250. <https://doi.org/10.1371/journal.pone.0259250>

Editor: Susan A. Bartels, Queen's University at Kingston, CANADA

Received: February 26, 2021

Accepted: October 16, 2021

Published: November 4, 2021

Copyright: © 2021 Bolarinwa et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The dataset is freely available at <https://dhsprogram.com/data/available-datasets.cfm> and can be accessed after request has been made and request intent has been approved by DHS member. No specific access is required all requests are judged independently based on request made. The authors did not have any special access privileges that others would not have.

Funding: The authors received no specific funding for this work.

Abstract

Background

High maternal mortality ratio in sub-Saharan Africa (SSA) has been linked to inadequate medical care for pregnant women due to limited health facility delivery utilization. Thus, this study, examined the association between age at first childbirth and health facility delivery among women of reproductive age in Nigeria.

Methods

The study used the most recent secondary dataset from Nigeria's Demographic and Health Survey (NDHS) conducted in 2018. Only women aged 15–49 were considered for the study (N = 34,193). Bi-variate and multivariable logistic regression models were used to examine the association between age at first birth and place of delivery. The results were presented as crude odds ratios and adjusted odds ratios (aOR) with corresponding 95% confidence intervals (CIs). Statistical significance was set at $p < 0.05$.

Results

The results showed that the prevalence of health facility deliveries was 41% in Nigeria. Women who had their first birth below age 20 [aOR = 0.82; 95% (CI = 0.74–0.90)] were less likely to give birth at health facilities compared to those who had their first birth at age 20 and above.

Conclusion

Our findings suggest the need to design interventions that will encourage women of reproductive age in Nigeria who are younger than 20 years to give birth in health facilities to avoid

Competing interests: The authors have declared that no competing interest exist.

the risks of maternal complications associated with home delivery. Such interventions should include male involvement in antenatal care visits and the education of both partners and young women on the importance of health facility delivery.

Background

The Sustainable Development Goal (SDG) 3 aims to ensure healthy lives and promote well-being for all ages [1]. However, high maternal mortality ratio (MMR), especially in low-and middle-income countries (LMICs), has remained worrisome [2–4]. In line with this, SDG 3.1 calls for the global MMR reduction to less than 70 per 100,000 live births by the year 2030 [1].

Although many countries have reduced their MMRs, causing a general global decline, efforts to reduce MMRs of countries in sub-Saharan Africa (SSA) have largely stagnated [5]. Nigeria's maternal and child health indices are among the worst in the world. In 2015, the World Health Organization (WHO) estimated that 19% of global maternal deaths occurred in Nigeria [5], with an estimated MMR of 814 per 100,000 live births, and this placed a lifetime risk of maternal death at 1 in 22 in contrast to 1 in 4900 in developed countries [5]. However, the 2018 Nigeria Demographic Health Survey (NDHS) estimated MMR of 512 per 100,000 live births. Despite this reduction, Nigeria still remains one of the countries with the highest MMR globally [6].

Neonatal mortality is another serious public health threat in SSA [7]. In 2019, SSA had an estimate of 27 per 1000 live births, while Nigeria reported 36 per 1000 live births out of the estimate [8]. Similar to MMR, efforts to reduce neonatal mortality in SSA, especially in Nigeria, have been relatively slow [9]. Between 2010 and 2019, only a 13% reduction in neonatal mortality ratio was achieved in Nigeria [8,10]. The high prevalence of maternal and neonatal deaths in SSA and Nigeria has been linked to inadequate medical care for pregnant women, resulting in low health facility delivery [11–14]. Health facility delivery is an important aspect of maternal care because of the presence of skilled birth professionals [12], as well as an enabling environment where resources necessary for effective management of obstetric complications are available [15,16].

A WHO report showed a 50 percent reduction in Egypt's maternal mortality when the number of births assisted by qualified professionals doubled [17]. This implies that maternal and neonatal mortality in Nigeria can be reduced by ensuring that women have better maternal healthcare services, especially health facility delivery [18,19]. Although home delivery may be relatively cheaper for these women, evidence suggests that homedelivery can pose a serious health risk to both the mother and infant [20–22].

Other than financial constraints, other identified factors for poor utilization of maternal health services in SSA include maternal age, maternal and partner literacy level, poor knowledge about obstetric danger signs, cultural beliefs, poor referral practices, scarcity of trained health workers, and poor coordination among staff [18,19]. These barriers have been found in Nigeria [11,23,24].

Studies in the past have suggested age at first delivery as a determinant of health facility delivery in Nigeria [11,23]. For instance, a study conducted among women in a semi-urban settlement in Zaria, Northern Nigeria, found that 77% of women who had first pregnancies before age 18 delivered at home compared to their older women counterparts with 56% home deliveries [11]. This implies that younger women in Nigeria tend to deliver at home compared to older women [11,23].

Similarly, the 2013 NDHS conducted estimated that 23% of Nigerian women aged 15–19 had begun childbearing, of which 17 percent out of this had their first child, and 5% were

pregnant with their first child [25]. The 2018 NDHS also showed that the median age at first birth among women age 25–49 was 20.4 years. This means that half of the women aged 25–49 gave birth for the first time before age 21 [6].

Therefore, there is a need to examine how age at first birth is associated with health facility delivery in Nigeria. Hence, this study aimed at examining the role age at first birth plays in health facility delivery among women of reproductive age in Nigeria. The result of this study would be useful to public health policymakers in building the right approaches to take in encouraging health facility delivery in Nigeria.

Materials and methods

Study design and data source

This study involved a cross-sectional analysis of secondary data from NDHS conducted in 2018. We utilized data from the birth records (NGIR7ADT) in the women's file. According to Corsi and Neuman [26], the Demographic Health Survey (DHS) is a nationally representative survey usually carried out to estimate sociodemographic, health, and other health-related indicators. The survey collects information on several indicators, including the place of childbirth and age at first childbirth. The survey is conducted in over 85 LMICs globally [26]. The 2018 NDHS employed a two-stage stratified sampling technique to recruit the respondents for the survey. A detailed explanation of the sampling procedures can be found in Aliaga and Ruilin [27] study and the 2018 NDHS report [6]. The 2018 NDHS dataset consisted of 41,821 individual women aged 15–49. 40,666 occupied households were selected for the sample, and 40,427 were interviewed successfully, resulting in a response rate of 99% [6]. Out of this, 34,193 of the respondents with complete cases on the variables of interest in this study were included in the final analysis. We relied on strengthening the reporting of observational studies in Epidemiology in writing this manuscript [28].

Variables

Outcome variable. The main outcome variable was the place of delivery. NDHS asks questions on place of delivery within five years preceding the survey, and the response options received were respondent's home, other homes, government hospital, government health center, government health post, other public sectors, private hospital/clinic, other private sectors, and other health facilities. Those who responded respondent's home and other homes were classified as "home delivery," while the remaining responses were grouped as "health facility delivery". This classification was informed by previous studies [29–32].

Key explanatory variable. The main explanatory variable was the age at first childbirth. This was assessed using the variable "age of respondent at first childbirth". The response options ranged from 12 years to 48 years. The age at first childbirth was then created by recoding the responses into "below 20 years" and "20 years and above". This categorization was informed by previous findings on age at first birth in Nigeria [6,25] and elsewhere [33].

Covariates. A total of thirteen covariates were studied. The covariates were selected based on their significant association with health facility delivery [29,30,34,35] and their availability in the DHS dataset. The covariates include maternal age, place of residence, parity, level of education, wealth quintile, current working status, marital status, religion, visit to the health facility in the past 12 months, permission to get medical help, distance to the health facility, region, and exposure to media. For this study, several of the covariates were recoded. The pre-existing coding for the place of residence, distance to the health facility, permission to get medical help, region, and visited health facility last 12 months were maintained and used in the final analysis. Maternal age was recoded into "15–24", "25–34", and "35+". The level of

education was re-classified into “no education”, “primary”, and “secondary or higher”. Wealth index was recoded as “poor”, “middle”, and “rich” by recoding poorer wealth quantile and poorest wealth quantile as “poor”, richer wealth quantile and richest wealth quantile as “Rich”, Middle wealth quantile still retains the categorisation as “Middle”. This categorization has been utilized elsewhere as well [36,37]. Also, marital status was recoded as “single”, “married”, and “others”. Parity was re-grouped into “1–3”, “4–6”, and “7+”. The women’s current working status was ground into “yes” and “no”. Religious affiliation of the women was recoded as “Christian”, “Islam” and traditional and other religious were recategorized as “others”. Exposure to media was created from three (3) variables (frequency of watching television, frequency of reading newspaper/magazine, and frequency of listening to the radio). All the three variables had the same response options (not at all less than once a week and at least once a week). The responses were recoded as “No = not at all” and “Yes” (less than once a week and at least once a week). Later, an index called the media exposure variable was created. In the current study, media exposure was defined as a woman with exposure in all three variables (watching television, reading newspaper/magazine, and listening to the radio).

Data analysis

Data were analyzed using Stata version 16.0 (Stata Corporation, College Station, TX, USA). A descriptive analysis was first performed to determine the proportion of health facility delivery, with the result presented using percentages. A Pearson chi-square test analysis was later carried out to examine the relationship between health facility and age at first birth and the studied covariates. Also, bivariate and multivariable binary logistic regression analyses were performed to determine the association between health facility and age at first birth and selected covariates. The results were presented using crude odds ratio [38] and adjusted odds ratio (aOR) at 95% confidence intervals (CIs). The level of statistical significance was set at $p < 0.05$ in the chi-square and regression analyses. A multicollinearity test was conducted using the Variance Inflation Factor (VIF), and we found no evidence of multicollinearity among the studied variables. During the analysis, the women’s sample weights (v005/1,000,000) were applied to obtain unbiased estimates. According to the DHS guidelines and the survey command (SVY) in Stata, it was used to adjust the data’s complex sampling structure in both the chi-square and regression analyses.

Ethical approval

This was a secondary analysis of data, and therefore no ethical approval was required since the data is available in the public domain. Further information about the DHS data usage and ethical standards is available at <http://goo.gl/ny8T6X>. The authors of this manuscript did not collect the data, we sought permission from the MEASURE DHS website, access to the latest dataset of Nigeria was provided after our intent for the request was assessed and approved on the 20th of March 2021. The DHS surveys are ethically accepted by the ORC Macro Inc. Ethics Committee and the Ethics Boards of partner organizations in different countries, such as the Ministries of Health. The interviewed women gave either written or verbal consent during each of the surveys and followed Helsinki’s declaration of ethical principles.

Results

Distribution of key independent variable, covariate, and place of delivery

[Table 1](#) presents results on the distribution of age at first childbirth, covariates, and place of delivery among women of reproductive ages in Nigeria. The prevalence of health facility

Table 1. Weighted distribution of key independent and covariate variables and place of delivery (n = 34,193).

Variables	Percentage (%)	Place of delivery (%)		p-value (χ^2)
		Home	Health facility	
Age at first birth				<0.001
20 years and above	42.04	39.80	60.20	
Below 20 years	57.96	72.94	27.06	
Age				<0.001
15–24	23.82	66.25	33.75	
25–34	50.84	56.20	43.80	
35 +	25.34	57.82	42.18	
Place of residence				<0.001
Urban	38.52	36.34	63.66	
Rural	61.48	73.20	26.8	
Parity				<0.001
1–3	46.46	49.07	50.93	
4–6	34.29	61.19	38.81	
7+	19.25	79.09	20.91	
Level of education				<0.001
No education	46.38	85.98	14.02	
Primary	14.93	56.4	43.60	
Secondary and above	38.70	27.68	72.32	
Wealth index				<0.001
Poor	44.91	83.06	16.94	
Middle	20.60	58.08	41.92	
Rich	34.50	28.24	71.76	
Currently working				<0.001
No	32.41	71.32	28.68	
Yes	67.59	53.10	46.9	
Marital status				<0.001
Single	1.71	40.89	59.11	
Married	92.63	60.08	39.92	
Others	5.66	46.08	53.20	
Religion				<0.001
Christian	35.99	30.85	69.15	
Islam	63.48	74.83	25.17	
Others	0.53	75.30	24.70	
Visited health facility last 12 months				<0.001
No	46.68	65.42	34.58	
Yes	53.32	53.39	46.61	
Region				<0.001
North Central	13.51	50.13	49.87	
North East	18.17	74.50	25.50	
North West	36.73	84.42	15.58	
South East	10.02	17.18	82.82	
South-South	8.68	43.83	56.17	
South West	12.89	16.79	83.21	
Permission to get medical help				<0.001
Big problem	11.80	72.40	27.60	
Not a big problem	88.20	57.21	42.79	

(Continued)

Table 1. (Continued)

Variables	Percentage (%)	Place of delivery (%)		p-value (χ^2)
		Home	Health facility	
Distance to the health facility				<0.001
Big problem	28.09	69.3	30.7	
Not a big problem	71.91	54.98	45.02	
Exposure to media				<0.001
No	38.87	81.18	18.82	
Yes	61.13	44.9	55.10	
Nigeria = 34,193	100	59.00	41.00	

Weighted NDHS, 2018.

<https://doi.org/10.1371/journal.pone.0259250.t001>

deliveries was 41%. Age at first birth shows that 72.94% of home deliveries were among women who had their first birth below the age of 20. The chi-square (χ^2) results showed a significant relationship ($p < 0.001$) between age at first childbirth all selected co-variables, and place of delivery.

Association between age at first birth and health facility delivery in Nigeria

Table 2 shows the results of age at first birth and co-variables associated with health facility delivery in Nigeria. There is a significant relationship between the key independent variable

Table 2. Bivariate and multivariable logistics regression results on the predictors of health facility delivery (n = 34,193).

Variables	Unadjusted		Adjusted	
	cOR	95% CI	aOR	95% CI
Age at first birth (RC = 20 year & above)				
Below 20 years	0.24***	0.22–0.27	0.82***	0.74–0.90
Age group (RC = 15–24)				
25–34	1.53***	1.39–1.68	1.01	0.90–1.13
35 +	1.43***	1.28–1.60	1.21*	1.03–1.42
Place of residence (RC = Urban)				
Rural	0.21***	0.17–0.25	0.79**	0.68–0.92
Parity (RC = 1–3 children)				
4–6	0.61***	0.57–0.66	0.78***	0.70–0.87
7+	0.25***	0.23–0.28	0.65***	0.55–0.76
Level of education (RC = No education)				
Primary	4.74***	0.06–5.53	1.74***	1.53–1.97
Secondary and above	16.03***	13.90–18.47	2.82***	2.44–3.25
Wealth status (RC = Poor)				
Middle	3.54***	2.48–3.18	1.70***	1.50–1.93
Rich	12.46***	10.57–14.69	2.69***	2.32–3.12
Currently working (RC = No)				
Yes	2.20***	1.99–2.43	1.17**	1.06–1.30
Marital status (RC = Single)				
Married	0.46***	0.37–0.57	1.24	0.98–1.57
Others	0.79	0.61–1.01	1.05	0.79–1.39
Religion (RC = Christian)				

(Continued)

Table 2. (Continued)

Variables	Unadjusted		Adjusted	
	cOR	95% CI	aOR	95% CI
Islam	0.15***	0.13–0.18	0.58***	0.48–0.69
Others	0.15***	0.08–0.28	0.43**	0.26–0.70
Region (RC = North Central)				
North East	0.34***	0.27–0.44	0.65***	0.53–0.78
North West	0.19***	0.14–0.24	0.32***	0.27–0.39
South East	4.85***	3.53–6.66	1.69***	1.10–1.51
South South	1.29*	1.00–1.65	0.44***	0.43–0.64
South West	4.98***	3.85–6.45	2.14***	1.11–1.53
Visited health facility in last 12 months (RC = No)				
Yes	1.65***	1.50–1.81	1.46***	1.33–1.60
Permission to get Medical help (RC = Big problem)				
Not a big problem	1.96***	1.63–2.36	1.30***	1.12–1.51
Distance to health facility (RC = Big problem)				
Not a big problem	1.85***	1.62–2.11	1.30***	1.16–1.45
Exposure to media (RC = No)				
Yes	5.29***	4.71–5.95	1.44***	1.30–1.60

Weighted NDHS, 2018.

RC = Recode; cOR = unadjusted odds ratios; aOR = adjusted odds ratios; CI = confidence interval.

* $p < 0.05$

** $p < 0.01$

*** $p < 0.001$.

<https://doi.org/10.1371/journal.pone.0259250.t002>

(age at first birth) and health facility delivery among women of reproductive age in Nigeria. The adjusted odds ratio shows that women who had their first birth below age 20 [aOR = 0.82; 95% (CI = 0.74–0.90)] were less likely to give birth at health facilities compared to those who had their first birth at age 20 and above. Among the selected co-variables, age of respondents, place of residence, parity, level of education, wealth index, currently working, religion, region, visit to health facility, Permission to get Medical help, distance to health facility and exposure to media were significant with health facility delivery.

Discussion

This study assessed the age at first birth and its association with health facility delivery among women of reproductive age in Nigeria. This is important to prioritize the healthcare needs of women of reproductive age, considering that place of delivery determines the quality of care received by women of reproductive age in Nigeria.

Our study showed a prevalence of 59% homedeliveries in Nigeria among women of reproductive-aged 15–49. This result is higher than a study conducted by Olusanya and Alakija [39] that reported 51.4% of non-utilization of facility-based maternity services in Lagos, Nigeria. In the same vein, the study result is higher compared to a study conducted in Tanzania that reported 35.5% [40] but lower than a study conducted in Ethiopia with 81.1% prevalence of homedelivery among the same characteristics of women [41].

We also found a significant relationship between the key independent variable (age at first birth) and health facility delivery among women of reproductive age in Nigeria. The adjusted odds ratio showed that women who had their first birth below age 20 were less likely to give

birth at health facilities compared to those who had their first birth at age 20 and above. This could be due to the stigmatization that goes with unintended pregnancy outside marriage, and the women might prefer home delivery to avoid being stigmatized by the health attendants. Similar studies in sub-Saharan Africa showed that choosing a health facility as the place of delivery among women of reproductive age increased with age [42,43].

Among the selected co-variables, age of respondents, place of residence, parity, level of education, wealth index, currently working, religion, region, visit to the health facility, permission to get medical help, distance to health facility and exposure to media were significantly associated with health facility delivery.

Our study showed that respondents who were residing in the rural area of Nigeria were less likely to give birth at the health facility than those from the urban area. This is evidently showing the rural-urban inequality that exists in accessing health care facilities. Rural residents face numerous barriers to healthcare, which limits them from conveniently accessing and obtaining the appropriate care needed [20,44].

Women with number of children between 4 and 6 were less likely to give birth at the health facility alongside women with 7 children and above. Inadequate resources resulting from large family size and negative experiences from previous pregnancies have been identified as barriers to seeking maternal healthcare services by women with many children [45,46].

Religious beliefs and practices significantly determine a woman's place of birth delivery. Women practicing Islam were less likely to give birth at the health facility compared to their Christian counterparts. This finding is based on religious grounds, whereby women who practice Islam prefer a female birth attendant for delivery, hence, avoid visiting health care facilities so as not to be attended to by a male healthcare worker [47–50].

In the same vein, the women who reside in the North East and South-South regions of Nigeria were less likely to give birth at the health facilities compared to those residing in the South West region. Despite the fact that the South-South region of Nigeria is rich in oil, yet the rate of facility delivery is low within the region [31,46]. This could be due to the disparities in socioeconomic status, distribution of healthcare services, and culture in Nigeria's various regions [51]. Similarly, a study conducted in Kenya found a disparity in place of delivery by region as women residing in the Northern region were less likely to utilise health facilities for delivery [52].

We found that the odds of choosing to give birth at the health facility were higher among women age 35 and above. This is contrary to another study conducted in SSA, which showed that the odds of choosing a health facility as the place of delivery declined with age among reproductive age women [29]. Women who had secondary education and above were more likely to give birth at the health facility than women with primary education or no education. This shows that the level of education can influence a woman's choice of a place to give birth. This finding is consistent with studies that reported that a woman's educational status is significantly associated with the health facility delivery [53–56].

Our study also showed that rich women were more likely to give birth at Nigeria's health facilities. This is in concordance with a study conducted in Ghana that wealthy women were more likely to utilize health facilities for birth delivery [57]. Similarly, women who are currently working were more likely to give birth at the health facility. The out-of-pocket payment system for healthcare services poses a huge barrier for unemployed women who do not have any means of livelihood to access the health facility to give birth, which also affects poor women [51].

The findings of this research showed that women residing closer to the health facility were more likely to give birth at the health facility than women who reside far from the health facility. The distance to health facilities has been recognized as a determining factor for the

previously utilized healthcare facilities [58]. Also, women who were exposed to mass media were more likely to give birth at the health facilities than their counterparts who were not exposed to mass media. This shows that the mass media plays a significant role in positively influencing the better practice of women's reproductive health behavior [29,59]. This aligns with a spatial analysis study conducted in Nigeria in 2014 [46] and a recent study conducted in Ethiopia [60].

Strengths and limitations

This study's major strength is using the current nationally representative survey, which makes the study's findings generalizable to women of reproductive age in Nigeria. Despite this strength, there are some limitations inherent in this study that must be acknowledged. Firstly, because the age at first childbirth and place of delivery were both self-reported, this study is subject to recall bias. Thus, respondents may over-or under-report their age at first childbirth or the place of delivery which could affect the findings' interpretations. Secondly, the study dataset was cross-sectional, so it was only possible to conclude an association between age at first childbirth and health facility delivery and not causality. Thirdly, Nigeria is diverse, and the geopolitical zones have socio-cultural differences dating back to precolonial times; hence, the finding on age at first childbirth (and other covariates) may not hold across all the six zones. Hence, future studies may consider addressing this limitation.

Conclusion and recommendation

Our findings call for the need to implement health education programmes among women of reproductive age in Nigeria to increase awareness on the importance of giving birth at the health facilities to avoid the risks of maternal complications associated with home delivery. These health education programmes should be targeted at women of reproductive age younger than 20 years, those with no education, those currently not working, the poor, those currently residing in the Northern part of the country, those practicing Islam and those not exposed to mass media. Since one of the factors contributing to homedelivery was wealth index, and those who are in the poor wealth index group were less likely to utilize healthcare facilities for delivery, then Nigeria government and non-governmental organization should make healthcare payment more affordable for every woman of reproductive age by establishing a separate health payment scheme that will be targeted to women within poor wealth index.

Author Contributions

Conceptualization: Obasanjo Afolabi Bolarinwa.

Formal analysis: Obasanjo Afolabi Bolarinwa, Olalekan Seun Olagunju.

Investigation: Obasanjo Afolabi Bolarinwa.

Methodology: Obasanjo Afolabi Bolarinwa, Richard Gyan Aboagye, Abdul-Aziz Seidu.

Software: Obasanjo Afolabi Bolarinwa.

Validation: Obasanjo Afolabi Bolarinwa.

Writing – original draft: Obasanjo Afolabi Bolarinwa, Effiong Fortune, Richard Gyan Aboagye, Abdul-Aziz Seidu, Ugochinyere Ijeoma Nwagbara, Edward Kwabena Ameyaw, Bright Opoku Ahinkorah.

Writing – review & editing: Obasanjo Afolabi Bolarinwa, Effiong Fortune, Richard Gyan Aboagye, Abdul-Aziz Seidu, Olalekan Seun Olagunju, Ugochinyere Ijeoma Nwagbara, Edward Kwabena Ameyaw, Bright Opoku Ahinkorah.

References

1. World Health Organization [WHO]. World health statistics 2016: monitoring health for the SDGs sustainable development goals: World Health Organization; 2016.
2. World Health Organization [WHO]. Maternal Mortality: World Health Organization. 2014.
3. Yayla M. Maternal mortality in developing countries. *Journal of perinatal medicine*. 2003; 31(5):386–91. <https://doi.org/10.1515/JPM.2003.059> PMID: 14601259
4. Girum T, Wasie A. Correlates of maternal mortality in developing countries: an ecological study in 82 countries. *Maternal health, neonatology and perinatology*. 2017; 3(1):1–6. <https://doi.org/10.1186/s40748-017-0059-8> PMID: 29142757
5. World Health Organization [WHO]. Trends in maternal mortality: 1990–2015: estimates from WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division: World Health Organization; 2015.
6. National Population Commission—NPC, ICF. Nigeria Demographic and Health Survey 2018—Final Report. Abuja, Nigeria: NPC and ICF; 2019.
7. Andegiorgish AK, Andemariam M, Temesghen S, Ogbai L, Ogbe Z, Zeng L. Neonatal mortality and associated factors in the specialized neonatal care unit Asmara, Eritrea. *BMC public health*. 2020; 20(1):1–9. <https://doi.org/10.1186/s12889-019-7969-5> PMID: 31898494
8. United Nations Children’s Fund (UNICEF). The neonatal period is the most vulnerable time for a child 2019 [Available from: <https://data.unicef.org/topic/child-survival/neonatal-mortality/>].
9. Lawn JE, Kerber K, Enweronu-Laryea C, Cousens S, editors. 3.6 million neonatal deaths—what is progressing and what is not? *Seminars in perinatology*; 2010: Elsevier. <https://doi.org/10.1053/j.semperi.2010.09.011> PMID: 21094412
10. Rajaratnam JK, Marcus JR, Flaxman AD, Wang H, Levin-Rector A, Dwyer L, et al. Neonatal, postneonatal, childhood, and under-5 mortality for 187 countries, 1970–2010: a systematic analysis of progress towards Millennium Development Goal 4. *The Lancet*. 2010; 375(9730):1988–2008.
11. Idris S, Gwarzo U, Shehu A. Determinants of place of delivery among women in a semi-urban settlement in Zaria, northern Nigeria. *Annals of African medicine*. 2006; 5(2):68–72.
12. Nuamah GB, Agyei-Baffour P, Mensah KA, Boateng D, Quansah DY, Dobin D, et al. Access and utilization of maternal healthcare in a rural district in the forest belt of Ghana. *BMC pregnancy and childbirth*. 2019; 19(1):1–11. <https://doi.org/10.1186/s12884-018-2145-y> PMID: 30606156
13. Victora CG, Requejo JH, Barros AJ, Berman P, Bhutta Z, Boerma T, et al. Countdown to 2015: a decade of tracking progress for maternal, newborn, and child survival. *The Lancet*. 2016; 387(10032):2049–59.
14. Panel AP. Maternal health. Geneva: Investing in the Lifeline of Healthy Societies and Economies Policy Brief, Africa Progress Panel. 2010.
15. Bell JS, Curtis SL, Alayón S. Trends in delivery care in six countries. 2003.
16. Morgan A, Jimenez Soto E, Bhandari G, Kermode M. Provider perspectives on the enabling environment required for skilled birth attendance: a qualitative study in western Nepal. *Tropical Medicine & International Health*. 2014; 19(12):1457–65. <https://doi.org/10.1111/tmi.12390> PMID: 25252172
17. World Health Organization [WHO]. Success factors for women’s and children’s health: Egypt. 2015.
18. Kyei-Nimakoh M, Carolan-Olah M, McCann TV. Access barriers to obstetric care at health facilities in sub-Saharan Africa—a systematic review. *Systematic reviews*. 2017; 6(1):1–16. <https://doi.org/10.1186/s13643-016-0385-3> PMID: 28077170
19. Geleto A, Chojenta C, Musa A, Loxton D. Barriers to access and utilization of emergency obstetric care at health facilities in sub-Saharan Africa: a systematic review of literature. *Systematic reviews*. 2018; 7(1):1–14. <https://doi.org/10.1186/s13643-017-0670-9> PMID: 29304876
20. Kifle MM, Kesete HF, Gaim HT, Angosom GS, Araya MB. Health facility or home delivery? Factors influencing the choice of delivery place among mothers living in rural communities of Eritrea. *Journal of Health, Population and Nutrition*. 2018; 37(1):1–15.
21. Wanjira C, Mwangi M, Mathenge E, Mbugua G. Delivery practices and associated factors among mothers seeking child welfare services in selected health facilities in Nyandarua South District, Kenya. *BMC public health*. 2011; 11(1):1–9. <https://doi.org/10.1186/1471-2458-11-360> PMID: 21599994

22. Ahinkorah BO. Non-utilization of health facility delivery and its correlates among childbearing women: a cross-sectional analysis of the 2018 Guinea demographic and health survey data. *BMC health services research*. 2020; 20(1):1–10. <https://doi.org/10.1186/s12913-020-05893-0> PMID: 33167985
23. Okeshola FB, Sadiq IT. Determinants of home delivery among Hausa in Kaduna south local government area of Kaduna state, Nigeria. *American International Journal of contemporary research*. 2013; 3(5):78–85.
24. Envuladu E, Agbo H, Lassa S, Kigbu J, Zoakah A. Factors determining the choice of a place of delivery among pregnant women in Russia village of Jos North, Nigeria: achieving the MDGs 4 and 5. *International Journal of Medicine and Biomedical Research*. 2013; 2(1):23–7.
25. National Population Commission—NPC/Nigeria, ICF International. *Nigeria Demographic and Health Survey 2013*. Abuja, Nigeria: NPC/Nigeria and ICF International; 2014.
26. Corsi DJ, Neuman M, Finlay JE, Subramanian S. Demographic and health surveys: a profile. *International journal of epidemiology*. 2012; 41(6):1602–13. <https://doi.org/10.1093/ije/dys184> PMID: 23148108
27. Aliaga A, Ruilin's R, editors. *Cluster optimal sample size for demographic and health surveys*. 7th International Conference on Teaching Statistics—ICOTS; 2006.
28. Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *International journal of surgery*. 2014; 12(12):1495–9. <https://doi.org/10.1016/j.ijsu.2014.07.013> PMID: 25046131
29. Adde KS, Dickson KS, Amu H. Prevalence and determinants of the place of delivery among reproductive age women in sub-Saharan Africa. *Plos one*. 2020; 15(12):e0244875. <https://doi.org/10.1371/journal.pone.0244875> PMID: 33382825
30. Dickson KS, Adde KS, Amu H. What influences where they give birth? Determinants of place of delivery among women in rural Ghana. *International journal of reproductive medicine*. 2016; 2016. <https://doi.org/10.1155/2016/7203980> PMID: 28101522
31. Alabi TA, Atinge S, Ejim C, Adejoh SO. Does where mothers live matter? Regional variations in factors influencing place of child delivery in Nigeria. *Journal of Biosocial Science*. 2020:1–21. <https://doi.org/10.1017/S0021932020000747> PMID: 33303042
32. Lukman Solanke B, Adebayo Rahman S, Olasupo Ogunjuyigbe P. To what extent do community characteristics drive health facility delivery? Findings among Women who had recent live births in Nigeria. *Health & Social Care in the Community*. 2020. <https://doi.org/10.1111/hsc.13133> PMID: 32783320
33. Soltanian A, Davar S, Akhgar MM, Mahjub H, Karami M. Modeling the Factors Affecting the First Birth in the Family's Fertility in Hamedan Province. *Journal of Pharmaceutical Research International*. 2019:1–11.
34. Gebregziabher NK, Zeray AY, Abtey YT, Kinfe TD, Abrha DT. Factors determining choice of place of delivery: analytical cross-sectional study of mothers in Akordet town, Eritrea. *BMC public health*. 2019; 19(1):1–11. <https://doi.org/10.1186/s12889-018-6343-3> PMID: 30606151
35. Yaya S, Bishwajit G, Uthman OA, Amouzou A. Why some women fail to give birth at health facilities: a comparative study between Ethiopia and Nigeria. *PloS one*. 2018; 13(5):e0196896. <https://doi.org/10.1371/journal.pone.0196896> PMID: 29723253
36. Oduaran C, Udomboso CG. Assessing Impacts of Indicators of Child Wealth Index in Africa: A Study of Four Regions in Sub-Saharan Africa. *Commonwealth Youth & Development*. 2019;17(2).
37. Lakew D, Tesfaye D, Mekonnen H. Determinants of stillbirth among women deliveries at Amhara region, Ethiopia. *BMC pregnancy and childbirth*. 2017; 17(1):1–7. <https://doi.org/10.1186/s12884-016-1183-6> PMID: 28049520
38. Colaceci S, Zambri F, Marchetti F, Trivelli G, Rossi E, Petruzzo A, et al. 'A sleeping volcano that could erupt sooner or later'. Lived experiences of women with multiple sclerosis during childbearing age and motherhood: A phenomenological qualitative study. *Multiple Sclerosis and Related Disorders*. 51:102938. <https://doi.org/10.1016/j.msard.2021.102938> PMID: 33882427
39. Olusanya B, Alakija O, Inem V. Non-uptake of facility-based maternity services in an inner-city community in Lagos, Nigeria: an observational study. *Journal of biosocial science*. 2010; 42(3):341–58. <https://doi.org/10.1017/S0021932009990526> PMID: 19951456
40. Muhunzi S, Ngocho JS, Mwanamsangu A, Sanga L, Hiza H, Msuya SE, et al. Prevalence, predictors and reasons for home delivery amongst women of childbearing age in Dodoma Municipality in central Tanzania. *African Health Sciences*. 2020; 20(4):1933–42. <https://doi.org/10.4314/ahs.v20i4.52> PMID: 34394260

41. Ayele G, Tilahun M, Merdikyos B, Animaw W, Taye W. Prevalence and associated factors of home delivery in Arbaminch Zuria district, southern Ethiopia: community based cross sectional study. *Science*. 2015; 3(1):6–9.
42. Dunlop CL, Benova L, Campbell O. Effect of maternal age on facility-based delivery: analysis of first-order births in 34 countries of sub-Saharan Africa using demographic and health survey data. *BMJ open*. 2018; 8(4):e020231. <https://doi.org/10.1136/bmjopen-2017-020231> PMID: 29654024
43. Benova L, Macleod D, Radovich E, Lynch CA, Campbell OM. Should I stay or should I go?: consistency and switching of delivery locations among new mothers in 39 Sub-Saharan African and South/Southeast Asian countries. *Health policy and planning*. 2017; 32(9):1294–308. <https://doi.org/10.1093/heapol/czx087> PMID: 28981668
44. Yaya S, Uthman OA, Okonofua F, Bishwajit G. Decomposing the rural-urban gap in the factors of under-five mortality in sub-Saharan Africa? Evidence from 35 countries. *BMC Public Health*. 2019; 19(1):616. <https://doi.org/10.1186/s12889-019-6940-9> PMID: 31113395
45. Ali AAA, Osman MM, Abbaker AO, Adam I. Use of antenatal care services in Kassala, eastern Sudan. *BMC Pregnancy and Childbirth*. 2010; 10(1):67. <https://doi.org/10.1186/1471-2393-10-67> PMID: 20973972
46. Gayawan E. Spatial analysis of choice of place of delivery in Nigeria. *Sexual & Reproductive Health-care*. 2014; 5(2):59–67. <https://doi.org/10.1016/j.srhc.2014.01.004> PMID: 24814440
47. Adewuyi EO, Khanal V, Zhao Y, David L, Bamidele OD, Auta A. Home childbirth among young mothers aged 15–24 years in Nigeria: a national population-based cross-sectional study. *BMJ open*. 2019; 9(9):e025494. <https://doi.org/10.1136/bmjopen-2018-025494> PMID: 31537553
48. Adewuyi EO, Auta A, Khanal V, Bamidele OD, Akuoko CP, Adefemi K, et al. Prevalence and factors associated with underutilization of antenatal care services in Nigeria: A comparative study of rural and urban residences based on the 2013 Nigeria demographic and health survey. *PLoS one*. 2018; 13(5):e0197324. <https://doi.org/10.1371/journal.pone.0197324> PMID: 29782511
49. Solanke BL, Oladosu OA, Akinlo A, Olanisebe SO. Religion as a social determinant of maternal health care service utilisation in Nigeria. *African Population Studies*. 2015; 29(2).
50. Ganle JK. Why Muslim women in Northern Ghana do not use skilled maternal healthcare services at health facilities: a qualitative study. *BMC International Health and Human Rights*. 2015; 15(1):10.
51. Wagle RR, Sabroe S, Nielsen BB. Socioeconomic and physical distance to the maternity hospital as predictors for place of delivery: an observation study from Nepal. *BMC pregnancy and childbirth*. 2004; 4(1):1–10. <https://doi.org/10.1186/1471-2393-4-1> PMID: 15005809
52. Kitui J, Lewis S, Davey G. Factors influencing place of delivery for women in Kenya: an analysis of the Kenya demographic and health survey, 2008/2009. *BMC pregnancy and childbirth*. 2013; 13(1):1–10. <https://doi.org/10.1186/1471-2393-13-40> PMID: 23414104
53. Afework MF, Admassu K, Mekonnen A, Hagos S, Asegid M, Ahmed S. Effect of an innovative community based health program on maternal health service utilization in north and south central Ethiopia: a community based cross sectional study. *Reproductive Health*. 2014; 11(1):28. <https://doi.org/10.1186/1742-4755-11-28> PMID: 24708848
54. Kanini CM. Utilization of skilled birth attendants among women of reproductive age in Central District, Kitui County: Thesis. *Reproductive Health Of Kenyatta University*; 2012.
55. Babalola S, Fatusi A. Determinants of use of maternal health services in Nigeria—looking beyond individual and household factors. *BMC Pregnancy and Childbirth*. 2009; 9(1):43. <https://doi.org/10.1186/1471-2393-9-43> PMID: 19754941
56. Onah HE, Ikeako LC, Iloabachie GC. Factors associated with the use of maternity services in Enugu, southeastern Nigeria. *Social Science & Medicine*. 2006; 63(7):1870–8. <https://doi.org/10.1016/j.socscimed.2006.04.019> PMID: 16766107
57. Dankwah E, Zeng W, Feng C, Kirychuk S, Farag M. The social determinants of health facility delivery in Ghana. *Reproductive health*. 2019; 16(1):1–10. <https://doi.org/10.1186/s12978-018-0662-9> PMID: 30621726
58. Adebayo AM, Asuzu MC. Utilisation of a community-based health facility in a low-income urban community in Ibadan, Nigeria. *African journal of primary health care & family medicine*. 2015; 7(1):1–8.
59. Rahman M, Curtis SL, Chakraborty N, Jamil K. Women’s television watching and reproductive health behavior in Bangladesh. *SSM-population health*. 2017; 3:525–33. <https://doi.org/10.1016/j.ssmph.2017.06.001> PMID: 29349243
60. Ketemaw A, Tareke M, Dellie E, Sitotaw G, Deressa Y, Tadesse G, et al. Factors associated with institutional delivery in Ethiopia: a cross sectional study. *BMC health services research*. 2020; 20(1):1–6. <https://doi.org/10.1186/s12913-020-05096-7> PMID: 32234043