Harper, Damian, Liefeith, Andreas ORCID logoORCID: https://orcid.org/0000-0003-2121-3219 and Thomas, Aaron (2015) Repeated acceleration and deceleration ability (RADA) in semi-professional soccer players. In: UK Strength and Conditioning Association's 11th Annual Conference, 1-2 August 2015, Chesford Grange Warwickshire, United Kingdom.. (Unpublished)

Downloaded from: https://ray.yorksj.ac.uk/id/eprint/2666/

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. Institutional Repository Policy Statement

# RaY

Research at the University of York St John

For more information please contact RaY at <a href="mailto:ray@yorksi.ac.uk">ray@yorksi.ac.uk</a>

## Repeated acceleration and deceleration ability (RADA) in male semi-professional soccer players


Harper, D.J., Liefeith, A. & Thomas, A. Department of Applied Human Sciences, York St John University, York, England

## **Purpose**

Contemporary developments in soccer match play demands players to execute more frequent bouts of short explosive sprints, and consequently be able to control for sudden and rapid reductions in velocity (Barnes *et al.*, 2014). In fact recent research has highlighted the prevalence and importance of executing repeated rapid accelerations (ACC) and decelerations (DEC) (Akenhead *et al.*, 2014; Russell *et al.*, 2015). A reduced proficiency to maintain repeated rapid ACC's and DEC's could have significant performance and injury implications. Therefore, the aims of this study was to develop a new repeated ACC and DEC test that could be used to evaluate the repeated ACC and DEC ability (RADA) in soccer players, but could also be applied to other team sports that demand repeated bouts of ACC and DEC.

#### Methods

10 male semi-professional soccer players (age 23.06 ± 6.08years, mass 76.86 ± 8.13kg, stature 1.78 ± 0.04m) completed the RADA test. RADA test consisted of a 20m maximal ACC (with 10m-split time) followed by a 5m rapid DEC that proceeded a 3m backpedal, prior to a final 8m re-ACC. This sequence was completed 6 times with a 30s active recovery period (figure 1). Split times at 10, 20, 22 and 25m were used for final analysis and recorded using dual-beam timing gates (Smart Speed, Fusion Sport, Australia). Average velocity and ACC (positive and negative) for each section of the RADA test was calculated and test performance evaluated with best, mean and percentage decrement (Fitzimmons et al., 1993). Pearson's coefficients were also used to investigate the association between distinct aspects of the RADA test.



### Results

**Table 1.** Descriptive data (mean ± standard deviation (SD); 90% confidence interval [CI]) for each section of the RADA

RADA test variables	Mean ± SD	90% CI	
ACC performance variables			
10mACC <sub>best</sub> (s)	$1.87 \pm 0.08$	1.74-2.00	
10mACC <sub>best</sub> (m.s <sup>-2</sup> )	2.87 ± 0.27	2.43-3.31	
10mACC <sub>mean</sub> (s)	1.96 ± 0.05	1.88-2.04	
$10 \text{mACC}_{\text{mean}} (\text{m.s}^{-2})$	2.61 ± 0.15	2.36-2.86	
10mACC <sub>%dec</sub> (%)	8.54 ± 5.82	0-18.11	
20mACC <sub>best</sub> (s)	$3.28 \pm 0.09$	3.13-3.43	
$20 \text{mACC}_{\text{best}} (\text{m.s}^{-2})$	$2.20 \pm 0.09$	2.05-2.35	
20mACC <sub>mean</sub> (s)	$3.37 \pm 0.09$	3.13-3.43	
$20 \text{mACC}_{\text{mean}} (\text{m.s}^{-2})$	2.11 ± 0.11	1.93-2.29	
20mACC <sub>%dec</sub> (%)	3.88 ± 1.78	0.95-6.81	
DEC performance variables			
5mDEC <sub>best</sub> (s)	1.16 ± 0.08	1.03-1.29	
$5mDEC_{best} (m.s^{-2})$	-5.13 ± 0.55	4.23-6.03	
5mDEC <sub>mean</sub> (s)	$1.26 \pm 0.08$	1.13-1.39	
$5mDEC_{mean} (m.s^{-2})$	-4.60 ± 0.40	3.94-5.26	
5mDEC <sub>%dec</sub> (%)	$10.02 \pm 5.13$	1.58-18.46	
3mDEC <sub>best</sub> (s)	$0.82 \pm 0.07$	0.70-0.94	
$3mDEC_{best} (m.s^{-2})$	-4.61 ± 0.83	3.24-5.98	
$3mDEC_{mean}(s)$	$0.91 \pm 0.08$	0.78-1.04	
$3mDEC_{mean}(m.s^{-2})$	-3.79 ± 0.65	2.72-4.86	
3mDEC <sub>%dec</sub> (%)	17.71 ± 6.31	7.33-28.09	
RADA performance variables			
RADA <sub>best</sub> (s)	4.52 ± 0.10	4.36-4.68	
RADA <sub>mean</sub> (s)	$4.63 \pm 0.10$	4.47-4.79	
RADA <sub>%dec</sub> (%)	2.28 ± 0.98	0.67-3.89	

**Table 2.** Best predictors of RADA. r = correlation coefficient (\*\* = P < 0.001, \* = P < 0.05),  $r^2 = coefficient$  of determination, descriptor = magnitude of correlation interpreted using Hopkins (2000) scale (<0.1 trivial, 0.1-0.3 small, 0.3-0.5 moderate, 0.5-0.7 large, 0.7-0.9 very large, 0.9-1.0 perfect).

Best Predictors of RADA	r	r²	Descriptor
RADA <sub>%dec</sub> DEC <sub>3%dec</sub>	0.780**	61%	Very Large
RADA <sub>best</sub> DEC <sub>3best</sub>	0.644*	41%	Large
ACC <sub>10%dec</sub> ACC <sub>10best</sub>	0.789**	62%	Very Large

## **Conclusions**

RADA<sub>decrement</sub> was significantly correlated to the  $DEC_{3mean}$  ability. A reduction in DEC ability is likely to have huge impact on ability to execute repeated high intensity bouts of ACC's and DEC's, indicating DEC's seem to be highly sensitive to fatigue. Interestingly  $DEC_{3best}$  was also significantly correlated to  $RADA_{best}$ , reiterating that DEC is critical to overall performance.  $ACC_{10best}$  significantly correlated to  $ACC_{10decrement}$ , suggesting players that can produce high ACC have less ability to maintain this quality when repeated ACC's are required. Small correlations between all ACC and DEC abilities seem to suggest ACC and DEC are distinct movement skills therefore requiring unique training interventions. Further investigation should focus on identifying physical and technical qualities required for proficient RADA in particular the DEC component, and to determine the validity, reliability and sensitivity of this test.

#### References

Akenhead, R., Hayes, P. R., Thompson, K. G., & French, D. (2013). Diminutions of acceleration and deceleration output during professional football match play. *Journal of Science and Medicine in Sport / Sports Medicine Australia*, *16*(6), 556–61.

Barnes, C., Archer, D. T., Hogg, B., Bush, M., & Bradley, P. S. (2014). The Evolution of Physical and Technical Performance Parameters in the English Premier League. *International Journal of Sports Medicine*, *35*, 1095–1100.

Fitzsimmons, M., Dawson, B., Ward, D., & Wilkinson, A. (1993). Cycling and running tests of repeated sprint ability. Australian Journal of Science and Medicine in Sport, 25, 82-87.

Hopkins, WG. (2000) A new view of statistics. Available at :http://www.sportsci.org/resource/stats/effectmag.html

Russell, M., Sparkes, W., Northeast, J., Cook, C. J., Love, T. D., Bracken, R. M., & Kilduff, L. P. (2014). Changes in acceleration and deceleration capacity throughout professional soccer match-play. *Journal of Strength and Conditioning Research*. In press.