

Est.  
1841

YORK  
ST JOHN  
UNIVERSITY

Schrodt, Franziska, Bailey, Joseph J, Rijdsdijk, Kenneth F., Kissling, W. Daniel, Seijmonsbergen, Arie C., Van Ree, Derk, Hjort, Jan, Lawley, Russell S., Williams, Christopher N., Anderson, Mark G., Beier, Paul, Van Beukering, Pieter, Boyd, Doreen S., Brilha, Jose, Carcavilla, Luis, Dahlin, Kyla M., Gill, Joel C., Gordon, John E., Gray, Murray, Grundy, Mike, Hunter, Malcolm M., Lawler, Joshua J., Monge-Ganuzas, Manu, Royse, Katherine R., Stewart, Iain, Record, Sydne, Turner, Woody, Zarnetske, Phoebe L. and Field, Richard (2019) Opinion: To advance sustainable stewardship, we must document not only biodiversity but geodiversity. Proceedings of the National Academy of Sciences (PNAS, USA), 116 (33). pp. 16155-16158.

Downloaded from: <https://ray.yorks.ac.uk/id/eprint/3999/>

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version:

<https://www.pnas.org/content/116/33/16155>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repositories Policy Statement](#)

# RaY

Research at the University of York St John

For more information please contact RaY at  
[ray@yorks.ac.uk](mailto:ray@yorks.ac.uk)



# To advance sustainable stewardship, we must document not only biodiversity but geodiversity

Franziska Schrodtr<sup>a,1</sup>, Joseph J. Bailey<sup>b</sup>, W. Daniel Kissling<sup>c</sup>, Kenneth F. Rijdsdijk<sup>c</sup>, Arie C. Seijmonsbergen<sup>c</sup>, Derk van Ree<sup>d,e</sup>, Jan Hjort<sup>f</sup>, Russell S. Lawley<sup>g</sup>, Christopher N. Williams<sup>g</sup>, Mark G. Anderson<sup>h</sup>, Paul Beier<sup>i</sup>, Pieter van Beukering<sup>o</sup>, Doreen S. Boyd<sup>a</sup>, José Brilha<sup>j</sup>, Luis Carcavilla<sup>k</sup>, Kyla M. Dahlin<sup>l</sup>, Joel C. Gill<sup>m</sup>, John E. Gordon<sup>n</sup>, Murray Gray<sup>o</sup>, Mike Grundy<sup>p</sup>, Malcolm L. Hunter<sup>q</sup>, Joshua J. Lawler<sup>r</sup>, Manu Monge-Ganuzas<sup>s</sup>, Katherine R. Royse<sup>g</sup>, Iain Stewart<sup>t</sup>, Sydne Record<sup>u</sup>, Woody Turner<sup>v</sup>, Phoebe L. Zarnetske<sup>w</sup>, and Richard Field<sup>a</sup>

Rapid environmental change is driving the need for complex and comprehensive scientific information that supports policies aimed at managing natural resources through international treaties, platforms, and networks. One successful approach for delivering such information has been the development of essential variables for climate (1), oceans (2), biodiversity (3), and sustainable development goals (4) (ECVs, EOVs, EBVs, and ESDGVs, respectively). These efforts have improved consensus on terminology and identified essential sets of measurements for characterizing and monitoring changes on our planet. In doing so, they have advanced science and informed policy. As an important but largely unanticipated consequence, conceptualizing these variables has also given rise to discussions regarding data discovery, data access, and governance of research infrastructures. Such discussions are vital to ensure effective storage, distribution, and use of data among management agencies, researchers, and policymakers (5, 6).

Although the current essential variables frameworks account for the biosphere, atmosphere, and some aspects of the hydrosphere (1–4), they largely overlook geodiversity—the variety of abiotic features



**Mining is one example of the human impact on geodiversity. Active mines cause a decrease in local biodiversity, but in some cases they can provide an important habitat for specialized and rare species after the mine has been abandoned. Image credit: Shutterstock/1968.**

and processes of the land surface and subsurface (7). Analogous to biodiversity, geodiversity is important for the maintenance of ecosystem functioning and services (8), and areas high in geodiversity have been

<sup>a</sup>School of Geography, University of Nottingham, Nottingham, NG7 2RD, United Kingdom; <sup>b</sup>School of Humanities, Religion and Philosophy, York St John University, YO31 7EX, United Kingdom; <sup>c</sup>Department of Theoretical and Computational Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, 1090 GE Amsterdam, Netherlands; <sup>d</sup>Unit Geo-engineering, Deltares, 2600 MH Delft, Netherlands; <sup>e</sup>Faculty of Science, Institute for Environmental Studies, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, Netherlands; <sup>f</sup>Geography Research Unit, University of Oulu, 90014, Oulu, Finland; <sup>g</sup>GeoAnalytics and Modelling Directorate, British Geological Survey, Nottingham, NG12 5GG, United Kingdom; <sup>h</sup>The Nature Conservancy, Boston, MA 02111; <sup>i</sup>School of Forestry, Northern Arizona University, Flagstaff, AZ 86011-5018; <sup>j</sup>Institute of Earth Sciences, Pole of the University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal; <sup>k</sup>Geological and Mining Institute of Spain, 28003 Madrid, Spain; <sup>l</sup>Department of Geography, Environment, and Spatial Sciences, Michigan State University, East Lansing, MI 48824; <sup>m</sup>British Geological Survey, Environmental Science Centre, Nottingham, NG12 5GG, United Kingdom; <sup>n</sup>School of Geography and Sustainable Development, University of St Andrews, St Andrews, Scotland KY16 9AL, United Kingdom; <sup>o</sup>School of Geography, Queen Mary University of London, London, E1 4NS, United Kingdom; <sup>p</sup>Agriculture and Food, CSIRO, St Lucia, Queensland 4067, Australia; <sup>q</sup>Department of Wildlife, Fisheries, and Conservation Biology, University of Maine, Orono, ME 04469; <sup>r</sup>School of Environmental and Forest Sciences, University of Washington, Seattle, WA 98195; <sup>s</sup>Service of Urdaibai Biosphere Reserve, Basque Government, 48011 Bilbao, Spain; <sup>t</sup>School of Geography, Earth, and Environmental Sciences, Plymouth University, Plymouth, PL4 8AA, United Kingdom; <sup>u</sup>Department of Biology, Bryn Mawr College, Bryn Mawr, PA 19010; <sup>v</sup>Earth Science Division, NASA Headquarters, Washington, DC, 20546; and <sup>w</sup>Department of Forestry, Department of Fisheries and Wildlife, Michigan State University, East Lansing, MI 48824

Author contributions: F.S., J.J.B., W.D.K., K.F.R., A.C.S., D.v.R., J.H., R.S.L., C.N.W., M.G.A., P.B., P.v.B., and R.F. designed research; F.S. performed research; F.S. and J.C.G. analyzed data; F.S., J.J.B., W.D.K., K.F.R., A.C.S., D.v.R., J.H., R.S.L., C.N.W., M.G.A., P.B., P.v.B., D.S.B., J.B., L.C., K.M.D., J.C.G., J.E.G., M. Gray, M. Grundy, M.L.H., J.J.L., M.M.-G., K.R.R., I.S., S.R., W.T., P.L.Z., and R.F. wrote the paper. F.S., J.J.B., and R.F. conceived the idea with major contributions by W.D.K., K.F.R., A.C.S., D.v.R., J.H., and R.S.L. as well as further input by all co-authors for conceptualization and methodology. F.S. led the writing with major contributions by J.J.B. and R.F., as well as further contributions by all other co-authors. F.S. designed and made Fig. 1; J.C.G. designed and made Table S1, and both received contributions from all co-authors.

The authors declare no conflict of interest.

Published under the PNAS license.

Any opinions, findings, conclusions, or recommendations expressed in this work are those of the authors and have not been endorsed by the National Academy of Sciences.

<sup>1</sup>To whom correspondence may be addressed. Email: Franziska.Schrodtr1@nottingham.ac.uk.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1911799116/-/DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1911799116/-/DCSupplemental).



Meanwhile, extraction and storage of vast quantities of soil and rock that are byproducts from mining metals, as well as associated land requirements, are currently not part of integrated broad-scale management frameworks (16).

### Essential Variables

In presenting the EGV concept, we aim to 1) complement and augment existing essential variables (ECVs, EOVs, EBVs, and ESDGVs) (Fig. 1), 2) improve global coordination of monitoring strategies, and 3) advance communication between policymakers and geoscientists. To achieve these goals, we propose a framework for policymakers and researchers to guide future definitions of relevant measurements that capture the key elements of geodiversity. We define EGVs as abiotic state and process variables related to geology, geomorphology, soils, and hydrology 1) relevant to natural resource management and human well-being, conservation, or ecology; 2) complementary to (and not duplicating) the other suites of essential variables; and 3) feasible and cost effective to measure (Fig. 1). In Table S2, we propose a candidate set of variables.

Some aspects of EGVs are already used by international conservation organizations; these provide a solid basis for further integrating EGVs into global treaties and international conventions. For example, the International Union for Conservation of Nature (IUCN) refers to the relevance of geodiversity for the conservation of natural resources within three resolutions titled “Conservation of Geodiversity and Geological Heritage,” “Valuing and Conserving Geoheritage Within the IUCN Programme,” and “Conservation of Moveable Geological Heritage.” Furthermore, the United Nations Educational, Scientific and Cultural Organization recognizes the outstanding universal value of geodiversity elements with their inclusion in both the World Heritage List (in May 2019, 95 properties in 53 countries worldwide) and in the Global Geoparks Network (140 Geoparks in 40 countries, as of May 2019). Many protected areas have the preservation of geodiversity and geoheritage as a goal of their management planning, including the Spanish network of Biosphere Reserves, Australia’s New South Wales National Parks, and the US National Park Service. Meanwhile, the Food and Agriculture Organization coordinates a Global Soil Partnership that seeks to monitor the state of global soils and improve the governance and effectiveness of soil information.

Data and information products to measure changes in EGVs at management-relevant timescales are increasingly available and sometimes linked to global observatories, such as the Global Earth Observation System of Systems, with its Societal Benefit Areas (SBAs). However, these mainly cover natural hazards such as floods, earthquakes, and landslides (e.g., SBA disaster resilience). Where dangers are more diffuse or related to natural resource use, EGVs are not yet available (e.g., relating to global sand extraction and domestication of soil resources).

### Making EGVs Work

Overall, despite the clear global importance of geodiversity, very limited international efforts have been devoted to developing measures that support decision making for supranational and global policy targets and SDGs [although there have been efforts to do so in the past (17)]. Geodiversity is highly relevant, for example, to the IUCN World Parks Congress, the World Conservation Congress, the Convention on Biological Diversity’s Aichi targets, SDGs, and the Sendai Framework for Disaster Risk Reduction (Table S2). In Table S1, we specifically link EGVs to the 17 SDGs and four Sendai Framework priorities.

We advocate a holistic approach that recognizes and tracks the integrity of the abiotic and biotic components of geosystems and ecosystems as the most effective means to address global environmental challenges. Following the examples of the ECV, EOV,

**Attaining a sustainable circular economy and safeguarding our natural resources, while also accounting for population growth, further urbanization, and improved well-being, will require international consideration of material flows and their impacts across terrestrial and aquatic systems globally.**

and EBV communities, we recommend collaborative development of comprehensive and interoperable databases of geodiversity globally, following common protocols, a standardized terminology (e.g., controlled vocabularies), and a consistent metadata reporting. We further recommend forming an expert panel, for example within the Group on Earth Observation framework, to further develop the conceptual framework of EGVs. Finally, we encourage better communication with policymakers about the importance of considering EGVs in international conventions and policy documents. This could be enhanced by applying a “geosystem services” concept, which would complement the successful ecosystem services concept whose use within a policy and international treaties context was advanced by the Millennium Ecosystem Assessment (11). Better communication would also be enhanced by applying the recently published International Panel on Climate Change communication recommendations.

We now have the technical capacity and experience from other scientific communities to describe abiotic characteristics of Earth’s surface and subsurface and to develop holistic and parsimonious measures of geosystem and ecosystem structure, function, and risks. Attaining a sustainable circular economy and safeguarding our natural resources, while also accounting for population growth, further urbanization, and improved well-being, will require international consideration of material flows and their impacts across terrestrial and aquatic systems globally. This will entrench a more holistic approach to nature, improving our efforts to designate protected areas and enhance the management of

natural resources. Doing so is essential for safeguarding biodiversity, geodiversity, ecosystem, and geosystem services in a rapidly changing world and for integrating and balancing the economic, social, and environmental dimensions of sustainable development.

## Acknowledgments

F.S. acknowledges support by the University of Nottingham Anne McLaren fellowship; J.H. received funding from the Academy of Finland project 315519; W.D.K. acknowledges support from the University of Amsterdam Faculty Research Cluster “Global Ecology.”

- 1 S. Bojinski et al., The concept of essential climate variables in support of climate research, applications, and policy. *Bull. Am. Meteorol. Soc.* **95**, 1431–1443 (2014).
- 2 A. J. Constable et al., Developing priority variables (“ecosystem Essential Ocean Variables”—eEOVs) for observing dynamics and change in Southern Ocean ecosystems. *J. Mar. Syst.* **161**, 26–41 (2016).
- 3 H. M. Pereira et al., Ecology. Essential biodiversity variables. *Science* **339**, 277–278 (2013).
- 4 B. Reyers, M. Stafford-Smith, K. H. Erb, R. J. Scholes, O. Selomane, Essential variables help to focus sustainable development goals monitoring. *Curr. Opin. Environ. Sustain.* **26**, 97–105 (2017).
- 5 W. D. Kissling et al., Towards global interoperability for supporting biodiversity research on essential biodiversity variables (EBVs). *Biodiversity* **16**, 99–107 (2015).
- 6 M. D. Wilkinson et al., The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data* **3**, 160018 (2016).
- 7 M. Gray, *Geodiversity. Valuing and Conserving Abiotic Nature* (Wiley, London), ed. 2, 2013).
- 8 J. Brilha, M. Gray, D. I. Pereira, P. Pereira, Geodiversity: An integrative review as a contribution to the sustainable management of the whole of nature. *Environ. Sci. Policy* **86**, 19–28 (2018).
- 9 J. J. Bailey, D. S. Boyd, J. Hjort, C. P. Lavers, R. Field, Modelling native and alien vascular plant species richness: At which scales is geodiversity most relevant? *Glob. Ecol. Biogeogr.* **26**, 763–776 (2017).
- 10 J. Hjort, J. E. Gordon, M. Gray, M. L. Hunter, Jr, Why geodiversity matters in valuing nature’s stage. *Conserv. Biol.* **29**, 630–639 (2015).
- 11 C. C. D. F. Van Ree, P. J. H. van Beukering, Geosystem services: A concept in support of sustainable development of the subsurface. *Ecosyst. Serv.* **20**, 30–36 (2016).
- 12 E. Nickless, Delivering sustainable development goals: The need for a new international resource governance framework. *Ann. Geophys.* **60**, 1–6 (2017).
- 13 T. Markus, “The Alpine Convention’s Soil Conservation Protocol: A model regime?” in *International Yearbook of Soil Law and Policy*, H. Ginzky, I. Heuse, T. Qin, O. Ruppel, P. Wegerdt, Eds. (Springer, Cambridge, 2016), pp. 149–164.
- 14 M. Bendixen et al., Promises and perils of sand exploitation in Greenland. *Nat. Sustain.* **2**, 98–104 (2019).
- 15 M. L. C. M. Henckens, C. M. J. Ryngaert, P. P. J. Driessen, E. Worrell, Normative principles and the sustainable use of geologically scarce mineral resources. *Resour. Policy* **59**, 351–359 (2018).
- 16 A. Alaoma, N. Voulvoulis, Mineral resource active regions: The need for systems thinking in management. *AIMS Environ. Sci.* **5**, 78–95 (2018).
- 17 A. R. Berger, Assessing rapid environmental change using geoindicators. *Environ. Geol.* **32**, 36–44 (1997).