

Schrodt, Franziska, Bailey, Joseph J ORCID logoORCID: https://orcid.org/0000-0002-9526-7095, Rijsdijk, Kenneth F., Kissling, W. Daniel, Seijmonsbergen, Arie C., Van Ree, Derk, Hjort, Jan, Lawley, Russell S., Williams, Christopher N., Anderson, Mark G., Beier, Paul, Van Beukering, Pieter, Boyd, Doreen S., Brilha, Jose, Carcavilla, Luis, Dahlin, Kyla M., Gill, Joel C., Gordon, John E., Gray, Murray, Grundy, Mike, Hunter, Malcolm M., Lawler, Joshua J., Monge-Ganuzas, Manu, Royse, Katherine R., Stewart, Iain, Record, Sydne, Turner, Woody, Zarnetske, Phoebe L. and Field, Richard (2019) Opinion: To advance sustainable stewardship, we must document not only biodiversity but geodiversity. Proceedings of the National Academy of Sciences (PNAS, USA), 116 (33). pp. 16155-16158.

Downloaded from: https://ray.yorksj.ac.uk/id/eprint/3999/

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version: https://www.pnas.org/content/116/33/16155

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. Institutional Repositories Policy Statement

RaY

Research at the University of York St John
For more information please contact RaY at ray@yorksj.ac.uk

Opinion: To advance sustainable stewardship, we must document not only biodiversity but geodiversity

Authors: Franziska Schrodt¹, Joseph J. Bailey², W. Daniel Kissling³, Kenneth F. Rijsdijk³, Arie C. Seijmonsbergen³, Derk van Ree^{4,9}, Jan Hjort⁵, Russell S. Lawley⁶, Christopher N. Williams⁶, Mark Anderson⁷, Paul Beier⁸, Pieter van Beukering⁹, Doreen S. Boyd¹, José Brilha¹⁰, Luis Carcavilla¹¹, Kyla M. Dahlin¹², Joel C. Gill¹³, John E. Gordon¹⁴, Murray Gray¹⁵, Mike Grundy¹⁶, Malcolm L. Hunter¹⁷, Joshua J. Lawler¹⁸, Manu Monge-Ganuzas¹⁹, Katherine R. Royse⁶, Iain Stewart²⁰, Sydne Record²¹, Woody Turner²², Phoebe L. Zarnetske²³, Richard Field¹

Rapid environmental change is driving the need for complex and comprehensive scientific information that supports policies aimed at managing natural resources through international treaties, platforms, and networks. One successful approach for delivering such information has been the development of Essential Variables for climate (1), oceans (2), biodiversity (3), and sustainable development goals (4) (ECVs, EOVs, EBVs, and ESDGVs, respectively). These efforts have improved consensus on terminology, and identified essential sets of measurements for characterizing and monitoring changes on our planet. In doing so, they have advanced science and informed policy. As an important but largely unanticipated consequence, conceptualizing these variables has also given rise to discussions regarding data discovery, data access, and governance of research infrastructures. Such discussions are vital to ensure effective storage, distribution, and use of data among management agencies, scientists, and policymakers (5, 6).

¹ School of Geography, University of Nottingham, Nottingham, NG7 2RD, UK

² Geography, School of Humanities, Religion and Philosophy, York St John University, YO31 7EX, UK

³ Department Theoretical and Computational Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, P.O. Box 94240, 1090 GE Amsterdam, Netherlands

⁴ Unit Geo-engineering, Deltares, 2600 MH Delft, Netherlands

⁵ Geography Research Unit, University of Oulu, P.O. Box 3000, 90014, Oulu, Finland

⁶ Geo-Analytics and Modelling Directorate, British Geological Survey, Nottingham, NG12 5GG, UK

⁷ The Nature Conservancy, Boston, MA, USA

⁸ School of Forestry, Northern Arizona University, Flagstaff AZ 86011-5018, USA

⁹ Faculty of Science, Institute for Environmental Studies, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

¹⁰ Institute of Earth Sciences, Pole of the University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal

¹¹ Geological and Mining Institute of Spain, Madrid, Spain

¹² Department of Geography, Environment, and Spatial Sciences, Michigan State University, East Lansing, MI, USA

¹³ British Geological Survey, Environmental Science Centre, Nottingham, NG12 5GG, UK

¹⁴ School of Geography and Sustainable Development, University of St Andrews, St Andrews, Scotland KY16 9AL, UK

¹⁵ School of Geography, Queen Mary University of London, London E1 4NS, UK

¹⁶ Agriculture and Food, CSIRO, St Lucia, Qld 4067, Australia

¹⁷ Department of Wildlife, Fisheries, and Conservation Biology, University of Maine, Orono, ME, USA

¹⁸ School of Environmental and Forest Sciences, University of Washington, USA

¹⁹ Service of Urdaibai Biosphere Reserve, Basque Government, Spain

²⁰ School of Geography, Earth, & Environmental Sciences, Plymouth University, Plymouth PL4 8AA, UK

²¹ Department of Biology, Bryn Mawr College, Bryn Mawr, 19010, USA

²² Earth Science Division, NASA Headquarters, Washington, DC, USA

²³ Department of Forestry, Department of Fisheries and Wildlife, Michigan State University, East Lansing, MI, USA

Although the current Essential Variables frameworks account for the biosphere, atmosphere, and some aspects of the hydrosphere (1–4), they largely overlook geodiversity—the variety of abiotic features and processes of the land surface and subsurface (7) (fig. 1). Analogous to biodiversity, geodiversity is important for the maintenance of ecosystem functioning and services (8), and areas high in geodiversity have been shown to support high biodiversity (9). Thus, consideration of geodiversity is an important part of developing nature-based solutions to global environmental challenges and demands for natural resources, particularly in relation to human wellbeing and ecosystem functioning.

And yet, despite many facets of sustainable development being underpinned by access to geological assets, key elements of geodiversity are yet to be incorporated into policy documents and international conventions. We therefore propose Essential Geodiversity Variables (EGVs) describing features and processes of Earth's abiotic surface and subsurface, to advance science and sustainable stewardship, complementing the existing Essential Variables. These EGVs will enable more holistic and better-informed monitoring efforts, decision making, and responses to global change.

Broad Scope

The scope of geodiversity covers a wide range of policy areas, including terrestrial and marine conservation, sustainable use of natural resources, public health, natural hazard management, recreation, and tourism (e.g., see the "Conserving Nature's Stage" (10) and geosystem services (11) concepts). For example, abiotic features, including geothermal springs, inspired the creation of the world's first national park, Yellowstone. This park aimed specifically to safeguard geodiversity, and a century later its geothermal springs were the discovery site for *Thermus aquaticus*, a bacterium containing a thermostable enzyme that is used to amplify DNA segments and is thus the foundation of modern gene technology.

Another illustration of the critical importance of understanding and monitoring geodiversity globally concerns resource extraction. Removal of natural resources can decrease geological or mineral diversity, negatively impact local ecosystems (due to toxic extraction methods), and conflict with human rights. Notably, mobile phones with touch screens contain 54 elements of the periodic table, many of which are unevenly distributed in nature around the world and thus represent resource security concerns. Continued resource extraction is essential for achieving the Sustainable Development Goals (SDGs), but trade-offs with biodiversity conservation and human rights need to be explicitly addressed. For example, the transition to renewable energy sources will require the extraction of new minerals (e.g. materials for solar panels), as will greater urbanization and fertilizers for enhanced food security (12). Some tools and concepts necessary for incorporating such trade-offs are already available but not commonly applied within the context of the SDGs, e.g. in form of the geosystem services approach (11).

And while some EGVs such as groundwater are already considered within international legislation (e.g. the EU Groundwater Directive), most are underrepresented or not effective over wider regions. For example, currently the only binding international convention specifically on soil conservation is the 1991 Alpine Convention's Soil Conservation Protocol (13) (fig. 1), which omits most of the Earth's surface. Similarly, extraction of sand—a key ingredient in building materials and electronics—remains largely unregulated, despite rising global demand for this finite resource and wide-ranging devastating environmental consequences resulting from its extraction (14). The demand for minerals is rising globally, yet their extraction lacks international governance (15). Meanwhile, extraction and storage of vast quantities of soil and rock that are by-products from mining metals, as well as associated land requirements, are currently not part of integrated broad-scale management frameworks (16).

Essential Variables

In presenting the EGV concept, we aim to 1) complement and augment existing essential variables (ECVs, EOVs, EBVs and ESDGVs) (fig. 1); 2) improve global coordination of monitoring strategies; and 3)

advance communication between policymakers and geoscientists. To achieve these goals, we propose a framework for policymakers and scientists to guide future definitions of relevant measurements that capture the key elements of geodiversity. We define EGVs as abiotic state and process variables related to geology, geomorphology, soils, and hydrology (i) relevant to natural resource management and human wellbeing, conservation, or ecology; (ii) complementary to (and not duplicating) the other suites of Essential Variables; and (iii) feasible and cost effective to measure (fig. 1, table S2).

Some aspects of EGVs are already used by international conservation organizations; these provide a solid basis for further integrating EGVs into global treaties and international conventions. For example, the International Union for Conservation of Nature (IUCN) refers to the relevance of geodiversity for conservation of natural resources within three resolutions titled "Conservation of geodiversity and geological heritage," "Valuing and conserving geoheritage within the IUCN Programme," and "Conservation of moveable geological heritage." Furthermore, the United Nations Educational, Scientific and Cultural Organization (UNESCO) recognizes the outstanding universal value of geodiversity elements with their inclusion in both the World Heritage List (in May 2019, 95 properties in 53 countries worldwide) and in the Global Geoparks Network (140 Geoparks in 40 countries, as of May 2019). Many protected areas have the preservation of geodiversity and geoheritage as a goal of their management planning, including the Spanish network of Biosphere Reserves, Australia's New South Wales National Parks, and the U.S. National Park Service. Meanwhile, the Food and Agriculture Organization (FAO) coordinates a Global Soil Partnership that seeks to monitor the state of global soils and improve the governance and effectiveness of soil information.

Data and information products to measure changes in EGVs at management-relevant timescales are increasingly available and sometimes linked to global observatories, such as the Global Earth Observation System of Systems (GEOSS), with its Societal Benefit Areas (SBAs). However, these mainly cover natural hazards such as floods, earthquakes, and landslides (e.g. SBA disaster resilience). Where dangers are more diffuse or related to natural resource use, EGVs are not yet available (e.g. relating to global sand extraction and domestication of soil resources).

Making EGVs Work

Overall, despite the clear global importance of geodiversity, very limited international efforts have been devoted to developing measures that support decision-making for supra-national and global policy targets and SDGs (though there have been efforts to do so in the past (17)). Geodiversity is highly relevant, for example, to the IUCN World Parks Congress, the World Conservation Congress, the Convention on Biological Diversity's Aichi targets, SDGs and the Sendai Framework for Disaster Risk Reduction (also see table S1 and table S2 in the supplementary materials).

We advocate a holistic approach that recognizes and tracks the integrity of the abiotic and biotic components of geo- and ecosystems, as the most effective means to address global environmental challenges. Following the examples of Essential Variables for the climate (ECV), ocean (EOV), and biodiversity (EBV) communities, we recommend collaborative development of comprehensive and interoperable databases of geodiversity globally, following common protocols, a standardized terminology (e.g. controlled vocabularies) and a consistent metadata reporting. We further recommend forming an expert panel, for example within the Group on Earth Observation framework to further develop the conceptual framework of EGVs. Finally, we encourage better communication with policymakers about the importance of considering EGVs in international conventions and policy documents. This could be enhanced by applying a "geosystem services" concept, which would complement the successful ecosystem services concept whose use within a policy and international treaties context was advanced by the Millennium Ecosystem Assessment (11), and by applying the recently published International Panel on Climate Change (IPCC) communication recommendations.

We now have the technical capacity and experience from other scientific communities to describe abiotic characteristics of Earth's surface and subsurface, and to develop holistic and parsimonious measures of geo-and ecosystem structure, function, and risks. Attaining a sustainable circular economy and safeguarding our natural resources, while also accounting for population growth, further urbanization, and improved well-being, will require international consideration of material flows and their impacts across terrestrial and aquatic systems globally. This will entrench a more holistic approach to nature, improving our efforts to designate protected areas and enhance management of natural resources. Doing so is essential for safeguarding biodiversity, geodiversity, ecosystem and geosystem services in a rapidly changing world, and for integrating and balancing the economic, social, and environmental dimensions of sustainable development.

CAPTIONS

Pic 1:

Mining is one example of the human impact on geodiversity. Active mines cause a decrease in local biodiversity, which then typically blossoms again after the mine's been abandoned. Image credit: Shutterstock/1968

Pic 2:

Figure 1: Schematic proportions of the Earth covered by existing Essential Biodiversity Variables (EBV, green), Essential Climate Variables (ECV, light blue), Essential Ocean Variables (EOV, dark blue), and by our proposed Essential Geodiversity Variables (EGVs, orange). Although life occurs throughout the ocean environment, EOVs refer predominately to abiotic aspects such as ocean physics and biogeochemistry, which do not overlap with EBVs (by definition exclusively covering biotic aspects). Consequently, the EBV box does not extend across the whole Earth surface cover axis. Some Essential Variables do overlap, as indicated by the striped sections—e.g. zooplankton diversity is both an EBV and EOV, while surface water is both an ECV and EGV. Several major international conventions (right) monitor and assess networks associated with each Essential Variable concept.

References and Notes

- 1. S. Bojinski et al., The concept of essential climate variables in support of climate research, applications, and policy. Bull. Am. Meteorol. Soc. 95, 1431–1443 (2014).
- 2. A. J. Constable et al., Developing priority variables ("ecosystem Essential Ocean Variables" eEOVs) for observing dynamics and change in Southern Ocean ecosystems. J. Mar. Syst. 161, 26–41 (2016).
- 3. H. M. Pereira et al., Essential biodiversity variables. Science. 339, 277–278 (2013).
- 4. B. Reyers, M. Stafford-Smith, K. H. Erb, R. J. Scholes, O. Selomane, Essential Variables help to focus Sustainable Development Goals monitoring. Curr. Opin. Environ. Sustain. 26, 97–105 (2017).
- 5. W. D. Kissling et al., Towards global interoperability for supporting biodiversity research on essential biodiversity variables (EBVs). Biodiversity. 16, 99–107 (2015).
- 6. M. D. Wilkinson et al., The FAIR Guiding Principles for scientific data management and stewardship. Sci. Data. 3, 160018 (2016).
- 7. M. Gray, Geodiversity. Valuing and Conserving Abiotic Nature (Wiley, London, ed. 2, 2013).
- 8. J. Brilha, M. Gray, D. I. Pereira, P. Pereira, Geodiversity: An integrative review as a contribution to the sustainable management of the whole of nature. Environmental Science & Policy 86, 19-28 (2018).

- 9. J. J. Bailey, D. S. Boyd, J. Hjort, C. P. Lavers, R. Field, Modelling native and alien vascular plant species richness: At which scales is geodiversity most relevant? Global Ecology and Biogeography 26, 763–776 (2017).
- 10. J. Hjort, J. E. Gordon, M. Gray, M. L. Hunter, Why geodiversity matters in valuing nature's stage. Conserv. Biol. 29, 630–639 (2015).
- 11. C. C. D. F. Van Ree, P. J. H. van Beukering, Geosystem services: A concept in support of sustainable development of the subsurface. Ecosyst. Serv. 20, 30–36 (2016).
- 12. E. Nickless, Delivering Sustainable Development Goals: The Need for a New International Resource Governance Framework. Ann. Geophys. 60, 1–6 (2017).
- 13. T. Markus, in International Yearbook of Soil Law and Policy, H. Ginzky, I. Heuse, T. Qin, O. Ruppel, P. Wegerdt, Eds. (Springer, Chambridge, 2017), 149–164 (2016).
- 14. M. Bendixen. I. Overeem, M. T. Rosing, A. A. Bjørk, K. H. Kjær, A., Kroon, G. Zeitz, L. Lønsmann Iversen, Promises and perils of sand exploitation in Greenland. Nature sustainability 2, 98-104 (2019).
- 15. M. L. C. M. Henckens, C. M. J. Ryngaert, P. P. J. Driessen, E. Worrell, Normative principles and the sustainable use of geologically scarce mineral resources. Resources Policy 59, 351-359 (2018)
- 16. A. Alaoma, N. Voulvoulis, Mineral resource active regions: The need for systems thinking in management. AIMS Envir. Sci. 5, 78-95 (2018)
- 17. A. R. Berger, Assessing rapid environmental change using geoindicators. Environ. Geol. 32, 36–44 (1997).

Funding: FS acknowledged support by the University of Nottingham Anne McLaren fellowship; JH received funding from the Academy of Finland project 315519; WDK acknowledges support from the University of Amsterdam Faculty Research Cluster 'Global Ecology'.

Author contributions: FS, JJB and RF conceived the idea with major contributions by WDK, KFR, ACS, DvR, JH and RSL as well as further input by all co-authors for conceptualization and methodology. FS led the writing with major contributions by JJB and RF, as well as further contributions by all other co-authors. FS designed and made Fig. 1, JCG designed and made Table S1, and both received contributions from all co-authors.

Competing interests: All authors declare no conflict of interests.



