

Est.
1841

YORK
ST JOHN
UNIVERSITY

Dechant, Pierre-Philippe ORCID

logoORCID: <https://orcid.org/0000-0002-4694-4010> (2016) A conformal geometric algebra construction of the modular group. In: Alterman Conference on Geometric Algebra, 1st - 9th August 2016, Brasov, Romania. (Unpublished)

Downloaded from: <https://ray.yorks.ac.uk/id/eprint/4005/>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at ray@yorks.ac.uk



A conformal geometric algebra construction of the modular group


Pierre-Philippe Dechant

Mathematics Department, University of York

Alterman Conference Brasov – August 6th, 2016


- 1 The modular group and the braid group
- 2 Motivation: Moonshine
- 3 Clifford algebra and the conformal construction
- 4 A CGA construction of the modular group

Torus





- A **nice manifold**, compact, 2 real-dimensional, or 1-complex dimensional
- The **unique Calabi-Yau 1-fold**
- The product of **two circles**: $T^2 = S^1 \times S^1$
- A torus is actually **topologically flat**, can cut open along the two circles to get a **parallelogram**

Torus embedding in the plane




- One can **tile** the plane with parallelograms in many ways (**lattice**)
- One can embed the **torus** if one picks such a lattice and **periodically identifies** opposite sides
- Considering the bottom left hand corner as the origin, and the top right hand corner as a number in the complex plane then this number is called the **complex structure of the torus τ**

Symmetries of the embedding



- **Many** possible embeddings from the same torus: **redundancy**
- **Winding** around the torus more often $\tau \rightarrow \tau + 1$
- **Flipping** the long and the short side of the torus $\tau \rightarrow -\frac{1}{\tau}$

The modular group




- The **modular generators** $T : \tau \rightarrow \tau + 1$, $S : \tau \rightarrow -\frac{1}{\tau}$ generate the modular group
- They satisfy the **abstract relations** $\langle S, T \mid S^2 = I, (ST)^3 = I \rangle$
($T^\infty = I$)
- Keyhole **fundamental region**

The modular group as $SL(2, \mathbb{Z})$

- $SL(2, \mathbb{Z})$: $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $ad - bc = 1$
- Subgroup of $SL(2, \mathbb{R})$, and the group of **Moebius** transformations of the plane, i.e. the 2D **conformal** group $PGL(2, \mathbb{C})$

The braid group




- B_n : the group of **braiding**s of n strands
- Presentation

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \ (1 \leq i \leq n-2), \sigma_i \sigma_j = \sigma_j \sigma_i \ (|i-j| \geq 2) \rangle$$
- B_3 is a **double cover** of the modular group

Modular functions

$\mathbb{H} := \{\tau \in \mathbb{C} \mid \text{Im}(\tau) > 0\}$, $q = \exp(2i\pi\tau)$; $SL(2, \mathbb{R})$ action on \mathbb{H} :




$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \tau = \frac{a\tau + b}{c\tau + d}, \quad ad - bc = 1$$

Take G a discrete subgroup of $SL(2, \mathbb{R})$ commensurable with $SL(2, \mathbb{Z})$ then f_G is a **modular function** for G iff

- $f_G : \mathbb{H} \rightarrow \mathbb{C}$ is **meromorphic**
- $f_G(\tau) = f_G\left(\frac{a\tau + b}{c\tau + d}\right) \quad \forall \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G$
- $f_G(A \cdot \tau) = \sum_{n=-\infty}^{\infty} b_n q^{n/N} \quad \forall A \in SL(2, \mathbb{Z})$ and some N, b_n depending on A with $b_n = 0$ for $n < -M$, $M \in \mathbb{N}$

Modular forms




$\Phi(q)$ a **modular form** of $SL(2, \mathbb{Z})$ of **weight** k if

$$\Phi\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^k \Phi(\tau) \quad \forall \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$$

- **Dedekind eta-function** $\eta(q) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n)$
- **Theta functions** from an even unimodular lattice L :
 $\theta_L(z) = \sum_{\lambda \in L} \exp \pi i \|\lambda\|^2 z$
- **Eisenstein series** from a lattice Λ : $E_k(\Lambda) = \sum_{0 \neq \lambda \in \Lambda} \lambda^{-k}$ a modular form of weight k

Weak Jacobi forms



weak Jacobi form of weight k and index m

$$\Phi\left(\frac{\mu}{c\mu+d}, \frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^k \exp\left(\frac{2i\pi mc\mu^2}{c\tau+d}\right) \Phi(\mu, \tau)$$

$$\forall \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}), \mu \in \mathbb{C}, \tau \in \mathbb{H}, z = \exp(2i\pi\mu), q = \exp(2i\pi\tau)$$

$$\Phi(\mu + a\tau + b, \tau) = \exp(-2i\pi m(a^2\tau + 2b\mu)) \Phi(\mu, \tau) \quad \forall a, b \in \mathbb{Z}$$

Modular forms and number theory



- **Number theorists** are very active in modular things
- Connections with **elliptic curves**, Taniyama-Shimura, Andrew Wiles' proof of **Fermat's theorem**
- Extremely clever people, formidably **hard** (a lot worse than what's on the slides)

- 1 The modular group and the braid group
- 2 Motivation: Moonshine**
- 3 Clifford algebra and the conformal construction
- 4 A CGA construction of the modular group

Moonshine



- Connection between two very different areas of Mathematics: **finite simple groups** and **modular forms**
- **Moonshine** = crazy, unlikely connection; **insubstantial**; illegal distilling of information from character table
- **Monstrous** = belonging to the Monster group, **enormous**, amazing


The Monster

- **Finite simple** groups: 18 **series** and 26 **sporadic** (exceptional)
- **Monster**: the **largest** sporadic group
- **Order**:

$$2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 11^2 \cdot 13^3 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71 \sim 8 \cdot 10^{53}$$
- **194 irreducible representations**: 1, 196883, 21296876, ...
- 20 sporadic groups are subquotients of the Monster: **The Happy Family**
- 6 are not: the **Pariahs**



The Klein j -invariant




- j -invariant $j(\tau) = \frac{\Theta_{E_8}(\tau)^3}{\eta(\tau)^{24}} - 744$
- **Hauptmodul** for the genus 0 group $G = SL(2, \mathbb{Z})$
- **Ogg**: genus 0 iff p is
2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 41, 47, 59, 71 offering a bottle
of **Jack Daniel's whiskey**
- Periodic: Fourier expansion **coefficients** wrt $q = e^{2\pi i\tau}$
- $$j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + \dots$$

Monstrous Moonshine


- **Mysterious connection** between two very different areas (**discrete** and **non-discrete**) of Mathematics noticed by **John McKay 1978**
- Monster $1, 196883, 21296876, \dots$
- Klein $j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + \dots$
- $196884 = 196883 + 1, 21493760 = 21296876 + 196883 + 1, \dots$
- Conway, Norton conjectured, Atkin, Fong, and Smith showed a moonshine module exists in 1980 (vertex operator algebra)
- Constructed by Richard **Borcherds** in 1992

The Mathieu group M_{24}




- One of 5 sporadic groups (first to be discovered) named after Mathieu: $M_{11}, M_{12}, M_{22}, M_{23}, M_{24}$
- Multiply transitive **permutation** groups
- $244823040 = 3 \cdot 16 \cdot 20 \cdot 21 \cdot 22 \cdot 23 \cdot 24 = 2^{10} \cdot 3^3 \cdot 5 \cdot 7 \cdot 11 \cdot 23$
- Arises as **automorphism group** of the **extended binary Golay code**, **Leech lattice** or **Steiner system** $S(5, 8, 24)$
- 26 irreps 1, 23, 45, 231, 252, 253, 483, 770, 990, 1035, 1265, 1771, 2024, 2277, 3312, 3520, 5313, 5544, 5796, 10395

K3 – Kähler, Kummer, Kodaira (mountain K2 in Kashmir, André Weil)



- 2 complex-dimensional manifold
- $T^2 \times T^2$ and K3 are the **only Calabi-Yau 2-folds**
- **Kähler** manifold, Ricci-flat, **compact**
- Often used in **string theory compactifications** along with T^2
- From orbifold resolutions, Kummer surfaces etc – all **diffeomorphic**

String Theory



- A supersymmetric **point particle** evolving along a closed **loop** in space-time
- A closed **superstring** sweeps out a 2D **torus**
- **Index of Dirac operator** and its **stringy** generalisation **Ramond** operator
- Compactifications on **tori** and **K3** manifolds
- Interesting (roughly) **modular quantities** and properties arise in string theory: e.g. $N = 4$ **characters**, **elliptic genus** = counts different **states**

K3 elliptic genus

- Known since 80s, **innocuous** enough looking

$$E_{K3}(z, q) = 8 \left(\frac{\theta_2^2(z, q)}{\theta_2^2(1, q)} + \frac{\theta_3^2(z, q)}{\theta_3^2(1, q)} + \frac{\theta_4^2(z, q)}{\theta_4^2(1, q)} \right)$$

- A weak Jacobi form of weight $k = 0$ and index $m = 1$
- Special values give **Euler characteristic**, \hat{A} **Dirac** index, σ **Hirzebruch** signature
- If **rewrite** this in terms of $N = 4$ characters of a specific **string theoretic** non-linear sigma model describing superstring propagation on a **K3** surface (!) one gets

$$E_{K3}(\tau, z) = -2Ch(0; \tau, z) + 20Ch(1/2; \tau, z) + e(q)Ch(\tau, z)$$

- Coefficients in the **q -series** are

$$e(q) = 90q + 462q^2 + 1540q^3 + 4554q^4 + 11592q^5 + \dots$$

Mathieu Moonshine – Eguchi, Ooguri, Tachikawa 2010

- **Similar Moonshine phenomenon** connecting finite simple groups and modular forms (ish..)
- **elliptic genus** of an $\mathcal{N} = 4$ SCFT compactified on a **K3-surface** (Taormina, Eguchi 80s)
- Finite simple group: **Mathieu M_{24}** 45,231,770,2277,5796...

- Elliptic genus is

$$E_{K3}(\tau, z) = -2Ch(0; \tau, z) + 20Ch(1/2; \tau, z) + e(q)Ch(\tau, z)$$

- All the coefficients in the q -series

$$e(q) = 90q + 462q^2 + 1540q^3 + 4554q^4 + 11592q^5 + \dots$$
 are

twice the dimension of some M_{24} irrep

Mathieu Moonshine – state of the field

- Generally **pretty lost** what's going on
- **Partial results**, e.g. proving coefficients are **positive even integers**, dimensions of M_{24} irreps etc
- **Nothing** in the string theory actually **can have full M_{24} symmetry!**?
- Elliptic genus is invariant under **surfing** across different **Kummer surfaces** despite counting different states in different theories etc e.g. given by symmetry group of D_4 lattice i.e. **binary tetrahedral group**
- Generalisation **umbral moonshine** connecting M_{24} to 23 other symmetry groups of **Niemeyer lattices**

Mathieu Moonshine – state of the field

- **Number theorists** chime in: weakly holomorphic mock modular form of weight $1/2$ on $SL(2, \mathbb{Z})$ with shadow $\eta(q)^3$.
All **extremely technical**
- My construction of **exceptional root systems** from Thursday ties in with these things: lattices, symmetry groups of Kummer surfaces, McKay correspondence, Trinities, Moonshine
- **New approach** to **modular symmetry**?

- 1 The modular group and the braid group
- 2 Motivation: Moonshine
- 3 Clifford algebra and the conformal construction**
- 4 A CGA construction of the modular group

Clifford Algebra and orthogonal transformations

- **Inner product** is symmetric part $a \cdot b = \frac{1}{2}(ab + ba)$
- Reflecting a in b is given by $a' = a - 2(a \cdot b)b = -bab$ (b and $-b$ **doubly cover** the same reflection)
- Via **Cartan-Dieudonné** theorem any orthogonal (/conformal/modular) transformation can be written as **successive reflections**

$$x' = \pm n_1 n_2 \dots n_k x n_k \dots n_2 n_1 = \pm A x \tilde{A}$$

- The conformal group $C(p, q) \sim SO(p+1, q+1)$ so can use these for **translations, inversions** etc as well

Conformal Geometric Algebra

- Go to $e_0, e_1, e_2, e_3, e, \bar{e}$, with $e_0^2 = 1, e_i^2 = -1, e^2 = 1, \bar{e}^2 = -1$
- Define two **null** vectors $n \equiv e + \bar{e}, \bar{n} \equiv e - \bar{e}$
- Can **embed** the 4D vector $x = x^\mu e_\mu = te_0 + xe_1 + ye_2 + ze_3$ as a **null vector in 6D** (also normalise $\hat{X} \cdot e = -1$)

$$\hat{X} = \frac{1}{\lambda^2 - x^2} (x^2 n + 2\lambda x - \lambda^2 \bar{n})$$

- So neat thing is that **conformal transformations** are now done by **rotors** (except inversion which is a reflection) – distances are given by **inner products**


Conformal Transformations in CGA

$$\hat{X} = \frac{1}{\lambda^2 - x^2} (x^2 n + 2\lambda x - \lambda^2 \bar{n})$$

- **Reflection:** spacetime $F(-axa) = -aF(x)a$
- **Rotation:** spacetime $F(Rx\tilde{R}) = RF(x)\tilde{R}$, $R = \exp(\frac{ab}{2\lambda})$
- **Translation:** $F(x+a) = R_T F(x)\tilde{R}_T$ for $R_T = \exp(\frac{na}{2\lambda}) = 1 + \frac{na}{2\lambda}$
- **Dilation:** $F(e^\alpha x) = R_D F(x)\tilde{R}_D$ for $R_D = \exp(\frac{\alpha}{2\lambda} e\bar{e})$
- **Inversion:** Reflection in extra dimension e : $F(\frac{x}{x^2}) = -eF(x)e$
 sends $n \leftrightarrow \bar{n}$
- **Special conformal transformation:** $F(\frac{x}{1+ax}) = R_S F(x)\tilde{R}_S$ for
 $R_S = R_I R_T R_I$

- 1 The modular group and the braid group
- 2 Motivation: Moonshine
- 3 Clifford algebra and the conformal construction
- 4 A CGA construction of the modular group**

Modular group



- Modular generators: $T : \tau \rightarrow \tau + 1$, $S : \tau \rightarrow -1/\tau$
- $\langle S, T \mid S^2 = I, (ST)^3 = I \rangle$ CGA: $R_Y X \tilde{R}_Y$
- CGA: $T_X = \exp\left(\frac{ne_1}{2}\right) = 1 + \frac{ne_1}{2}$ and $S_X = e_1 e$ (slight issue of complex structure $\tau =$ complex number, not vector in the 2D real plane so map $e_1 : x_1 e_1 + x_2 e_2 \leftrightarrow x_1 + x_2 e_1 e_2 = x_1 + ix_2$)
- $(S_X T_X)^3 = -1$ and $S_X^2 = -1$
- So a 3-fold and a 2-fold rotation in conformal space

Braid group

- $(S_X T_X)^3 = -1$ and $S_X^2 = -1$ is inherently **spinorial**
- Of course Clifford construction gives a **double cover**
- The **braid group** is a double cover
- So **Clifford** construction gives the **braid group double cover** of the **modular group**
- $\sigma_1 = \tilde{T}_X$ and $\sigma_2 = T_X S_X T_X$ satisfying $\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2$
($= S_X$)

Braid group

- $(S_X T_X)^3 = -1$ and $S_X^2 = -1$ is inherently **spinorial**
- Of course Clifford construction gives a **double cover**
- The **braid group** is a double cover
- So **Clifford** construction gives the **braid group double cover** of the **modular group**
- $\sigma_1 = \tilde{T}_X = \exp(-ne_1/2)$ and $\sigma_2 = T_X S_X T_X = \exp(-\bar{n}e_1/2)$ satisfying $\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 (= S_X)$
- Might not be known?

Where to go from here?

- Routinely do Clifford **complex analysis** in the plane
- Could look at **meromorphic functions**
- Look at functions in the complex plane in **CGA** representation
- Consider **modular symmetry** in this setup: **modular functions, modular forms, weak Jacobi forms** etc
- Perhaps the spinorial approach opens up **new techniques** to deal with the formidable algebra?