

Est.
1841

YORK
ST JOHN
UNIVERSITY

Bartel, S. J., Sherry, S. B., Smith, Martin
M. ORCID logoORCID: <https://orcid.org/0000-0002-4754-3032>,
Stewart, S. H., Glowacka, M. and Speth, T. A. (2020) Social
influences on binge drinking in emerging adults: Which social
network members matter most? Substance abuse.

Downloaded from: <https://ray.yorks.ac.uk/id/eprint/4228/>

The version presented here may differ from the published version or version of record. If
you intend to cite from the work you are advised to consult the publisher's version:
<https://www.tandfonline.com/doi/full/10.1080/08897077.2019.1709604>

Research at York St John (RaY) is an institutional repository. It supports the principles of
open access by making the research outputs of the University available in digital form.
Copyright of the items stored in RaY reside with the authors and/or other copyright
owners. Users may access full text items free of charge, and may download a copy for
private study or non-commercial research. For further reuse terms, see licence terms
governing individual outputs. [Institutional Repository Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at ray@yorks.ac.uk

Substance Abuse



**Social influences on binge drinking in emerging adults:
Which social network members matter most?**

Journal:	<i>Substance Abuse</i>
Manuscript ID	WSUB-2019-3444
Manuscript Type:	Brief Report
Keywords:	binge drinking, emerging adults, alcohol, social network

SCHOLARONE™
Manuscripts

Abstract

Background: Binge drinking peaks in emerging adulthood and is associated with a myriad of negative consequences. Research indicates social network members have a significant influence on binge drinking. In particular, theory suggests that drinking habits of romantic partners and peers have a stronger influence on emerging adult binge drinking than do drinking habits of siblings and parents. We investigated the relative influences of siblings, parents, romantic partners, and peers on emerging adults' binge drinking using a multi-source design and a robust measure of binge drinking. We hypothesized that peer and romantic partner binge drinking would more strongly predict emerging adult (target) binge drinking than would parent and sibling binge drinking. *Methods:* We recruited 321 participants (targets) aged 17-25 years, alongside 882 members of their social network (influencers). Targets and influencers completed self-report measures of binge drinking (frequency, quantity, and self-perception). *Results:* Structural equation modeling revealed the direct effect from romantic partner binge drinking to target binge drinking was significant. In contrast, the direct effects from peer, parent, and sibling binge drinking to target binge drinking were non-significant. *Conclusion:* In emerging adulthood, romantic partners appear to exert the strongest social influences on binge drinking.

Keywords: binge drinking, emerging adults, alcohol, social network

BINGE DRINKING INFLUENCERS IN EMERGING ADULTS

2

Social influences on binge drinking in emerging adults:

Which social network members matter most?

Binge drinking—the consumption of 5+ drinks in two-hours (4+ for women)¹ peaks in early adulthood² and is associated with health and social risks such as accidents, lost productivity, and relationship problems³. As these negative consequences are especially prevalent in emerging adulthood (17-25 years)⁴, identifying factors, such as social influences, that contribute to binge drinking in emerging adulthood is vital.

Emerging adults are embedded within social networks where members influence one another's alcohol use via active (e.g., explicit offers of alcohol) and passive (e.g., modeling, perceived norms) mechanisms⁵. Developmental theorists suggest that peers and romantic partners are the most important sources of social influence for emerging adults⁶⁻⁷. During emerging adulthood, young people shift from spending most of their time with parents and siblings to spending more time with peers and romantic partners⁴. As such, emerging adults rely more on peers and partners than family for support and intimacy⁴.

Indeed, among emerging adults, peer use of alcohol is a robust predictor of regular and hazardous drinking⁸, and both active and passive influences from peers independently predict binge drinking⁹. Similarly, the alcohol use of partners influences individual binge drinking over the short- and the long-term and predicts changes in alcohol consumption between adolescence and emerging adulthood^{10,11,12}.

In contrast, the impact of parent and sibling alcohol use on the alcohol use of emerging adults is unclear. Studies on sibling influence in emerging adulthood suggest the influence of siblings is modest, and tends to occur from older siblings, especially if siblings are close in age and the same sex¹³. Research examining the impact of parental use suggests that parental alcohol

1
2
3
4 use increases the likelihood that emerging adults will also drink alcohol¹⁴; however, relative to
5
6 the influence of peer drinking, parental drinking might have a smaller impact on emerging
7
8 adults' drinking¹⁴.
9

10 **Advancing Literature on Binge Drinking and Social Influence in Emerging Adults**

11
12
13 Research has focused primarily on specific social network members or undifferentiated
14
15 social networks rather than contributions of multiple types of network members on emerging
16
17 adult alcohol use. Our aim was to address this by determining the most important influencers.
18
19 Moreover, research has typically included participant perceptions of network members' alcohol
20
21 use instead of having network members directly report on their alcohol use⁸. As perceptions of
22
23 others' alcohol use can be biased, it is unclear if the actual alcohol use of network members is an
24
25 important influence¹⁵. Therefore, we collected self-reported alcohol use directly from
26
27 influencers. Our study is the first to compare the influence of multiple types of network members
28
29 using direct reports. Finally, we advanced the literature by measuring binge drinking as a latent
30
31 variable composed of frequency, severity, and perceptions of binge drinking. Building on theory
32
33 and research^{4,10}, we hypothesized that the binge drinking of peers and romantic partners, but not
34
35 the binge drinking of parents and siblings, will predict target binge drinking.
36
37
38
39

40 **Methods**

41 **Participants**

42
43 We recruited 321 undergraduate emerging adult targets (M age = 19.5; range 17-25;
44
45 72.9% female; 80.4% Caucasian) from XXX as part of a larger study (XXX). We also contacted
46
47 1,680 social network members (influencers) of our targets; 962 influencers responded (M = 3.28
48
49 influencers/target). Influencers were comprised of 166 mothers, 99 fathers, 65 sisters, 37
50
51 brothers, 3 spouses, 80 dating partners, 373 friends, 52 roommates, 4 classmates, 3 co-workers,
52
53
54
55
56
57
58
59
60

BINGE DRINKING INFLUENCERS IN EMERGING ADULTS

4

42 other relatives (e.g., grandparent), and 32 “others.” Influencers categorized as “other” and “other relatives” were excluded, as were influencers who did not specify their relationship to the target ($n = 6$). There was subsequently a total of 882 influencers: 265 parents, 102 siblings, 432 peers, and 83 romantic partners. Influencer characteristics are in Table 1.

---TABLE 1 HERE---

Measures

Binge drinking frequency. Consistent with the NIAAA¹, targets and influencers were asked, “during the past 7 days, how often did you have 5 (4 for women) or more drinks containing any kind of alcohol within a 2-hour period?” Targets and influencers responded on a 12-point scale from “0 times” to “10+ times”. This item correlates strongly with the NIAAA’s measure of binge drinking frequency¹ ($r = .68-.83$)¹¹.

Binge drinking severity. Targets and influencers were asked, “What is the greatest number of drinks you consumed in a 2-hour period during the past 7 days?”, as a measure of binge drinking severity¹¹.

Binge drinking self-perceptions. Targets and influencers were asked three items assessing perceptions of their binge drinking (“During the past 7 days, there were times when I rapidly drank a very large amount of alcohol within a 2 hour period;” “The average person would be amazed if s/he knew how much alcohol I consumed within a 2 hour period (during the past 7 days);” “During the past 7 days, there were times when I drank what other people would regard as an unusually large amount of alcohol within a 2 hour period”)¹¹. Items were rated on a 5-point scale from 1 (*strongly disagree*) to 5 (*strongly agree*). This scale has adequate reliability ($\alpha = .77-.87$)¹¹.

Procedure

1
2
3
4 Targets completed paper-and-pencil versions of measures and provided the email address
5
6 of five social network members (influencers) whom they had known for at least three months
7
8 and had contact with at least twice a week. Eligible influencers completed measures online.
9
10 Targets were compensated with one credit towards a psychology course. Influencers were
11
12 entered into 1-of-20 \$50 draws.
13
14

15 Results

17 Descriptive Statistics

19 Means, standard deviations, bivariate correlations, and Cronbach's alphas appear in Table
20
21 2. Following Cohen's guidelines¹⁶ for small, moderate, and large effects ($r = .10, .30, .50$), target
22
23 binge drinking had a moderate positive relationship with peer binge drinking and a large positive
24
25 relationship with partner binge drinking. In contrast, the relationships between target binge
26
27 drinking with parent and sibling binge drinking were small and non-significant ($p < .05$).
28
29
30

31 ---TABLE 2 HERE---

33 Structural Equation Modeling

35 We tested if parent, sibling, peer, and/or partner binge drinking were predictors of target
36
37 binge drinking using structural equation modeling with FIML estimation in AMOS¹⁷. We
38
39 evaluated fit using CFI, TLI, and RMSEA. CFI and TLI $\geq .95$, and RMSEA $\leq .05$ indicate good
40
41 fit¹⁸.
42
43
44

45 **Measurement model.** The measurement model (Figure 1) fit well: $\chi^2 (80) = 128.12$, CFI
46
47 = .968; TLI = .952; RMSEA = .043, 90% CI [.029-.057]. Factor loadings were significant ($p <$
48
49 $.001$) and $> .40$.
50
51

52 ---FIGURE 1 HERE---

54 *Figure 1.* Measurement model. Ovals represent latent variables. Rectangles represent observed
55
56 indicators. Double-headed black arrows represent significant correlations ($p < .05$). Double
57
58
59
60

BINGE DRINKING INFLUENCERS IN EMERGING ADULTS

6

headed grey arrows represent non-significant correlations ($p > .05$). Single headed black arrows represent significant loadings ($p < .05$). Pairwise deletion was utilized. BD = binge drinking.

Structural model. The structural model (Figure 2) had the identical fit indices as listed above for the measurement model. Only romantic partner binge drinking predicted target binge drinking ($\beta = .69, p < .001$).

----FIGURE 2 HERE---

Figure 2. Structural Model. Significant direct effects ($p < .05$) are represented by single-headed black arrows. Non-significant direct effects ($p > .05$) are represented by single-headed grey arrows. Double headed arrows signify covariances. Estimates are standardized.

Discussion

We tested the effects of the binge drinking patterns of social network members on the binge drinking patterns of emerging adults using a multi-source design and a psychometrically-robust binge drinking measure. We investigated the relative influence of different social network member groups by examining various social network members in a single model.

As hypothesized, parent and sibling binge drinking did not predict target binge drinking. Our results align with theory and research suggesting that the relative influence of parents and siblings decreases with age^{4,6,14}. Similarly, consistent with Bartel and colleagues¹⁰ and Mushquash and colleagues¹¹ and as hypothesized, romantic partners' binge drinking predicted target binge drinking. The large size of this effect probably reflects the important influence of romantic partners during emerging adulthood and the cross-sectional nature of our study, which likely captured selection and socialization effects.

People select romantic partners based on the similarity of characteristics, values, and behaviors, which may lead people to choose partners with similar levels of alcohol use¹². This similarity-based selection of partners may be especially important given findings that the degree

BINGE DRINKING INFLUENCERS IN EMERGING ADULTS

7

1
2
3
4 of similarity in alcohol use between partners impacts relationship quality¹². As such, the
5
6 similarity of binge drinking between partners in our sample may partially reflect this selection
7
8 process. Additionally, once in a relationship, partners may further reduce discrepancies between
9
10 their alcohol use patterns and that of their partners to improve relationship quality and meet
11
12 needs for approval and acceptance¹⁹. This socialization effect may be particularly strong in
13
14 romantic partner contexts, given social learning theory suggests that the behavior of those who
15
16 are highly valued is most likely to be emulated⁵, and developmental theory suggests that
17
18 romantic partners are important in emerging adulthood⁷. Thus, partners might influence *how*
19
20 *much* individuals consume alcohol²⁰.

21
22
23
24 Partners might also influence *why* individuals drink alcohol; individuals may observe
25
26 their partners' motivations for drinking (e.g., drinking to cope with anxiety) and adopt those
27
28 reasons for drinking themselves, which can result in an escalation of drinking behavior²⁰.
29
30
31 Ultimately, the combined effects of selection and socialization might result in a "drinking
32
33 partnership"²¹ which encourages binge drinking and may explain our strong direct effects of
34
35 partners on target binge drinking.
36
37

38
39 Unlike partner binge drinking and contrary to hypotheses, peer binge drinking did not
40
41 predict target binge drinking. Broad peer networks have an impact on binge drinking while in
42
43 general single peers, except for "drinking buddies," might not²²⁻²³. Thus, we might not have
44
45 included enough peers to capture the influence of a peer network or might not have included the
46
47 "right" single peers. Secondly, peer influence might be the result of descriptive or injunctive
48
49 norms (how much peers are perceived to drink; what is perceived to be accepted by the peer
50
51 group) rather than the actual drinking habits of specific individuals. Third, as romantic partners
52
53 are likely present in many peer-based interactions, some of the variance from peers may have
54
55
56
57
58
59
60

1
2
3
4 been captured and accounted for by partners in the model. Nonetheless, our results accord with
5
6 developmental theory suggesting that during the transition from adolescence to emerging
7
8 adulthood, the importance of peers declines, while the importance of romantic partners
9
10 increases²⁴.

11 12 **Limitations and Future Directions**

13
14
15 Our results are limited by our predominately female, Caucasian, and university-educated
16
17 sample and may not be generalizable to more diverse groups of emerging adults. Future research
18
19 should adopt a longitudinal design. Additionally, a broader peer network was not captured by our
20
21 study, nor was the influence of different types of peers (e.g., drinking buddies vs. casual
22
23 acquaintances). The impact of these distinct sources of peer influence should be investigated and
24
25 compared to the influence of romantic partners. Finally, though our results suggest parental
26
27 binge drinking is not a significant influence, parents might influence emerging adults via
28
29 mechanisms other than personal alcohol use (e.g., rules about drinking)²⁵.

30 31 32 **Concluding Remarks**

33
34
35
36 When investigating the social networks of emerging adults, partners are a particularly
37
38 influential type of network member when it comes to binge drinking. The strength of romantic
39
40 partner influence has important implications for health-promotion campaigns, which could focus
41
42 on increasing awareness of partner influence. Additionally, the strength of romantic partner
43
44 influence also has implications for alcohol misuse interventions, which should not underestimate
45
46 the impact of partners on emerging adult binge drinking.
47
48

49
50 **Disclosure Statement:** The authors have no potential conflicts of interest to declare.
51
52
53
54
55
56
57
58
59
60

References

-
- ¹ National Institute on Alcohol Abuse and Alcoholism. NIAAA council approves definition of binge drinking. *NIAAA Newsletter*, 2004;3(3):1-4.
- ² Patrick ME, Terry-McElrath YM, Lanza ST, Jager J, Schulenberg JE, O'Malley PM. Shifting age of peak binge drinking prevalence: historical changes in normative trajectories among young adults aged 18 to 30. *Alcohol Clin Exp Res*. 2019;43(2):287-298.
- ³ Thavorncharoensap M, Teerawattananon Y, Yothasamut J, Lertpitakpong C. The economic impact of alcohol consumption: a systematic review. *Subst Abuse Treat Prev Policy*. 2009;4(1):20-31.
- ⁴ Arnett JJ. The developmental context of substance use in emerging adulthood. *J Drug Issues*. 2005;35(2):235-254.
- ⁵ Bandura A., Walters RH. *Social learning theory* (Vol. 1). Englewood Cliffs, NJ: Prentice-Hall; 1977.
- ⁶ Borsari B, Carey KB. Peer influences on college drinking: a review of the research. *J Subst Abuse*. 2001;13(4):391-424.
- ⁷ Larson R, Richards MH. Daily companionship in late childhood and early adolescence: changing developmental contexts. *Child Dev*. 1991;62(2):284-300.
- ⁸ Mason MJ, Zaharakis N, Benotsch EG. Social networks, substance use, and mental health in college students. *J Am Coll Health*. 2014;62(7):470-477.
- ⁹ Read JP, Wood MD, Capone C. A prospective investigation of relations between social influences and alcohol involvement during the transition into college. *J Stud Alcohol*. 2005;66(1):23-34.
- ¹⁰ Bartel SJ, Sherry SB, Molnar DS, et al. Do romantic partners influence each other's heavy episodic drinking? Support for the partner influence hypothesis in a three-year longitudinal study. *Addict Behav*. 2017;69:55-58
- ¹¹ Mushquash AR, Stewart SH, Sherry SB, Sherry DL, Mushquash CJ, MacKinnon AL. Depressive symptoms are a vulnerability factor for heavy episodic drinking: a short-term, four-wave longitudinal study of undergraduate women. *Addict Behav*. 2013;38(5):2180-2186.
- ¹² Wiersma JD, Fischer JL & Fitzpatrick J. The role of romantic partners' drinking and bingeing patterns in relationship quality and alcohol-related problems. In DiGuarde KI, *Binge Drinking Research Progress*. Hauppauge, NY: Nova Science; 2009:39-61.

BINGE DRINKING INFLUENCERS IN EMERGING ADULTS

1

1
2
3
4
5 ¹³ Boyle MH, Sanford M, Szatmari P, Merikangas K, Offord DR. Familial influences on
6 substance use by adolescents and young adults. *Can J Public Health*. 2001;92(3):206-209.
7

8 ¹⁴ Scholte RH, Poelen EA, Willemsen G, Boomsma DI, Engels RC. Relative risks of adolescent
9 and young adult alcohol use: the role of drinking fathers, mothers, siblings, and friends. *Addict*
10 *Behav*. 2008;33(1):1-4.
11

12
13 ¹⁵ Henry DB, Kobus K, Schoeny ME. Accuracy and bias in adolescents' perceptions of
14 friends' substance use. *Psychol Addict Behav*. 2011;25(1):80-89.
15

16 ¹⁶ Cohen J. A power primer. *Psychol Bull*. 1992;112(1):155-159.
17

18 ¹⁷ Arbuckle JL. *Amos 20 user's guide*. Chicago, IL: SPSS Inc; 2011.
19

20
21 ¹⁸ Kline RB. Convergence of structural equation modeling and multilevel modeling. In
22 Williams M. *Handbook of methodological innovation*. Thousand Oaks, CA: Sage; 2011: 562-
23 589.
24

25
26 ¹⁹ Baumeister RF, Leary MR. The need to belong: desire for interpersonal attachments as a
27 fundamental human motivation. *Psychol Bull*. 1995;117(3):497-529.
28

29 ²⁰ Kehayes IL, Mackinnon SP, Sherry SB, Leonard KE, Stewart SH. Drinking motives and
30 drinking behaviors in romantic couples: A longitudinal actor-partner interdependence model.
31 *Psychol Addict Behav*. 2019; 33: 208-220.
32

33
34 ²¹ Roberts LJ, Leonard KE. An empirical typology of drinking partnerships and their relationship
35 to marital functioning and drinking consequences. *J Marriage Fam*. 1998;60(2):515-526.
36

37 ²² Reifman A, Watson WK, McCourt A. Social networks and college drinking: probing processes
38 of social influence and selection. *Pers Soc Psychol Bull*. 2006;32(6):820-832.
39

40
41 ²³ Nogueira-Arjona R, Shannon T, Kehayes IL, Sherry SB, Keough MT, Stewart SH. Drinking to
42 keep pace: a study of the moderating influence of extraversion on alcohol consumption similarity
43 in drinking buddy dyads. *Addict Behav*. 2019;92:69-75.
44

45 ²⁴ Erikson EH. *Childhood and society (2nd ed.)*. New York: Norton; 1963.
46

47
48 ²⁵ Wetherill RR, Fromme K. Alcohol use, sexual activity, and perceived risk in high school
49 athletes and non-athletes. *J Adolesc Health*. 2007;41(3):294-301.
50

Table 1

Influencer characteristics

Type	Number included	Mean age in years	Relationship length in years
Parents	265	51.16 (37-68)	20.91 (.08-26)
Siblings	102	21.14 (14-34)	19.71 (3.25-40.0)
Romantic	83	21.40 (17-36)	2.59 (.17-13.16)
Peers	432	20.20 (11-63)	5.01 (.08-35.9)

Note: As parents were not required to be biological parents, the value of .08 may represent a step-parent.

Table 2

Bivariate correlations, means, standard deviations, possible range, actual range, and alpha reliability

Variable	1	2	3	4	5
1. Target BD total	—				
2. Parent BD total	.12	—			
3. Sibling BD Total	.15	.12	—		
4. Peer BD total	.27**	-.03	-.03	—	
5. Partner BD total	.60**	.05	.54**	.17	—
<i>M</i>	7.86	5.67	6.46	8.75	9.37
<i>SD</i>	6.17	3.61	5.14	5.94	6.82
Possible range of scores	3+	3+	3+	3+	3+
Actual range of scores	3-35	3-26	3-23	3-34	3-19
Cronbach's alpha (α)	.91	.77	.89	.88	.81

Note: Missing data handled with pairwise deletion. **BD** = binge drinking. ($N = 20$ [partner/sibling] to 221 [target/peer]). Cronbach's alpha calculated using summation of standardized frequency, standardized severity, and standardized total perception scale. See measures section for previous research on the alpha of the three-item perception scale.

* $p < .05$ ** $p < .01$.




Figure 1




Figure 2