# Tan, Xiao ORCID logoORCID: https://orcid.org/0009-0008-0960-4974, Pan, Gang ORCID logoORCID: https://orcid.org/0000-0003-0920-3018, Che, Wenxue, Man, Ying, Wang, Liyi, Li, Jinting and Zhao, Hui (2025) Adaptation responses of Dunaliella sp. and its natural symbiotic bacteria to sulfamethoxazole. Environmental Technology & Innovation, 38. p. 104171.

Downloaded from: https://ray.yorksj.ac.uk/id/eprint/11899/

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version: https://doi.org/10.1016/j.eti.2025.104171

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. Institutional Repository Policy Statement

# RaY

Research at the University of York St John For more information please contact RaY at <u>ray@yorksj.ac.uk</u> Contents lists available at ScienceDirect



# Environmental Technology & Innovation



journal homepage: www.elsevier.com/locate/eti

# Adaptation responses of *Dunaliella* sp. and its natural symbiotic bacteria to sulfamethoxazole

Xiao Tan<sup>a</sup>, Gang Pan<sup>a,c,e</sup>, Wenxue Che<sup>a</sup>, Ying Man<sup>a</sup>, Liyi Wang<sup>a</sup>, Jinting Li<sup>a</sup>, Hui Zhao<sup>a,b,c,d,\*</sup>

<sup>a</sup> College of Chemistry and Environmental Science, Guangdong Ocean University, Zhanjiang 524088, China

<sup>b</sup> Southern Marine Science and Engineering Guangdong Laboratory, Zhuhai 519082, China

<sup>c</sup> Research Center for Coastal Environmental Protection and Ecological Resilience, Guangdong Ocean University, Zhanjiang 524088, China

<sup>d</sup> Cooperative Research Center for Offshore Marine Environmental Change, Guangdong Ocean University, Zhanjiang 524088, China

<sup>e</sup> School of Humanities, York St John University, York YO31 7EX, UK

ARTICLE INFO

Keywords: Sulfamethoxazole Microalgae Natural symbiotic bacteria Quorum sensing Dunaliella

#### ABSTRACT

Microalgal-bacterial consortia provide a sustainable, low-cost solution for antibioticcontaminated wastewater remediation. Microalgae cultivation typically involves natural symbiotic bacteria, which are often overlooked. To fill this gap, this study evaluated the adaptive response of *Dunaliella* sp. and its natural symbiotic bacteria to sulfamethoxazole (SMX) through cultivation experiments and 16S rDNA sequencing. The results demonstrated strong SMX tolerance but limited removal efficiency by the consortium. The adaptive responses were characterized by increased bacterial diversity and connectivity, with quorum sensing bacteria-driven community changes and biofilm formation playing a critical role in this process. *Dunaliella* sp. exhibited strong antioxidant capacity (with catalase playing a key role) and increased extracellular polymeric substances secretion, which not only contributed to its tolerance but also promoted microalgae-bacteria interactions, enhancing consortium adaptability. These findings strengthen our understanding of the responses of natural symbiotic bacteria to antibiotics and advance the development of microalgae-bacteria symbiosis technology for wastewater treatment.

## 1. Introduction

The demand for mariculture continues to expand with the decline in wild marine fishery resources. Globally, antibiotic usage reached 93 million tons in 2017 and is projected to exceed 236 million tons by 2030, with 5.7 % used in aquaculture (Adenaya et al., 2023). However, antibiotic abuse leads to elevated residue levels, disrupting microbial communities, promoting antibiotic resistance genes (ARGs), and causing toxicity to aquatic organisms (Wang et al., 2022a, Peng et al., 2025). Sulfamethoxazole (SMX), one of the most widely used antibiotics in aquaculture, is typically detected in water bodies at concentrations ranging from 67.9 ng/L to 63.6 µg/L (Zhu et al., 2022). In shrimp farming ponds, SMX concentrations can reach up to 2.39 mg/L (Le and Munekage, 2004), and in pharmaceutical wastewater can also reach mg/L levels (Zhang et al., 2025). Even in natural seawater, the highest reported SMX concentration is 1454.2 ng/L (Niu et al., 2016). Traditional methods like adsorption, activated sludge, advanced oxidation, and electrochemistry face challenges including inconsistent efficiency, high costs, and the potential to promote the spread of ARGs (Lu

\* Corresponding author at: College of Chemistry and Environmental Science, Guangdong Ocean University, Zhanjiang 524088, China. *E-mail address:* huizhao1978@163.com (H. Zhao).

https://doi.org/10.1016/j.eti.2025.104171

Received 4 February 2025; Received in revised form 13 March 2025; Accepted 22 March 2025

Available online 26 March 2025

<sup>2352-1864/© 2025</sup> The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

et al., 2025). In contrast, microalgae-based biological treatment technologies are gaining attention for their low cost, sustainability, and environmental friendliness.

Microalgae are important components of aquatic ecosystems and have been shown to degrade antibiotics while removing nitrogen and phosphorus from wastewater (Chandel et al., 2022, Zhang et al., 2022a). However, most microalgae show obvious growth inhibition under the influence of moderate concentrations of SMX (Xiong et al., 2019b, Xiong et al., 2020, Zhang et al., 2022b), limiting their practical application. Recent studies suggest that introducing foreign microorganisms can enhance the adaptability and removal efficiency of microalgae to antibiotics. This improvement is attributed to symbiotic interactions, involving extracellular polymeric substances (EPS)-mediated adhesion, nutrient exchange, oxygen release, and alterations in bacterial community structure and function (da Silva Rodrigues et al., 2020, Hu et al., 2022, Wang et al., 2022b). However, consortia composed of microalgae and foreign bacterial communities exhibit significant uncertainty and instability.

In practice, microalgae are often cultivated as monocultures rather than pure cultures, naturally carrying symbiotic bacteria whose roles have been largely overlooked. Most research has focused on the combined effects of algae with foreign microorganisms, while the role of natural symbiotic bacteria has not been fully studied. To fill this knowledge gap, this study focused on a microalga from the genus *Dunaliella*, a unique group of halophilic, unicellular, eukaryotic microalgae widely distributed in marine and hypersaline environments (Wang et al., 2023). *Dunaliella* has shown high tolerance to adverse environments, including high salinity (Liska et al., 2004), heavy metals (Folgar et al., 2009), microplastics (Chae et al., 2019), and various organic pollutants (Quesnel et al., 2011, Xiong et al., 2016). However, few studies have explored its response to antibiotics (Kvíderová and Henley, 2005, Machado and Soares, 2019, Hom-Diaz et al., 2022), and no studies have investigated the impact of SMX on *Dunaliella* or its natural symbiotic bacteria.

This study investigated the adaptive responses of *Dunaliella* sp. and its natural symbiotic bacteria to different SMX concentrations by analyzing microalgal growth, nutrient utilization, antibiotic removal, stress response, and 16S rDNA sequencing. The objectives were to investigate *Dunaliella* sp.'s SMX removal efficiency and tolerance mechanisms, and explore the response and interactions of the natural symbiotic bacterial community under SMX stress. To our knowledge, this is the first study to investigate the impact of antibiotic stress on microalgae's natural symbiotic bacteria. These findings provide new insights into the application of algae-bacteria symbiosis technology in bioremediation and bioenergy production.

# 2. Materials and methods

#### 2.1. Chemicals and microalgae

Tested SMX (purity  $\geq$ 98 %, CAS No. 723–46–6) was purchased from Aladdin Biochemical Technology Co., LTD (Shanghai, China). Methanol and acetonitrile (HPLC grade) were purchased from Thermo Fisher Scientific Inc. (USA). Other reagents were analytical grade unless otherwise specified. The marine microalga *Dunaliella* sp. (SCSIO-45184) was obtained from the Guangdong Province Economic Microalgae Species and Quality Resource Database (China) and cultured in F/2 medium.

#### 2.2. Microalgal-bacterial consortium culture and exposure experiment

The microalgal-bacterial consortium was cultured in simulated mariculture wastewater with SMX concentrations of 0, 1, and 10 mg/L for ten days. Although these concentrations exceed environmental levels, they are commonly used in algae-bacteria remediation screening studies to identify robust combinations and amplify their adaptive traits (Hu et al., 2022, Zhang et al., 2022a). Sterilised natural seawater was used as the basic culture medium supplemented with 4 mg/L nitrate (NO<sub>3</sub>-N), 2 mg/L ammonium (NH<sub>4</sub><sup>+</sup>-N), and 0.5 mg/L phosphates ( $PO_4^3$ -P) according to prior investigations. Glucose and sodium bicarbonate were added as carbon sources.

Experiments were conducted in triplicate using 1 L Erlenmeyer flasks, with an initial cell density of approximately  $1.0 \times 10^4$  cells mL<sup>-1</sup>. Cultures were maintained at  $25 \pm 1$ °C under a 12 h light/12 h dark cycle and shaken three to four times daily. From day 0, culture suspensions were collected every two days to determine the biomass, pigments, nutrients, and SMX content. Additionally, EPS, activities of the five key antioxidant enzymes, and contents of glutathione (GSH) and malondialdehyde (MDA) were only measured on days 4 and 10.

#### 2.3. Natural symbiotic bacteria exposure experiment

Natural symbiotic bacteria were cultured in 250 mL Erlenmeyer flasks (three replicates) under the same SMX and medium conditions as the consortium. Before incubation, the consortium was centrifuged at 4500 rpm for 15 min. The supernatant was discarded, and 0.5 % sterile saline was added to achieve thorough resuspension. *Dunaliella* sp. ruptured completely due to the absence of cell walls, leaving a natural symbiotic bacterial suspension for incubation (Fig. S1). Culture suspensions were collected every two days for nutrient analysis.

#### 2.4. Physiological analyses

Microalgae were fixed with Lugol's solution and counted using a haemocytometer under a microscope. Total chlorophyll and carotenoid contents were determined at 665, 652, and 470 nm using a spectrophotometer (METASH UV-9000S, China), calculated as follows (Xiong et al., 2017):

X. Tan et al.

Chlorophyll a (Chla) =  $16.82 \ A_{665} - 9.28 \ A_{652}$ 

Chlorophyll b (Chlb) =  $36.92 A_{652} - 16.54 A_{665}$ 

# $Carotenoid = (1000 \ A_{470} - 1.91 Chla - 95.15 Chlb)/225$

EPS were collected by centrifugation (4500 rpm, 5 minutes, 4°C) and classified as slime EPS (Mao et al., 2022). Polysaccharides (PS) and proteins (PN) were determined using assay kits based on bicinchoninic acid and anthrone-sulfuric acid methods, respectively. GSH and MDA contents and the activities of superoxide dismutase (SOD), glutathione peroxidase (GPX), glutathione S-transferase (GST), glutathione reductase (GR), and catalase (CAT) were detected using assay kits. The results were obtained using an ultra-micro spectrophotometer (BIO-DL Micro Drop, China). All kits, except for PS (Youxuan Biotechnology Co., Ltd., China), were sourced from the Nanjing Jiancheng Bioengineering Institute, China.

# 2.5. Analysis of nutrients and SMX

The concentrations of NO<sub>3</sub>-N, nitrite (NO<sub>2</sub>-N), NH<sup>+</sup><sub>4</sub>-N, PO<sup>3</sup><sub>4</sub>-P, total nitrogen (TN), and total phosphorus (TP) were determined using a flow analyser (Skalar San++, Netherlands) according to the standard methods for seawater detection by the Ministry of Ecology and Environment of China. Method details are presented in Supplementary Material Table S1. Solid-phase extraction and high-performance liquid chromatography with fluorescence detection were used to analyze SMX content, as detailed in the Supplementary Materials Text S1.

#### 2.6. 16S rDNA high-throughput sequencing of natural symbiotic bacteria

On days 0, 4, and 10, microalgal-bacterial culture samples were collected. A 100 mL aliquot was centrifuged at 4500 rpm for 15 min, and the supernatant was separated. 0.5 % sterile saline was added to achieve thorough resuspension, facilitating lysis of the microalgae cells. The suspension and supernatant were filtered through a 0.22 µm acetate membrane and sent to Majorbio Bio-Pharm Technology Co. Ltd. (Shanghai, China) for DNA extraction, PCR amplification and high-throughput sequencing. After quality filtering, merging, clustering, and chimera removal, 1256,804 optimized sequences were obtained for analysis.

D4\_0, D4\_1, and D4\_10 were blank, 1 mg/L, and 10 mg/L groups on day 4; D10\_0, D10\_1, and D10\_10 were the same on day 10. Species classification and functional gene annotation were based on SILVA138/16S\_bacteria and Kyoto Encyclopedia of Genes and Genomes databases. Raw sequencing data can be accessed from the National Centre for Biotechnology Information under accession number PRJNA1155904.

# 2.7. Statistical analysis

The standard curve correlation coefficient for each detection index exceeded 0.995, and 10 % random replicates were set up with a relative deviation of less than 2 %. Chemical indicators were analyzed using IBM SPSS Statistics 27 and microbial analysis was performed on the Majorbio platform. Treatment groups differences were compared using one-way ANOVA (p < 0.05). Results were visualized using Origin 2021 or Adobe Illustrator 2023.

#### 3. Results

# 3.1. Effect of SMX on the growth and pigment synthesis of Dunaliella sp

The effects of SMX concentration on Dunaliella sp. growth are presented in Fig. 1. Dunaliella sp. initially grew rapidly, with growth



**Fig. 1.** (a) Trends in the contents of Chla, Chlb, and carotenoid; (b) Trends in the cell density of *Dunaliella* sp.; and (c) Trends in the N/P ratio within *Dunaliella* sp. cells under different SMX concentrations during 10 d of cultivation. Error bars represent standard deviation (n = 3). Different letters indicate significant differences (p < 0.05) between treatments on the same day and the same letters or "ns" indicate no differences.

slowing after day 4 and no inhibition observed. From day 4 onward, the 10 mg/L group showed significantly higher biomass than other groups, reaching 1.53 times that of the blank group by the end of the experiment. Pigment content followed a similar trend, with significantly higher Chla and carotenoids contents observed in the 10 mg/L group from day 6.



**Fig. 2.** Trends in nutrient utilization by microalgal-bacterial consortium (a, c, e, g) and natural symbiotic bacteria alone (b, d, f, h) under different SMX concentrations during 10 d of cultivation. Error bars represent standard deviation (n = 3). Different letters indicate significant differences (p < 0.05) between treatments on the same day and the same letters indicate no differences.

The N/P ratio in *Dunaliella* sp. (Fig. 1c) changed as follows: under nutrient sufficient conditions on day 2, the N/P ratio was close to the Redfield ratio (N/P = 16). By day 4, under phosphorus limitation, the N/P ratio rose above 30, higher in SMX-treated groups than the blank group. Subsequently, the N/P ratio gradually decreased and stabilised at approximately 20.

# 3.2. Nutrient utilisation of Dunaliella sp. and its natural symbiotic bacteria under SMX stress

The effect of SMX concentration on nutrient utilization by the microalgal-bacterial consortium is presented in Fig. 2(a, c, e, g). NH<sub>4</sub><sup>+</sup>-N was consumed first, nearly depleted by day 2 except for the 10 mg/L group. NO<sub>3</sub>-N contents were stable initially, with slight increases in SMX-treated groups, but rapidly declined from day 4 and were eventually completely removed. Additionally, from day 4 onward, SMX-treated groups significantly accelerated NO<sub>3</sub><sup>-</sup>-N consumption, with NO<sub>3</sub>-N in the 10 mg/L group fully depleted by day 8. In contrast, NO<sub>2</sub>-N accumulation increased with higher SMX concentrations, but was consumed once NO<sub>3</sub>-N and NH<sub>4</sub><sup>+</sup>-N were completely absorbed. Unlike nitrogen, PO<sub>4</sub><sup>3</sup>-P was not completely utilized, and its concentration exhibited a fluctuating reduction following an initial rapid decline.

Changes in the nutrient utilization patterns of natural symbiotic bacteria under SMX stress after separation from *Dunaliella* sp. are presented in Fig. 2(b, d, f, h). The bacteria presented high NH<sup>+</sup><sub>4</sub>-N utilization efficiency, with the blank group consuming 83.4 % by day 2 and reaching 91.7 % eventually, but efficiency suppressed significantly with increasing SMX concentrations. NO<sub>3</sub>-N in the blank group was initially consumed but rapidly released, while levels in other groups remained stable. NO<sub>2</sub>-N also showed accumulation, but its contents were negligible compared to those in the microalgal-bacterial consortium. PO<sup>3</sup><sub>4</sub>-P utilization followed a pattern similar to ammonia but with slightly lower efficiency, peaking at 52.9 % in the blank group.

## 3.3. SMX removal efficiency and its effect on MDA and EPS content

The SMX residue rates at different SMX concentrations are presented in Fig. 3a, with values of 84.71 % and 90.20 % for the 1 and 10 mg/L groups, respectively. On day 4, MDA content (Fig. 3b) in the 10 mg/L group was 7.68 times higher than in the blank group but returned to normal by day 10. Additionally, on day 4, the 10 mg/L group showed significantly higher PS and PN contents in the slime



**Fig. 3.** Trends in (a) SMX residue rate, (b) MDA content in *Dunaliella* sp. cells, (c) PS content in slime EPS and (d) PN content in slime EPS under different SMX concentrations during incubation periods. Error bars represent standard deviation (n = 3). Different letters indicate significant differences (p < 0.05) between treatments on the same day and the same letters indicate no differences.

EPS (Fig. 3c-d) than those in blank group. By day 10, PS and PN contents in the 10 mg/L group had decreased.

# 3.4. Response of antioxidant system to SMX

Activity of antioxidant system in *Dunaliella* sp. are shown in Fig. 4a. On day 4, CAT, SOD, and GR activities increased with SMX concentration. The CAT activity in the 10 mg/L group was significantly higher than that in the blank group, while GST and GPX activities were significantly lower. By day 10, the CAT activity remained elevated in the 10 mg/L group, GST continued to decrease and GR activity reversed. SOD and GPX activities, and GSH content showed almost no differences among the treatment groups.

PCA results (Fig. 4b-c) showed that the sample points on day 4 were more clustered within groups and exhibited clearer separation between groups compared to day 10. On day 4, PC1 and PC2 explained 54.1 % and 18.5 % of the total variance, respectively. MDA, CAT, GR, PS, and PN were positively correlated with SMX concentrations, while GST and GPX showed negative correlations. On day 10, PC1 and PC2 explained 38.9 % and 24.6 % of total variance, respectively. Notably, the effect of MDA was negligible. CAT and PN remained positively correlated with SMX concentration, while the response of other indicators to SMX concentration weakened.

# 3.5. 16 s rDNA analysis on natural symbiotic bacteria of Dunaliella sp

#### 3.5.1. Natural symbiotic bacteria community structure analysis

The alpha diversity index was used to assess the changes in diversity of natural symbiotic bacterial communities under different SMX concentrations (Fig. 5a). Significantly higher Shannon's index and lower Simpson's index values with increasing SMX concentrations, implied increased biodiversity. Beta diversity (Fig. 5b), analysed by principal coordinate analysis (PCoA), revealed that the



Fig. 4. Under different SMX concentrations, (a) trends in the activities of antioxidant enzymes in *Dunaliella* sp. cells, and PCA of various physiological indicators related to the antioxidant system and EPS on day 4 (b) and day 10 (c).

bacterial composition in stimulated mariculture wastewater differed from the initial F/2 medium, and SMX significantly affected it as early as day 4.

The natural symbiotic bacterial community of *Dunaliella* sp. consisted of four phyla, 38 genera, and 40 species (15 identified). Only one genus disappeared during the incubation, with 31 genera remaining in all treatment groups (Fig. 5c). Bacterial composition results (Fig. 5d) showed that the relative abundance of dominant genera increased in stimulated mariculture wastewater compared to the F/2 medium but were evidently inhibited as SMX concentration increased. *Methylophaga* and *Labrenzia* had considerably lower relative abundances in the 10 mg/L group than in the blank group, while *Ponticoccus* and *Mameliella* increased. LDA Effect Size (LEfSe) analysis (LDA threshold = 4, Fig. 5e) identified 12 significantly different genera, mostly enriched in *Alphaproteobacteria* and derived from moderate abundance genera in day 10 samples from the 10 mg/L group.

One-way correlation networks were used to analyse the correlations between species and the results are presented in Fig. 6a. *Proteobacteria* remained the core phyla, while the core genera shifted from *Methylophaga* and *Labrenzia* to *Ponticoccus* and *Mameliella* as SMX concentration increased. The number of linkages markedly increased with both the incubation period and SMX concentration. For example, on day 4, the 10 mg/L group had 286 linkages, significantly higher than the blank (210) and 1 mg/L (198) groups. By day 10, linkages further increased to 378 (10 mg/L) and 360 (1 mg/L), compared to 236 in the blank group. Additionally, the positive correlation ratio (Fig. 6b) in the 10 mg/L group (53.5 % on day 4 and 54.0 % on day 10) was higher than in the blank (45.6 % and 46.6 %) and 1 mg/L (44.8 % and 49.4 %) groups. These results indicate that the connectivity within the natural symbiotic bacterial community increased under SMX stress.

#### 3.5.2. Functional predictions

The BugBase phenotype prediction results for natural symbiotic bacteria under different SMX concentrations are presented in Fig. 7a. The relative abundance of gram-negative and facultative anaerobic bacteria increased with SMX concentrations. Notably, biofilm-forming phenotypes also increased significantly, while the potentially pathogenic phenotypes decreased significantly (Fig. 7b). Additionally, KEGG pathway analysis (Fig. 7c) revealed that metabolic pathways had the highest relative share of 74.8 % at level 1, while the trends at level 2 were essentially consistent across all groups on days 4 and 10. Furthermore, level 2 comparisons revealed significant changes in bacterial physiological processes with increasing SMX concentrations. Specifically, pathways related to xeno-biotic degradation and metabolism, amino acid metabolism, and lipid metabolism increased, while those associated with sugar biosynthesis and metabolism, energy metabolism, and cellular motility decreased.



**Fig. 5.** Analysis of the composition of natural symbiotic bacterial communities under different SMX concentrations. (a) Shannon and Simpson indices of OUT level, (b) PCoA on OUT level, (c) Venn diagram of the genera, (d) Relative abundance of the genus level and (e) LEfSe multilevel species hierarchical cladogram. \*, \*\*, and \*\*\* indicate different significant differences p < 0.05, p < 0.01, and p < 0.001, respectively.



Fig. 6. One-way correlation network analysis of natural symbiotic bacterial communities at different time points under varying SMX concentrations. The nodes represent the species at the phylum level and linkages indicate a correlation between species  $(|\mathbf{r}| > 0.8)$ .

# 4. Discussion

### 4.1. SMX stress enhances Dunaliella sp. biomass via reduced bacterial competition and adaptive responses

This study confirmed that *Dunaliella* sp. showed a strong adaptation to SMX, as evidenced by their biomass, pigment content, and nutrient utilization efficiency. The microalgal biomass increased likely resulted from SMX restricting bacterial nutrient competition (Zhou et al., 2017), further supported by natural symbiotic bacterial isolation and culture experiments showing SMX inhibited bacterial ammonia and phosphate utilization. Since no nitrifying bacteria were detected in the natural symbiotic bacteria, NH<sup>+</sup><sub>4</sub>-N consumption was primarily attributed to bacterial assimilation (Kuypers et al., 2018). The order of nitrogen utilization by *Dunaliella* sp. was consistent with that of most microalgae: NH<sup>+</sup><sub>4</sub>-N > NO<sup>-</sup><sub>3</sub>-N > NO<sup>-</sup><sub>2</sub>-N (Perez-Garcia et al., 2011). Due to their competitive relationship, SMX restricted bacterial NH<sup>+</sup><sub>4</sub>-N utilization, resulting in NH<sup>+</sup><sub>4</sub>-N accumulation and promoting early biomass buildup in *Dunaliella* sp. Additionally, excess NH<sup>+</sup><sub>4</sub>-N and SMX may have suppressed nitrate reductase and nitrite reductase activities in microalgal cells, leading to a stagnation in NO<sup>-</sup><sub>3</sub>-N consumption (Su, 2021). Once the microalgae adapted, the early accumulated biomass became a key driver of subsequent growth, enabling rapid NO<sup>3</sup>-N utilization in the later phase.

X. Tan et al.

Environmental Technology & Innovation 38 (2025) 104171



Fig. 7. Analysis of BugBase phenotype prediction (a), comparison of phenotypic differences (b), and KEGG pathway on level 1 and level 2 (c) of natural symbiotic bacteria communities under different SMX concentrations. \*, \*\*, and \*\*\* indicate different significant differences p < 0.05, p < 0.01, and p < 0.001, respectively.

In addition, bacteria can cause diseases in microalgae, such as gall formation (Ashen Jon and Goff Lynda, 2000), bleaching disease (Zozaya-Valdés et al., 2017), and hole-rotten disease (Wang et al., 2008). The reduced abundance of pathogenic phenotypes in natural symbiotic bacteria under SMX exposure may further contribute to the increased biomass. Furthermore, pigments content in *Dunaliella* sp. strongly correlated with biomass and can be influenced by oxidative stress and photosynthetic processes under abiotic stress (Chen et al., 2025). Chla can participate in the removal of reactive oxygen species in chloroplasts (Kasahara et al., 2002), whereas carotenoids can effectively quench excited chlorophyll, thereby preventing stress-induced damage to the photosynthetic system (Paliwal et al., 2015). Photosynthesis is the primary pathway for pigments production. Studies have found that at high concentrations (5 mg/L), SMX still can enhance photosynthesis in *Chlorella vulgaris* biofilms by promoting nitrogen uptake (Sun et al., 2025). In this study, the increased N/P ratio in *Dunaliella* sp. cells under SMX exposure also indicates enhanced nitrogen uptake, which may reflect improved photosynthesis and could be related to enhanced extracellular protein secretion and synthesis of defensive proteins (Gao et al., 2016). These findings suggest that SMX stress induced adaptive responses and enhanced photosynthesis in *Dunaliella* sp., stimulating growth. However, the molecular mechanisms behind this remain to be further explored.

#### 4.2. Dunaliella sp. exhibits strong tolerance but poor removal ability to SMX

The excellent adaptability of *Dunaliella* sp. to high SMX concentrations is noteworthy. Excessive EPS production under stress is a key microbial defense strategy against toxic chemicals (Wang et al., 2021). Microalgae of the genus *Dunaliella* often increase EPS secretion in response to pollutant stress (Folgar et al., 2009, Wang et al., 2023). PS and PN are essential components of EPS due to their unique structures and functions (Wang et al., 2019). This study also observed that *Dunaliella* sp. produced significantly more PS and PN during the initial response to SMX, indicating that EPS enhancement is one of the key mechanisms for adapting to SMX stress. Interestingly, as cultivation progressed, the PS and PN content in the SMX-treated group declined.

EPS includes soluble EPS and bound EPS (Mao et al., 2022). Soluble EPS exhibits weaker interactions with cells and is more environmentally sensitive (Li et al., 2025). In bound EPS, PN can support bacterial aggregation, cell-to-cell signaling, and structural stability, while PS can promote microbial adhesion and biofilm formation (More et al., 2014). Additionally, EPS can enhance the conversion between inorganic and organic carbon, thereby supporting the growth of heterotrophic organisms and stabilizing the microalgae-bacteria consortium (Zhou et al., 2024). The increased diversity and connectivity of the natural symbiotic bacterial

community may benefit from changes in EPS properties. The decrease in EPS content may be related to changes in its properties, potentially enhancing interactions between *Dunaliella* sp. and its natural symbiotic bacteria.

The antioxidant system is also vital for microalgae adapting to antibiotic stress (Xiao et al., 2024). The intracellular antioxidant system removes excess ROS through coordinated enzymatic and non-enzymatic reactions. SOD, CAT, and GPX catalyze the conversion of  $O_2^-$  to  $H_2O_2$  and its breakdown into water (Gill and Tuteja, 2010), while GST removes harmful electrophilic substances (Zhang et al., 2022b). MDA is an important biomarker for the assessment of cellular oxidative damage (Zhang et al., 2022b). Changes in MDA content show that *Dunaliella* sp. is sensitive to short-term acute SMX damage but has strong recovery capabilities. Correspondingly, CAT activity rapidly and significantly increased and maintained high levels, indicating its crucial role in *Dunaliella* sp.'s adaptation to SMX stress. It is a common response in many microalgae under antibiotic stress (Chen et al., 2020b), chen et al., 2025), and a study indicates that CAT is vital for enhancing SMX adaptability in acclimated *C. vulgaris* (Zhang et al., 2022b), suggesting CAT activity as a potential biomarker for assessing microalgal adaptability to antibiotics.

GSH is essential for maintaining the cellular reduced state, serving as a substrate for both GPX and GST, while its balance is regulated by GR (Gill and Tuteja, 2010). By day 4, increased GSH consumption enhanced GR activity, while GSH primarily adapted to SMX stress through non-enzymatic pathways, as SMX inhibited GST and GPX. By day 10, reduced SOD and GR activities and increased GSH content indicated the recovery of *Dunaliella* sp.'s antioxidant system, consistent with decreased MDA content.

*Dunaliella* sp. showed good adaptation to high SMX concentrations, but its removal efficiency was lower than that of most microalgae, such as *Nannochloris sp.* (32 %; (Bai and Acharya, 2016)), *Monoraphidium contortum* (42.3 %; (Costa-Ramos et al., 2025)), and *Scenedesmus obliquus* (16–29 %; (Xiong et al., 2019a)). Currently, microalgae of the genus *Chlorella* has shown the highest SMX removal efficiency, with wild *Chlorella vulgaris* achieving 76 % removal and an acclimated strain reaching 92.5 % (Zhang et al., 2022b). Microalgae remove antibiotics mainly through biosorption, bioaccumulation, and biodegradation (Xiao et al., 2024). Biosorption is a key step, and the cell wall plays a crucial role in this process. The diverse functional groups on the cell wall can adsorb various pollutants, and the cell wall and EPS exhibit coordinated effects in pollutant removal (Naveed et al., 2025). The study found that *Dunaliella tertiolecta* and the cell wall-free mutant of *Chlamydomonas reinhardtii* formed smaller aggregates under nanoparticle exposure compared to cell wall-containing microalgae (Oukarroum et al., 2012, Perreault et al., 2012). The absence of a cell wall likely weakens the biosorption capacity of *Dunaliella* sp., resulting in lower SMX removal efficiency.

In addition, The absence of a cell wall is not necessarily linked to increased sensitivity in microalgae (Ge et al., 2022). Therefore, the cell membrane undoubtedly plays a more critical role in cell wall-free microalgae. The increase in MDA content indicates damage to the structure and function of *Dunaliella* sp.'s cell membrane, which may also affect its interaction with SMX. Microalgal surfaces are usually negatively charged (Günan Yücel et al., 2021), whereas SMX exists primarily in a stable anionic form at pH > 5.7 (Wang and Wang, 2018). Under seawater conditions (pH ~8.0), SMX is likely to experience electrostatic repulsion. Antibiotics with higher lip-ophilicity (log Kow) are more easily adsorbed by microalgae, while SMX, with a log Kow of 0.89, exhibits strong hydrophilicity (log Kow < 1) (Wang et al., 2022b). Thus, antibiotic removal efficiency is also influenced by *Dunaliella* sp.'s cell membrane characteristics, but deeper mechanisms need analysis via scanning electron microscopy (SEM) and Fourier-transform infrared spectroscopy (FTIR).

#### 4.3. Sulfamethoxazole alters the composition, connection and function of natural symbiotic bacterial communities

SMX is a broad-spectrum antibiotic that inhibits bacterial growth by competitively inhibiting dihydrofolate synthase and disrupting folate biosynthesis (Rodrigo et al., 2022). Functional prediction (Fig. 8) revealed reduced abundances of folate biosynthesis, ribosomes, flagellar assembly, bacterial chemotaxis, and the TCA cycle under SMX stress, all critical for bacterial growth, reproduction, and motility, and common targets of antibiotics (Xiu et al., 2017, Zhang et al., 2021, Yu et al., 2022, Xu et al., 2024). Consistent with



Fig. 8. Certain significantly different (p < 0.05) functions among the top 40 in relative abundance at KEGG pathway level 3 in natural symbiotic bacterial communities under different SMX concentrations.

other microalgal-bacterial consortia (Hu et al., 2022, Wang et al., 2022b), SMX significantly inhibited the growth of natural symbiotic bacteria associated with *Dunaliella* sp. (Fig. S2).

However, in complete contrast, the diversity of the natural symbiotic bacterial community significantly increased. Similar responses have been observed in bacterial communities under antibiotic stress in vegetable root systems (Ren et al., 2024) and vertical-flow constructed wetlands (Chen et al., 2020a). Recently, it was discovered for the first time that a symbiotic system of river-derived bacteria and *Chlorella vulgaris* exhibited increased diversity under SMX stress (Wang et al., 2024). Another distinction is that the enhanced connectivity in *Dunaliella* sp.'s natural symbiotic bacterial community under SMX stress indicates improved stability and adaptability to adverse conditions (Yuan et al., 2021).

The contrasting trends in diversity and connectivity between natural and foreign symbiotic bacterial communities may result from their distinct adaptation strategies. In foreign symbiotic consortia, host microalgae influence community composition more strongly than environmental factors through selectively recruiting specific bacteria (Kimbrel et al., 2019). Additionally, the decline in diversity is associated with the dominance of specific antibiotic-degrading bacteria (Xie et al., 2020), which are maintained through the accumulation and transfer of resistance genes (da da Silva Rodrigues et al., 2020).

In contrast, natural symbiotic consortia have stable species numbers, indicating weaker host selection pressure. From the perspective of specific species changes, *Labrenzia* has multiple pollutant resistance genes and exhibits strong aromatic compound degradation capabilities (Rodrigues Gisele et al., 2018). However, its abundance decreased, and no significant changes were observed in degradation-related pathways, suggesting that natural symbiotic bacteria may rely less on degradation strategies. Additionally, *Methylophaga* can promote nitrogen fixation and increase under nitrogen-limited conditions (Shi et al., 2018). However, it relies on phytoplankton for carbon and energy and competes for cobalamin (Bertrand et al., 2015). Its decline suggests poor adaptation to SMX stress, but it may also act as an adjustment strategy, creating growth opportunities for other bacteria. Notably, the abundance of *Ponticoccus* and *Mameliella* increased, and these bacteria are characterized by active quorum sensing (QS) functions or the presence of related genes (Danish-Daniel et al., 2016, Chi et al., 2017).

QS-mediated interactions is an important factor in shaping bacterial community structure and regulating ecological behaviors such as biofilm formation and antibiotic resistance (Zhou et al., 2016). Biofilm formation is an important bacterial adaptation strategy to environmental stresses because it can protect bacterial cells and enhance their resistance (Wu et al., 2015). The increase in QS function and biofilm formation phenotype in this study suggest that QS bacteria likely contribute to enhanced community connectivity by regulating biofilm formation. Additionally, the upregulation of amino acid metabolism in natural symbiotic bacteria reflects both active adaptation to SMX stress and a stronger interaction with microalgae, facilitated by increased EPS secretion (Wang et al., 2018). As a medium for metabolic exchange and signal communication, EPS serves as a critical link in enhancing the adaptability of *Dunaliella* sp. and its natural symbiotic bacteria to SMX stress.

#### 4.4. Application prospects and challenges

In this study , most response trends generally follow a gradient pattern under different SMX concentration stresses, indicating that the symbiotic consortia can maintain stable adaptation mechanisms across a wide range of SMX levels. Although the symbiotic consortia show limited pollutant removal efficiency, its broad adaptability makes it a practical option for nutrient recovery in aquaculture wastewater with high antibiotic contamination risks. Its low adsorption and bioaccumulation characteristics also enhance its suitability as fishery feed, reducing source pollution risks while ensuring food safety. In antibiotic remediation, the biofilms from *Dunaliella* sp. symbiotic systems show great potential for integration with high-performance microalgae or degradative bacteria, which is conducive to enhancing the system's tolerance and stability. The response patterns of natural symbiotic bacteria provide guidance for foreign bacteria selection and community construction, which are beneficial for enhancing practical application potential.

Additionally, this study still has certain limitations: (1) The absence of a cell wall in *Dunaliella* sp. poses technical challenges for extracting bound EPS, limiting in-depth research on its role. (2) Considering the presence of symbiotic bacteria provides a more realistic understanding of microalgae's adaptation strategies, but understanding axenic microalgae's intrinsic traits is also crucial. However, removing symbiotic bacteria without stressing microalgae remains a major technical challenge. (3) Nutrient contents and microalgae growth determined the cultivation period. For practical applications, cost and efficiency factors require more efficient cultivation methods to improve feasibility. (4) Actual aquaculture wastewater may contain not only antibiotics but also risks of combined pollution from heavy metals and organic compounds, and environmental factors such as dissolved oxygen, pH, and salinity could also impact the symbiotic system. These are all important factors that cannot be overlooked in subsequent research and practical applications.

#### 5. Conclusion

This study revealed that the *Dunaliella* sp. and its natural symbiotic bacterial consortium exhibited strong tolerance to SMX but weak removal efficiency. QS bacteria-driven community changes and biofilm formation enhance the diversity and connectivity of natural symbiotic bacterial communities. *Dunaliella* sp.'s high tolerance relies on strong antioxidant capacity (with CAT playing a key role) and elevated EPS secretion. EPS plays a key role in enhancing the adaptability of the consortium by strengthening the interaction between *Dunaliella* sp. and its natural symbiotic bacteria. However, the mechanisms regulating their interactions require further investigation. These findings strengthen our understanding of the responses of natural symbiotic bacteria to antibiotics, providing new insights for improving the adaptability of microalgae-bacteria symbiosis technology and the construction of bacterial communities.

#### CRediT authorship contribution statement

**Zhao Hui:** Writing – review & editing, Supervision, Resources, Methodology, Funding acquisition, Conceptualization. **Pan Gang:** Writing – review & editing, Conceptualization. **Che Wenxue:** Visualization, Software, Methodology. **Tan Xiao:** Writing – original draft, Visualization, Validation, Software, Methodology, Investigation, Formal analysis, Data curation. **Li Jinting:** Investigation. **Man Ying:** Methodology, Conceptualization. **Wang Liyi:** Investigation.

#### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

The funding received for this research includes National Natural Science Foundation of China (No. 42076162) and Innovation Group Project of Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai) (No. 311020004).

#### Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.eti.2025.104171.

#### Data availability

Data will be made available on request.

#### References

- Adenaya, A., Berger, M., Brinkhoff, T., Ribas-Ribas, M., Wurl, O., 2023. Usage of antibiotics in aquaculture and the impact on coastal waters. Mar. Pollut. Bull. 188, 114645. https://doi.org/10.1016/j.marpolbul.2023.114645.
- Ashen Jon, B., Goff Lynda, J., 2000. Molecular and ecological evidence for species specificity and coevolution in a group of marine algal-bacterial symbioses. Appl. Environ. Microbiol 66 (7), 3024–3030. https://doi.org/10.1128/AEM.66.7.3024-3030.2000.
- Bai, X., Acharya, K., 2016. Removal of trimethoprim, sulfamethoxazole, and triclosan by the green alga Nannochloris sp. J. Hazard. Mater. 315, 70–75. https://doi.org/10.1016/j.jhazmat.2016.04.067.
- Bertrand, E.M., McCrow, J.P., Moustafa, A., Zheng, H., McQuaid, J.B., Delmont, T.O., Allen, A.E., 2015. Phytoplankton–bacterial interactions mediate micronutrient colimitation at the coastal Antarctic sea ice edge. Proc. Natl. Acad. Sci. 112 (32), 9938–9943. https://doi.org/10.1073/pnas.1501615112.

Chae, Y., Kim, D., An, Y.-J., 2019. Effects of micro-sized polyethylene spheres on the marine microalga Dunaliella salina: Focusing on the algal cell to plastic particle size ratio. Aquat. Toxicol. 216, 105296. https://doi.org/10.1016/j.aquatox.2019.105296.

Chandel, N., Ahuja, V., Gurav, R., Kumar, V., Tyagi, V.K., Pugazhendhi, A., Bhatia, S.K., 2022. Progress in microalgal mediated bioremediation systems for the removal of antibiotics and pharmaceuticals from wastewater. Sci. Total Environ. 825, 153895. https://doi.org/10.1016/j.scitotenv.2022.153895.

Chen, J., Tong, T., Jiang, X., Xie, S., 2020a. Biodegradation of sulfonamides in both oxic and anoxic zones of vertical flow constructed wetland and the potential degraders. Environ. Pollut. 265, 115040. https://doi.org/10.1016/j.envpol.2020.115040.

- Chen, J., Qin, B., Han, Y., Yang, S., 2025. Sulfamethoxazole and sulfaisoxazole stress on growth, photosynthesis, cell structure and oxidation system in *Microcystis* aeruginosa. Algal Res. 87, 103880. https://doi.org/10.1016/j.algal.2024.103880.
- Chen, S., Zhang, W., Li, J., Yuan, M., Zhang, J., Xu, F., Wang, L., 2020b. Ecotoxicological effects of sulfonamides and fluoroquinolones and their removal by a green alga (*Chlorella vulgaris*) and a cyanobacterium (*Chrysosporum ovalisporum*). Environ. Pollut. 263, 114554. https://doi.org/10.1016/j.envpol.2020.114554.
- Chi, W., Zheng, L., He, C., Han, B., Zheng, M., Gao, W., Gao, X., 2017. Quorum sensing of microalgae associated marine *Ponticoccus* sp. PD-2 and its algicidal function regulation. AMB Express 7 (1), 59. https://doi.org/10.1186/s13568-017-0357-6.
- Costa-Ramos, L.T.A., Senra, M.V.X., Nogueira, G.H.S., Ramalho-Júnior, R.R., Andrade, S.J., Santo, E.E., Matsudo, M.C., 2025. Unveiling the antibiotics removal ability of *Monoraphidium contortum*. Biochem. Eng. J. 217, 109686. https://doi.org/10.1016/j.bej.2025.109686.
- Danish-Daniel, M., Han Ming, G., Noor, M.E.M., Yeong, Y.S., Usup, G., 2016. Draft genome sequence of Mameliella alba strain UMTAT08 isolated from clonal culture of toxic dinoflagellate Alexandrium tamiyavanichii. Genom. Data 10, 12–14. https://doi.org/10.1016/j.gdata.2016.08.015.

Folgar, S., Torres, E., Pérez-Rama, M., Cid, A., Herrero, C., Abalde, J., 2009. Dunaliella salina as marine microalga highly tolerant to but a poor remover of cadmium. J. Hazard. Mater. 165 (1), 486–493. https://doi.org/10.1016/j.jhazmat.2008.10.010.

- Gao, Y., Lim, T.K., Lin, Q., Li, S.F.Y., 2016. Identification of cypermethrin induced protein changes in green algae by iTRAQ quantitative proteomics. J. Proteom. 139, 67–76. https://doi.org/10.1016/j.jprot.2016.03.012.
- Ge, J., Yang, Q., Fang, Z., Liu, S., Zhu, Y., Yao, J., Guan, W., 2022. Microplastics impacts in seven flagellate microalgae: Role of size and cell wall. Environ. Res. 206, 112598. https://doi.org/10.1016/j.envres.2021.112598.

Gill, S.S., Tuteja, N., 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. Plant Physiol. Biochem 48 (12), 909–930. https://doi.org/10.1016/j.plaphy.2010.08.016.

- Günan Yücel, H., Ertuğrul Karatay, S., Aksu, Z., Dönmez, G., 2021. Lithium (I) biofortified Dunaliella salina as a potential functional nutrition supplement. Algal Res 56, 102257. https://doi.org/10.1016/j.algal.2021.102257.
- Hom-Diaz, A., Jaén-Gil, A., Rodríguez-Mozaz, S., Barceló, D., Vicent, T., Blánquez, P., 2022. Insights into removal of antibiotics by selected microalgae (Chlamydomonas reinhardtii, Chlorella sorokiniana, Dunaliella tertiolecta and Pseudokirchneriella subcapitata). Algal Res 61, 102560. https://doi.org/10.1016/j. algal.2021.102560.
- Hu, G., Fan, S., Wang, H., Ji, B., 2022. Adaptation responses of microalgal-bacterial granular sludge to sulfamethoxazole. Bioresour. Technol. 364, 128090. https:// doi.org/10.1016/j.biortech.2022.128090.
- Kasahara, M., Kagawa, T., Oikawa, K., Suetsugu, N., Miyao, M., Wada, M., 2002. Chloroplast avoidance movement reduces photodamage in plants. Nature 420 (6917), 829–832. https://doi.org/10.1038/nature01213.

- Kimbrel, J.A., Samo, T.J., Ward, C., Nilson, D., Thelen, M.P., Siccardi, A., Mayali, X., 2019. Host selection and stochastic effects influence bacterial community assembly on the microalgal phycosphere. Algal Res. 40, 101489. https://doi.org/10.1016/j.algal.2019.101489.
- Kuypers, M.M.M., Marchant, H.K., Kartal, B., 2018. The microbial nitrogen-cycling network. Nat. Rev. Microbiol 16 (5), 263–276. https://doi.org/10.1038/ nrmicro.2018.9.
- Kvíderová, J., Henley, W.J., 2005. The effect of ampicillin plus streptomycin on growth and photosynthesis of two halotolerant chlorophyte algae. J. Appl. Phycol. 17 (4), 301–307. https://doi.org/10.1007/s10811-005-7293-6.
- Le, T.X., Munekage, Y., 2004. Residues of selected antibiotics in water and mud from shrimp ponds in mangrove areas in Viet Nam. Mar. Pollut. Bull. 49 (11), 922–929. https://doi.org/10.1016/j.marpolbul.2004.06.016.
- Li, S., Bai, Y., Li, Z., Wang, A., Ren, N.-Q., Ho, S.-H., 2025. Overlooked role of extracellular polymeric substances in antibiotic-resistance gene transfer within microalgae-bacteria system. J. Hazard. Mater. 488, 137206. https://doi.org/10.1016/j.jhazmat.2025.137206.
- Liska, A.J., Shevchenko, A., Pick, U., Katz, A., 2004. Enhanced photosynthesis and redox energy production contribute to salinity tolerance in dunaliella as revealed by homology-based proteomics. Plant Physiol. 136 (1), 2806–2817. https://doi.org/10.1104/pp.104.039438.
- Lu, Y., Zhou, X., Zheng, Y., Yang, H., Cao, W., 2025. How far do we still need to go with antibiotics in aquatic environments? Antibiotic occurrence, chemical-free or chemical-limited strategies, key challenges, and future perspectives. Water Res 275, 123179. https://doi.org/10.1016/j.watres.2025.123179.
- Machado, M.D., Soares, E.V., 2019. Sensitivity of freshwater and marine green algae to three compounds of emerging concern. J. Appl. Phycol. 31 (1), 399–408. https://doi.org/10.1007/s10811-018-1511-5.
- Mao, J., Gu, Z., Zhang, S., An, X., Lan, H., Liu, H., Qu, J., 2022. Protonated carbon nitride elicits microalgae for water decontamination. Water Res 222, 118955. https://doi.org/10.1016/j.watres.2022.118955.
- More, T.T., Yadav, J.S.S., Yan, S., Tyagi, R.D., Surampalli, R.Y., 2014. Extracellular polymeric substances of bacteria and their potential environmental applications. J. Environ. Manag. 144, 1–25. https://doi.org/10.1016/j.jenvman.2014.05.010.
- Naveed, S., Yu, Q., Szewczuk-Karpisz, K., Zhang, C., Rahman, S.-U., Ge, Y., 2025. Roles of extracellular polymeric substances in arsenic accumulation and
- detoxification by cell wall intact and mutant strains of *Chlamydomonas reinhardtii*. J. Environ. Sci. 152, 142–154. https://doi.org/10.1016/j.jes.2024.03.048. Niu, Z.-G., Zhang, K., Zhang, Y., 2016. Occurrence and distribution of antibiotic resistance genes in the coastal area of the Bohai Bay, China. Mar. Pollut. Bull. 107 (1), 245–250. https://doi.org/10.1016/j.marpolbul.2016.03.064.
- Oukarroum, A., Bras, S., Perreault, F., Popovic, R., 2012. Inhibitory effects of silver nanoparticles in two green algae, Chlorella vulgaris and Dunaliella tertiolecta. Ecotoxicol. Environ. Saf. 78, 80–85. https://doi.org/10.1016/j.ecoenv.2011.11.012.
- Paliwal, C., Pancha, I., Ghosh, T., Maurya, R., Chokshi, K., Vamsi Bharadwaj, S.V., Mishra, S., 2015. Selective carotenoid accumulation by varying nutrient media and salinity in *Synechocystis* sp. CCNM 2501. Bioresour. Technol. 197, 363–368. https://doi.org/10.1016/j.biortech.2015.08.122.
- Peng, T., Song, B., Wang, Y., Yuan, J., Yang, Z., Tang, L., 2025. Trophic transfer of sulfonamide antibiotics in aquatic food chains: a comprehensive review with a focus on environmental health risks. Environ. Pollut. 369, 125823. https://doi.org/10.1016/j.envpol.2025.125823.
- Perez-Garcia, O., Escalante, F.M.E., de-Bashan, L.E., Bashan, Y., 2011. Heterotrophic cultures of microalgae: Metabolism and potential products. Water Res 45 (1), 11–36. https://doi.org/10.1016/j.watres.2010.08.037.
- Perreault, F., Bogdan, N., Morin, M., Claverie, J., Popovic, R., 2012. Interaction of gold nanoglycodendrimers with algal cells (*Chlamydomonas reinhardtii*) and their effect on physiological processes. Nanotoxicology 6 (2), 109–120. https://doi.org/10.3109/17435390.2011.562325.
- Quesnel, D.M., Bhaskar, I.M., Gieg, L.M., Chua, G., 2011. Naphthenic acid biodegradation by the unicellular alga Dunaliella tertiolecta. Chemosphere 84 (4), 504–511. https://doi.org/10.1016/j.chemosphere.2011.03.012.
- Ren, J., Lu, H., Lu, S., Huang, Z., 2024. Impacts of sulfamethoxazole stress on vegetable growth and rhizosphere bacteria and the corresponding mitigation mechanism. Front. Bioeng. Biotechnol. 12. https://doi.org/10.3389/fbioe.2024.1303670.
- Rodrigo, M.K.D., Saiganesh, A., Hayes, A.J., Wilson, A.M., Anstey, J., Pickering, J.L., Barnett, T.C., 2022. Host-dependent resistance of Group A Streptococcus to sulfamethoxazole mediated by a horizontally-acquired reduced folate transporter. Nat. Commun. 13 (1), 6557. https://doi.org/10.1038/s41467-022-34243-3.
- Rodrigues Gisele, N., Lago-Lestón, A., Costa, R., Keller-Costa, T., 2018. Draft genome sequence of *Labrenzia* sp. Strain EL143, a coral-associated alphaproteobacterium with versatile symbiotic living capability and strong halogen degradation potential. Genome Announc. 6 (10). https://doi.org/10.1128/genomea.00132-18.
- Shi, F., Wei, X., Feng, J., Sun, Y., Zhu, L., 2018. Variation of bacterial community associated with *Phaeodactylum tricornutum* in response to different inorganic nitrogen concentrations. Acta Oceanol. Sin. 37 (12), 118–128. https://doi.org/10.1007/s13131-018-1272-7.
- da Silva Rodrigues, D.A., da Cunha, C.C.R.F., Freitas, M.G., de Barros, A.L.C., e Castro, P.B.N., Pereira, A.R., de Cássia Franco Afonso, R.J., 2020. Biodegradation of sulfamethoxazole by microalgae-bacteria consortium in wastewater treatment plant effluents. Sci. Total Environ. 749, 141441. https://doi.org/10.1016/j. scitotenv.2020.141441.
- Su, Y., 2021. Revisiting carbon, nitrogen, and phosphorus metabolisms in microalgae for wastewater treatment. Sci. Total Environ. 762, 144590. https://doi.org/ 10.1016/j.scitotenv.2020.144590.
- Sun, F., Jiao, Y., Liang, S., Zhuang, L.L., Zhang, J., 2025. The effect of sulfamethoxazole on the growth of microalgae biofilms and the internal transportation and transformation of nutrients in the biofilm. Environ. Res. 273, 121232. https://doi.org/10.1016/j.envres.2025.121232.
- Wang, G., Shuai, L., Li, Y., Lin, W., Zhao, X., Duan, D., 2008. Phylogenetic analysis of epiphytic marine bacteria on Hole-Rotten diseased sporophytes of Laminaria japonica. J. Appl. Phycol. 20 (4), 403–409. https://doi.org/10.1007/s10811-007-9274-4.
- Wang, H., Qi, B., Jiang, X., Jiang, Y., Yang, H., Xiao, Y., Wang, W., 2019. Microalgal interstrains differences in algal-bacterial biofloc formation during liquid digestate treatment. Bioresour. Technol. 289, 121741. https://doi.org/10.1016/j.biortech.2019.121741.
- Wang, J., Wang, S., 2018. Microbial degradation of sulfamethoxazole in the environment. Appl. Microbiol. Biotechnol. 102 (8), 3573–3582. https://doi.org/10.1007/s00253-018-8845-4.
- Wang, J., Guo, H., Cao, C., Zhao, W., Kwok, L.-Y., Zhang, H., Zhang, W., 2018. Characterization of the adaptive amoxicillin resistance of *Lactobacillus casei* zhang by proteomic analysis, 9. https://doi.org/10.3389/fmicb.2018.00292.
- Wang, Q., Geng, L., Gao, Z., Sun, Y., Li, X., Sun, S., Luo, Y., 2024. Microalgae enhances the adaptability of epiphytic bacteria to sulfamethoxazole stress and proliferation of antibiotic resistance genes mediated by integron. Environ. Sci. Technol. 58 (43), 19397–19407. https://doi.org/10.1021/acs.est.4c04925.
- Wang, S., Ji, B., Cui, B., Ma, Y., Guo, D., Liu, Y., 2021. Cadmium-effect on performance and symbiotic relationship of microalgal-bacterial granules. J. Clean. Prod. 282, 125383. https://doi.org/10.1016/j.jclepro.2020.125383.
- Wang, S.C., Liu, G.Z., Liu, F.F., 2023. Physiological and metabolic toxicity of polystyrene microplastics to Dunaliella salina. Environ. Pollut. 316, 120544. https://doi. org/10.1016/j.envpol.2022.120544.
- Wang, X., Lin, Y., Zheng, Y., Meng, F., 2022a. Antibiotics in mariculture systems: A review of occurrence, environmental behavior, and ecological effects. Environ. Pollut. 293, 118541. https://doi.org/10.1016/j.envpol.2021.118541.
- Wang, Y., Li, J., Lei, Y., Li, X., Nagarajan, D., Lee, D.-J., Chang, J.-S., 2022b. Bioremediation of sulfonamides by a microalgae-bacteria consortium analysis of pollutants removal efficiency, cellular composition, and bacterial community. Bioresour. Technol. 351, 126964. https://doi.org/10.1016/j. biortech.2022.126964.
- Wu, H., Moser, C., Wang, H.-Z., Høiby, N., Song, Z.-J., 2015. Strategies for combating bacterial biofilm infections. Int. J. Oral. Sci. 7 (1), 1–7. https://doi.org/10.1038/ ijos.2014.65.
- Xiao, Z., Meng, H., Li, S., Ning, W., Song, Y., Han, J., Ho, S.-H., 2024. Insights into the removal of antibiotics from livestock and aquaculture wastewater by algaebacteria symbiosis systems. Environ. Res. 257, 119326. https://doi.org/10.1016/j.envres.2024.119326.
- Xie, B., Tang, X., Ng, H.Y., Deng, S., Shi, X., Song, W., Liang, H., 2020. Biological sulfamethoxazole degradation along with anaerobically digested centrate treatment by immobilized microalgal-bacterial consortium: performance, mechanism and shifts in bacterial and microalgal communities. Chem. Eng. J. 388, 124217. https://doi.org/10.1016/j.cej.2020.124217.

- Xiong, J.Q., Kurade, M.B., Abou-Shanab, R.A.I., Ji, M.K., Choi, J., Kim, J.O., Jeon, B.H., 2016. Biodegradation of carbamazepine using freshwater microalgae *Chlamydomonas mexicana* and *Scenedesmus obliquus* and the determination of its metabolic fate. Bioresour. Technol. 205, 183–190. https://doi.org/10.1016/j. biortech.2016.01.038.
- Xiong, J.Q., Kurade, M.B., Kim, J.R., Roh, H.S., Jeon, B.H., 2017. Ciprofloxacin toxicity and its co-metabolic removal by a freshwater microalga Chlamydomonas mexicana. J. Hazard. Mater. 323, 212–219. https://doi.org/10.1016/j.jhazmat.2016.04.073.
- Xiong, J.Q., Kim, S.J., Kurade, M.B., Govindwar, S., Abou-Shanab, R.A.I., Kim, J.R., Jeon, B.H., 2019b. Combined effects of sulfamethazine and sulfamethoxazole on a freshwater microalga, *Scenedesmus obliquus*: toxicity, biodegradation, and metabolic fate. J. Hazard. Mater. 370, 138–146. https://doi.org/10.1016/j. ibazmat.2018.07.049.
- Xiong, J.Q., Govindwar, S., Kurade, M.B., Paeng, K.J., Roh, H.S., Khan, M.A., Jeon, B.H., 2019a. Toxicity of sulfamethazine and sulfamethoxazole and their removal by a green microalga, *Scenedesmus obliquus*. Chemosphere 218, 551–558. https://doi.org/10.1016/j.chemosphere.2018.11.146.
- Xiong, Q., Liu, Y.S., Hu, L.X., Shi, Z.Q., Ying, G.G., 2020. Levofloxacin and sulfamethoxazole induced alterations of biomolecules in *Pseudokirchneriella subcapitata*. Chemosphere 253, 126722. https://doi.org/10.1016/j.chemosphere.2020.126722.
- Xiu, P., Liu, R., Zhang, D., Sun, C., 2017. Pumilacidin-like lipopeptides derived from marine bacterium Bacillus sp. Strain 176 suppress the motility of vibrio alginolyticus. Appl. Environ. Microbiol 83 (12), e00450–00417. https://doi.org/10.1128/AEM.00450-17.
- Xu, Q., Ali, S., Afzal, M., Nizami, A.-S., Han, S., Dar, M.A., Zhu, D., 2024. Advancements in bacterial chemotaxis: utilizing the navigational intelligence of bacteria and its practical applications. Sci. Total Environ. 931, 172967. https://doi.org/10.1016/j.scitotenv.2024.172967.
- Yu, C., Armengaud, J., Blaustein, R.A., Chen, K., Ye, Z., Xu, F., Shen, C., 2022. Antibiotic tolerance and degradation capacity of the organic pollutant-degrading bacterium *Rhodococcus biphenylivorans* TG9T. J. Hazard. Mater. 424, 127712. https://doi.org/10.1016/j.jhazmat.2021.127712.
- Yuan, M.M., Guo, X., Wu, L., Zhang, Y., Xiao, N., Ning, D., Zhou, J., 2021. Climate warming enhances microbial network complexity and stability. Nat. Clim. Change 11 (4), 343–348. https://doi.org/10.1038/s41558-021-00989-9.
- Zhang, H., Duan, L., Li, S., Gao, Q., Li, M., Xing, F., Zhao, Y., 2025. Efficient sulfamethoxazole removal, water recovery and electricity generation were achieved simultaneously by osmotic microbial fuel cell in treating antibiotic wastewater. Sep. Purif. Technol. 352, 128157. https://doi.org/10.1016/j. seppur.2024.128157.
- Zhang, L., He, J., Bai, L., Ruan, S., Yang, T., Luo, Y., 2021. Ribosome-targeting antibacterial agents: advances, challenges, and opportunities. Med. Res. Rev. 41 (4), 1855–1889. https://doi.org/10.1002/med.21780.
- Zhang, Q., Li, N., Wang, B.L., Zhang, T., Ma, P.F., Zhang, W.X., Ying, L., 2022a. Capabilities and mechanisms of microalgae on nutrients and florfenicol removing from marine aquaculture wastewater. J. Environ. Manag. 320, 115673. https://doi.org/10.1016/j.jenvman.2022.115673.
- Zhang, Y., Wan, J., Li, Z., Wu, Z., Dang, C., Fu, J., 2022b. Enhanced removal efficiency of sulfamethoxazole by acclimated microalgae: tolerant mechanism, and transformation products and pathways. Bioresour. Technol. 347, 126461. https://doi.org/10.1016/j.biortech.2021.126461.
- Zhou, D., Zhang, C., Fu, L., Xu, L., Cui, X., Li, Q., Crittenden, J.C., 2017. Responses of the microalga Chlorophyta sp. to bacterial quorum sensing molecules (N-Acylhomoserine Lactones): aromatic protein-induced self-aggregation. Environ. Sci. Technol. 51 (6), 3490–3498. https://doi.org/10.1021/acs.est.7b00355.
- Zhou, J., Lyu, Y., Richlen, M.L., Anderson, D.M., Cai, Z., 2016. Quorum sensing is a language of chemical signals and plays an ecological role in algal-bacterial interactions. Crit. Rev. Plant Sci. 35 (2), 81–105. https://doi.org/10.1080/07352689.2016.1172461.
- Zhou, Y., Cui, X., Wu, B., Wang, Z., Liu, Y., Ren, T., Rittmann, B.E., 2024. Microalgal extracellular polymeric substances (EPS) and their roles in cultivation, biomass harvesting, and bioproducts extraction. Bioresour. Technol. 406, 131054. https://doi.org/10.1016/j.biortech.2024.131054.
- Zhu, X., Shen, C., Huang, J., Wang, L., Pang, Q., Peng, F., Xu, B., 2022. The effect of sulfamethoxazole on nitrogen removal and electricity generation in a tidal flow constructed wetland coupled with a microbial fuel cell system: Microbial response. Chem. Eng. J. 431, 134070. https://doi.org/10.1016/j.cej.2021.134070.
- Zozaya-Valdés, E., Roth-Schulze, A.J., Egan, S., Thomas, T., 2017. Microbial community function in the bleaching disease of the marine macroalgae Delisea pulchra. Environ. Microbiol 19 (8), 3012–3024. https://doi.org/10.1111/1462-2920.13758.