

Est.
1841

YORK
ST JOHN
UNIVERSITY

Springham, Matthew, Singh, Nav, Stewart, Perry, Jones, Ian, Norton-Sherwood, Charlie, May, Dominic, Salter, Jamie, Strudwick, Anthony J. and Shaw, Joseph W. (2025) The moderating effects of speed, strength and endurance capacities on match induced neuromuscular fatigue in U-18 English premier league academy football players: A hypothesis-generating case report. *International Journal of Sports Science & Coaching*.

Downloaded from: <https://ray.yorks.ac.uk/id/eprint/11976/>

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version:

<https://doi.org/10.1177/17479541251333927>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repositories Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at
ray@yorks.ac.uk

The moderating effects of speed, strength and endurance capacities on match induced neuromuscular fatigue in U-18 English premier league academy football players: A hypothesis-generating case report

International Journal of Sports Science
& Coaching
1–9

© The Author(s) 2025



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/17479541251333927

journals.sagepub.com/home/spo



Matthew Springham^{1,2} , Nav Singh², Perry Stewart², Ian Jones², Charlie Norton-Sherwood², Dominic May³, Jamie Salter⁴ , Anthony J. Strudwick⁵, and Joseph W. Shaw¹

Abstract

Purpose: To identify new hypotheses relating to the moderating effects of speed, strength and endurance capacities on match-induced neuromuscular fatigue. **Methods** 14 U-18 outfield players from one EPL academy team completed countermovement jump (CMJ) and isometric adductor (IADS) and posterior chain (IPCS) strength tests one day before match day (MD) (i.e., MD-1) and on MD + 2 around 8 competitive games. Explosive strength (CMJ), reactive strength (30 cm drop jump (DJ)), speed (10 m and 30 m), maximal strength (isometric mid-thigh pull (IMTP)) and endurance (1 km Time Trial (TT)) were measured at 4 time points across the season. Relationships between the average match induced change to each NMF measure and the average of each physical capacity measure were examined using Pearson's R (r) (when normally distributed) or Spearman's rank (ρ) (when not normally distributed) correlation coefficients. **Results:** Moderate positive relationships were observed between CMJ-JH change (δ) and DJ-RSI ($\rho = 0.36$; $p = 0.16$) and IMPT ($r = 0.46$; $p = 0.06$). Small negative relationships were observed between IADS-PF δ and 10 m speed ($\rho = 0.27$; $p = 0.29$), 30 m speed ($r = 0.22$; $p = 0.41$), CMJ-JH ($\rho = 0.29$; $p = 0.26$) and DJ-RSI ($\rho = 0.36$; $p = 0.16$), and between IPCS-PF δ and 10 m speed ($\rho = 0.20$; $p = 0.45$) and 1 km TT ($\rho = 0.26$; $p = 0.33$). **Conclusions:** These results generate important hypotheses relating to the potential mitigating effects that reactive strength, maximal strength and endurance capacities might exert on match induced NMF in U-18 academy football players. These relationships warrant further investigation in larger research designs spanning both older and younger player age groups (i.e., across other youth development and professional development phase players) and competition levels (i.e., elite and sub-elite players).

Keywords

Drop jump, Isometric mid-thigh pull, physical capacity testing, soccer, stretch shortening cycle

Reviewers: Chris Barnes (CB Sports Performance Ltd, UK)
Filipe Clemente (Polytechnic Institute of Viana do Castelo, Portugal)
Nuno Nunes (Solent University, UK)

¹Faculty of Sport, Allied Health and Performance Science, St Mary's University, UK

²Arsenal Performance and Research Team, Arsenal Football Club, London, UK

³Brighton and Hove Albion Football Club, Brighton, UK

⁴School of Sport, York St John University, York, UK

⁵Sport Science Department, West Bromwich Albion Football Club, West Midlands, UK

Corresponding author:

Matthew Springham, Faculty of Sport, Allied Health and Performance Science, Office R5 Faculty of Sport, St Mary's University, UK; Arsenal Performance and Research Team, Arsenal Football Club, London, UK. Email: matt.springham@stmarys.ac.uk

Introduction

Neuromuscular fatigue (NMF; i.e., a temporary reduction to the maximal force-generating capacity of muscle) is commonly implicated in the aetiology of injury risk in young football players.^{1,2} Indeed, a recent four-year prospective study reported that U-18 and U-19 year old academy football players had the highest injury burden of all academy age groups and that NMF was a causal factor for 64%, 49% and 27% of hamstring, adductor and quadriceps injuries in U-18 academy football players² respectively. The link between NMF and muscle injury risk might relate to fatigue-induced impairments in force production, reduced eccentric strength, and compromised muscle stiffness regulation, which collectively, might increase susceptibility to strain and tissue damage.³ Indeed, fatigued muscle might be less able to tolerate high-force muscle contractions and mechanical load, increasing the risk of strain injury, particularly during high-intensity match actions. Consequently, neuromuscular performance tests including the counter-movement jump (CMJ) and isometric posterior chain (IPCS) and adductor (IADS) strength tests are commonly used to track the neuromuscular status of players around games across the competitive season. Interventions can then be put in place (i.e., to reduce training and match loads) in order to help mitigate injury risk, with 'normal' training loads resuming following the return to baseline performance levels.

Improving the physical capacities that relate to football match demands (i.e., speed, strength and endurance) might improve match play physical performance potential and, in-turn, player 'readiness' (i.e., denoting the interplay between player 'fitness' and 'fatigue'^{4,5}). Conceptually, this might help to reduce the relative burden of match play and consequently, the magnitude of match induced NMF.^{4,5} Indeed, two previous investigations reported that greater endurance (i.e., Yo-YoIR1 performance) and strength (i.e., isometric mid-thigh pull; IMTP performance) capacities were associated with higher match loads and reduced match induced NMF in academy rugby league players.^{6,7} Though rugby league and football share similar high-intensity intermittent running demands, rugby league typically involves more frequent collisions and prolonged periods of high-force isometric and quasi-isometric actions (i.e., during tackling and contact scenarios), which might result in different NMF responses compared to football. In contrast, academy football match play typically involves a larger volume of acceleration, deceleration, high-speed and change of direction activity, resulting in a large exposure to high intensity eccentric muscle actions. While endurance and strength have been shown to mitigate match-induced NMF in rugby league,^{6,7} the extent to which these physical qualities influence NMF in academy football remains unclear. To date, only Constantine and colleagues⁸ have reported that the greater match running distances

achieved by 'stronger' (IPCS) U-18 academy football players were accompanied by greater and longer-lasting periods of NMF; attributed to the accrual of additional muscle damage.

Despite there being some research evidence available to describe the moderating effect that strength exerts on match induced NMF in U-18 academy football players,⁸ the moderating effects of speed, explosive strength, reactive strength, and endurance capacities have received no research attention in this age group. These physical qualities represent key determinants of performance in football and conceptually, could moderate NMF responses. For example, speed and explosive strength underpin high-intensity football actions including accelerating, sprinting and change of direction, which contribute substantially to neuromuscular strain. Reactive strength, (i.e., the athlete's ability to efficiently utilise the stretch-shortening cycle (SSC)), may influence the extent to which players can tolerate mechanical loads associated with these match actions. Additionally, it is well established that endurance capacity is fundamental for sustaining high-intensity efforts across a full match.⁹ Consequently, it might help to mitigate NMF by virtue of improving metabolic efficiency and recovery between high-intensity efforts.

Understanding how physical capacities interact with match-induced NMF in academy football players is particularly important owing to the fatigue related injury risk in this population.^{1,2} Unlike collision based sports examined before, (i.e., rugby league^{6,7}), where contact is proposed to contribute to fatigue, football-related NMF is likely driven more by the cumulative effects of repeated explosive efforts and eccentric muscle loading.¹⁰ Therefore, investigating how speed, explosive strength, reactive strength, and endurance capacities relate to NMF may provide new insights into how academy players can better tolerate match demands and mitigate injury risk and performance decrements across the competitive season.

Given the particular injury risk faced by U-18 academy football players and the influence that NMF can exert on exacerbating this relationship,^{1,2} the purpose of this investigation was to identify and examine new hypotheses relating to the moderating effects that speed, strength, and endurance capacities might exert on match-induced NMF in English Premier League (EPL) U-18 academy players. These insights might help coaches and practitioners to prioritise the development of physical qualities that best mitigate the fatiguing effects of match play.

Methods

Match induced neuromuscular fatigue testing

Fourteen elite-level (i.e., tier 5 as per the Participant Classification Framework¹¹) U-18 players (age = 17.0 ± 0.7; height = 1.82 ± 0.07 m; body mass = 73.5 ± 7.6 kg)

from one EPL academy team attended two testing sessions around 8 competitive league home games that were evenly distributed (i.e., 2 games per season quarter) across the 2023–2024 season. Sample size was determined by the available squad size and the number of player – match observations. While no a-priori power analysis was conducted owing to these practical limitations, the research design aimed to enhance statistical sensitivity by employing a repeated measures design across a single homogenous cohort. Consequently, both effect sizes (ES) and confidence intervals are reported in our results to aid interpretation of the findings (given the sample size constraint). Match load was measured using sports global positioning system (GPS) and micro electrical mechanical sensor (MEMS) devices (Statsports APEX, Belfast, Northern Ireland, UK), sampling at 18 Hz (GPS) and 952 Hz (tri-axial accelerometer). All games kicked off at 11:00 AM and testing was conducted at: 1) 09:00 the day before match day (MD), (i.e., MD-1), and 2) 09:00 two-days post-match (MD + 2). These time points were chosen based on our previous findings that match induced changes to the chosen NMF measures peaked at 48 h post-match in a similar cohort.¹⁰ On each test day players completed three CMJ, IADS and IPCS tests according to methods described previously.^{10,12} Variable selection for these tests was based on our previous research examining the reliability¹² and responsiveness¹⁰ of these tests to match play in a similar cohort. Accordingly, chosen test variables were jump height (JH) for the CMJ and peak force (PF) for IADS and IPCS. All data were collected during single-game weeks and no data were analysed from games in which extra time was played. Data from players who played >75 min were included in the analysis (mean ± SD = 85.8 ± 8.7 min). The match duration threshold was based on the combined need to ensure sufficient match exposure to induce NMF whilst maximising the total number of player-match observations. Indeed, our previous work, and other scientific literature indicate that NMF is typically observed when elite level academy football players exceed ~75 min of match play.^{10,13,14} Ethical approval (application number: 2021-21_230) was provided by the **Insert institution name following peer review**, Human Research Ethics Committee and written informed consent and parental consent were obtained from all players and a parent / legal guardian.

Physical capacity testing

Explosive strength (CMJ jump height), reactive strength (30 cm drop jump (DJ)), speed (10 m and 30 m linear speed), maximal strength (IMTP) and endurance (1 km Time Trial (TT)) capacities were measured at 4 time points evenly distributed across one competitive season under replicable conditions: 1) at the beginning of pre-season, 2) at the end of pre-season, 3), half-way through the competitive season and 4) at the end of the competitive

season. At each time point, three test efforts were completed for CMJ, DJ, linear speed and IMTP and one test effort was completed for 1 km TT. Jump, speed and strength measures were conducted in an environmentally controlled performance centre and the 1 km TT was conducted on an outdoor grass pitch at the team's training facility.

Within-participant variability for the physical performance tests was calculated using the coefficient of variation (CV) according to the following equation¹⁵:

$$CV\% = \frac{\text{Standard Deviation}}{\text{Mean}} \times 100$$

Average within-participant CV's for CMJ jump height, DJ RSI, IMTP, 10 m linear speed and 30 m linear speed across the experimental period were 2.85%, 6.58%, 2.37%, 1.33% and 0.85% respectively.

For all test dates, testing started at 10:30 AM the day after a recovery day, five days after the most recent game (i.e., MD-2). All players had routinely performed these tests for at least one full competitive season and were therefore considered to be highly familiar with all protocols. Prior to testing, players performed a standardised 8-min warm-up following a RAMP protocol: ~4-min of low-intensity running and gross-motor warm-up exercises and ~4-min of dynamic mobility and sprint mechanics exercises. Physical capacity test methods are explained in order of their administration below. 5 min of recovery were allocated between each performance test type.

Explosive strength. Explosive strength was assessed using CMJ and bilateral 30 cm drop jump (DJ) tests according to EPL academy testing recommendations. CMJ testing and force plate calibration procedures for all jump measures are described previously.^{10,12} Drop jump testing was performed from a 30 cm box onto dual force plates (ForceDecks FD4000, Vald Performance, Brisbane, AU), sampling at 1000 Hz according to methods described previously.¹⁶ Force-time curves were analysed automatically using proprietary software (ForceDecks Version 2.0.8000, Vald Performance, Brisbane, AU). Players performed three DJs, separated by ~30 s during which they were cued to 'step off of the box using their preferred foot', to 'land with both feet on the force plates at the same time' and to 'minimise ground contact time and jump as high as possible'.¹⁶ Players were required to keep their hands on their hips for the entirety of each jump. All jump testing was conducted by the same experienced practitioner. In cases where a measurement error was observed (i.e., 'jumping' or 'stepping down' from the box, 'tucking' or 'piking' the legs during the flight phase, or if they did not land on the force plates), data were omitted, and the player was asked to perform another repetition. Variables of interest for the CMJ and DJ were JH (flight time) and modified reactive strength index (RSI), respectively.

Modified RSI was calculated using the using the equation¹⁷:

$$\text{Modified RSI} = \frac{\text{JH (m)}}{\text{Contraction Time (s)}}$$

Speed

Speed was assessed using 10 m and 30 m linear speed tests according to EPL academy testing recommendations. Accordingly, testing was conducted indoors, on a FIFA approved artificial pitch, with speed gates (Brower TCi Timing System, Draper, USA) fixed at a height of 0.9 m. Prior to testing players completed three progressive submaximal and one maximal 20 m warm up sprints. After three minutes of recovery, players performed three maximal 30 m test sprints separated by 3-min of recovery. Players used a two-point start with their preferred foot positioned forwards behind a start line. Players were cued to 'sprint maximally through the three speed gates' which were positioned at 1) a zero line (1 m in front of a start line) 2) 10 m in front of the zero line and 3) 30 m in front of the zero line. Players were cued to avoid using a countermovement and self-selected when to initiate each sprint. Variables of interest were the fastest times (s) attained from the three maximal efforts for both the 10 m and 30 m tests.

Strength. Strength was measured using the IMTP according to methods described previously.¹⁸ Players performed three submaximal 4 s IMTP efforts progressing from 60% to 90% of perceived maximal effort followed by three maximal 'test' efforts with one minute of recovery allocated in between. Testing was conducted in a bespoke, fixed isometric strength testing rig (Blk Box Fitness, Newtownabbey, UK), with embedded dual force plates (ForceDecks FD4000, Vald Performance, Brisbane, AU), sampling at 1000 Hz. Acceptable posture was defined as adopting a knee angle between 120–145°, and a hip angle between 140–150° whilst maintaining the trunk in an upright position, confirmed using a goniometer (Physio Parts, Twickenham, UK).¹⁸ Players used an overhand clean grip, with the hands 'fixed' to the bar using pulling straps (Eleiko, Halmstad, Sweden).¹⁸ Prior to testing, players were asked to adopt a rigid torso whilst avoiding pretension or countermovement and were cued to 'push your feet into the ground and pull on the bar as fast and as hard as possible'. Players were given a countdown of '3, 2, 1, pull' and were cued to 'pull, pull, pull, pull' across the 4 s test and then to 'relax'.¹⁸ The best relative peak vertical force (N/kg) attained from the 3 test efforts was calculated automatically from force-time curves using proprietary software (ForceDecks Version 2.0.8000, Vald Performance, Brisbane, AU).

Endurance. Endurance was measured using a 1 km TT according to methods described previously,^{19,20} and as recommended by the English national Football Association.¹⁹ Players

completed ten 100 m shuttle runs between a zero and a 100 m line, which were marked out by cones, and were cued to complete the target distance as quickly as possible. Two test administrators were positioned at each line to ensure that all players completed the correct distance for each shuttle. Total time required to complete the test (s) was recorded.^{19,20}

Data analysis

The match induced change for each NMF test between MD-1 and MD+2 was calculated for each player - match observation. The average change for each NMF test across the experimental period was then calculated for each player. The average match induced change for each NMF test was then associated with the in-season average result for each physical capacity test for each player. This enabled examination of the relationship between match induced change to each NMF test and performance in each physical capacity test.

Statistical analysis

All statistical analysis was conducted in R version 4.3.3 (R Foundation for Statistical Computing, Vienna, Austria). Normality of data were examined using Shapiro-Wilk tests and by visual examination of Q-Q plots. The linear relationship between the mean match induced change for each NMF measure (i.e., CMJ, IPCS and IADS) and the season average physical capacity measure (i.e., CMJ, DJ, 10 m speed, 30 m speed and 1 km TT) was examined using Pearson's *R* (*r*) correlation coefficients if assumptions of linearity and homoscedasticity were met, and by Spearman Rank correlation coefficients (*rho*) if assumptions of normality were violated (i.e., Shapiro-Wilk: $p > 0.05$). Significance was set at $p < 0.05$. The ES of relationships were interpreted as: r or $rho < 0.1 = \text{trivial}$; $0.1-0.3 = \text{small}$; $0.3-0.5 = \text{moderate}$; and $>0.5 = \text{large}$.²¹

Results

Descriptive statistics

Average match duration, total distance, high metabolic load distance and sprint distance across the experimental period were 97.9 min, 10,291 m, 2073 m and 264 m.

Descriptive statistics for the match induced changes to NMF tests and physical capacity tests are shown in Tables 1 and 2, below.

The relationships between match induced changes to CMJ JH, IADS PF, IPCS PF and each physical capacity test are shown in Figures 1–3, below. Change to CMJ JH shared non-significant, *moderate* positive relationships with DJ RSI and IMPT (Figure 1). Change to IADS PF shared non-significant *small* negative relationships with 10 m Speed, 30 m Speed, CMJ JH and DJ RSI (Figure 2). Change to IPCS PF shared non-significant *small* negative

Table 1. Descriptive statistics for match induced NMF test measure changes between MD-1 and MD + 2.

NMF Test	Mean Change (\pm SD)	95% CI
CMJ JH (cm)	0.12 (2.45)	-1.18–1.42
IPCS PF (N)	-21.1 (37.48)	-41.1 to -1.23
IADS PF (N)	-34.6 (63.91)	-68.6 to -0.53

Note: CMJ JH, countermovement jump height; IPCS PF, isometric posterior chain strength peak force; IADS PF, isometric adductor peak force; cm, centimetre and N=newtons.

Table 2. Descriptive statistics for physical capacity test measures.

Physical capacity test	Mean (\pm SD)	95% CI
CMJ JH (cm)	40.8 (4.23)	38.6–43.1
DJ RSI	1.92 (0.44)	1.68–2.14
10 m Speed (s)	1.65 (0.07)	1.61–1.68
30 m Speed (s)	3.99 (0.15)	3.92–4.07
IMTP PF (N/kg)	37.0 (2.86)	35.5–38.5
1 km TT (min)	3.38 (0.15)	3.30–3.45

Note: CMJ JH, countermovement jump height; DJ RSI, drop jump reactive strength index; IMTP PF, isometric mid-thigh pull peak force; km, kilometre; TT, time trial; cm, centimetre; s, second; N/kg, newtons per kilogram body mass and min; minutes.

relationships with 10 m Speed and 1 km TT (Figure 3). All other relationships were neither statistically significant or practically important, (i.e., ES = *trivial*), (Figures 1–3).

Discussion

The purpose of this case report was to identify new hypotheses relating to the moderating effects of speed, strength, and endurance capacities on match-induced neuromuscular fatigue (NMF) in English Premier League (EPL) U-18 academy players. All interactions were not statistically significant; therefore, relationships should be interpreted in conjunction with the ES and 95% CI presented in Figures 1–3. Overall, our most important findings are that maximal (IMTP) ($p=0.06$) and reactive (DJ RSI) ($p=0.16$) strength qualities shared *moderate* positive relationships with match-induced changes to CMJ JH (Figure 1). These observations generate important hypotheses relating to the potential role that maximal and reactive strength might play in mitigating NMF in academy football players and warrant further investigation.

Our finding that maximal strength shared a *moderate* positive relationship with match-induced NMF (Figure 1) is consistent with previous research in young Rugby League^{6,7} and senior professional football²² players. Indeed, Johnson and colleagues reported that greater maximal strength (i.e., 3RM back squat) was associated with higher match loads and reduced match induced muscle damage^{6,7} and NMF (CMJ peak power)⁶ in elite

level young Rugby League players. It was proposed that strength training adaptations (i.e., muscle fibre type distribution, calpain activity and / or force generating capacity²³) provide protective effects against match induced muscle damage and NMF. Importantly, similar effects were also observed by Owen and colleagues,²² who reported that ‘stronger’ (3RM back squat) senior professional male football players exhibited less match-induced muscle damage than ‘weaker’ players at 48 h post-match. Overall, this finding supports a hypothesis that maximal strength might serve to ‘mechanically protect’ young football players against NMF. Current evidence indicates that this might occur by virtue of mitigating the mechanical fatigue consequences of match play.¹³

A potential explanation for this relationship relates to the role that strength adaptations might exert on enhancing fatigue resistance. For example, strength training can induce important changes to muscle fibre composition, potentially favouring a greater proportion of Type IIa fibres, which exhibit a higher oxidative capacity compared to more fatigable Type IIb fibres.²⁴ Consequently, we speculate that this adaptation might serve to improve force-generating capacity and resistance to NMF during match play. Additionally, it is also possible that ‘stronger’ players also exhibit greater tendon stiffness and fascicle length, which might contribute to improve efficiency in force transmission and by virtue of this, reduce eccentric-induced muscle damage, giving rise to NMF.^{25,26}

It is feasible that neural adaptations associated with strength, power and speed training could play a role in mitigating NMF and serve to pretext against NMF. For example, increased motor unit recruitment efficiency and rate coding might enable stronger players to maintain force output with lower relative burden and reduce the extent of NMF accumulated during match-play.²⁷ Moreover, enhanced co-contraction of the stabilising musculature might serve to improve joint stability and by virtue of this, also help to mitigate NMF induced by high-intensity match actions.²⁸ Future studies examining these relationships might examine muscle fibre composition, neuromuscular efficiency, and muscle-tendon unit adaptations to confirm these mechanisms.

Interestingly, our findings also suggest that reactive strength (DJ RSI) might exert a *moderate* beneficial effect on match induced change to CMJ JH ($p=0.16$), (Figure 1). Though no previous research has examined this relationship, conceptually this finding might be explained by training induced improvements to energy conservation (i.e., greater utilisation of elastic- as opposed to metabolic- energy during high-intensity match activities) and the resultant reduction to match induced muscle damage (i.e., owing to the reduced relative mechanical stress of high intensity match activities) expected with improved SSC efficiency.¹⁴ Indeed, improved SSC efficiency might enable greater utilisation of stored elastic




Figure 1. Linear relationships between match induced change to CMJ JH and physical capacity test performances: CMJ JH (cm), DJ RSI (m/s), 10 m speed (s), 30 m speed (s), IMTP PF (N) and 1 km TT (min). Pearson's R correlations are presented as r value, effect size, t value and p value. Spearman's Rank correlations are presented as ρ value, effect size, S value and p value. The grey area depicts the upper and lower 95% confidence intervals (CI).




Figure 2. Linear relationships between match induced change to IADS PF and physical capacity test performances: CMJ JH (cm), DJ RSI (m/s), 10 m speed (s), 30 m speed (s), IMTP PF (N) and 1 km TT (min). Pearson's R correlations are presented as r value, effect size, t value and p value. Spearman's Rank correlations are presented as ρ value, effect size, S value and p value. The grey area depicts the upper and lower 95% confidence intervals (CI).




Figure 3. Linear relationships between match induced change to IPCS PF and physical capacity test performances: CMJ JH (cm), DJ RSI (m/s), 10 m Speed (s), 30 m Speed (s), IMTP PF (N) and 1 km TT (min). Pearson's R correlations are presented as r value, effect size, t value and p value. Spearman's Rank correlations are presented as ρ value, effect size, S value and p value. The grey area depicts the upper and lower 95% confidence intervals (CI).

energy in the muscle-tendon unit and reduce reliance on metabolically demanding contractile work.²⁵ Additionally, improved SSC function is associated with increased tendon stiffness, which in-turn, might enhance the ability to store and release elastic energy.²⁹ Conceptually, these mechanisms might go some way to explain this relationship, by improving mechanical efficiency and reducing the relative mechanical stress imposed on the lower limb musculature during high-intensity match actions, helping to mitigate NMF.

A further explanation for this relationship might relate to the beneficial role that improved SSC function might exert on muscle fibre strain, by altering the muscle-tendon interaction characteristics. For example, with greater tendon compliance, a more substantial proportion of muscle and tendon length change might occur in the tendon as opposed to muscle fibres, which might serve to reduce the degree of match-induced eccentric muscle damage.³⁰ Indeed, this mechanism might be particularly relevant during high-force eccentric match actions that likely contribute most substantially to NMF.³¹ Additionally, enhanced motor unit synchronisation and pre-activation strategies, might also facilitate improved force transmission and joint stabilisation during explosive actions,³² reducing the need for compensatory muscle activity and excessive co-contraction, which could otherwise accelerate NMF.

The extent to which reactive strength qualities might help to mitigate NMF in *specific* musculature remains an important question. We speculate that enhancing SSC function might serve to optimise force attenuation and energy return during dynamic movements, and on this basis suggest that it is possible that reactive strength might exert more profound protective effects on muscle groups that are more dependent on elastic energy storage (i.e., the quadriceps and calf-Achilles complex), as opposed to those that are likely to be more susceptible to match-induced eccentric damage, (i.e., the hamstrings). Of note, the quadricep musculature plays an important role in force attenuation during landing and deceleration tasks in football.²⁹ Therefore, greater SSC efficiency, characterised by enhanced tendon stiffness and neuromuscular pre-activation, might help to reduce fibre strain and metabolic cost during repeated high-intensity movements.²⁵ Indeed, this might go some way to explain why players with superior reactive strength could exhibit less match-induced CMJ impairment. Overall, the relationship shared between DJ RSI and match induced changes to CMJ JH generates a further hypothesis that reactive strength qualities might also serve to mitigate match induced NMF in EPL U-18 academy football players. Future research examining the interplay between SSC efficiency, match loads, and neuromuscular performance are required to confirm these hypotheses.

Compared to the anterior chain musculature, the posterior chain and adductor muscles experience particularly high eccentric loading during sprinting and change of direction tasks during match play. Given their role in deceleration and force transfer, these muscle groups might be particularly prone to muscle damage and (consequently), NMF.²⁸ Since the protective benefits of reactive strength are primarily linked to elastic energy return rather than eccentric force attenuation, it is possible that fatigue in the posterior chain and adductor musculature is less influenced by reactive strength and more dependent on maximal and eccentric strength characteristics. Indeed, this may help explain our finding that 10 m ($p=0.29$) and 30 m ($p=0.41$) linear speed shared small negative relationships with match-induced changes in IADS PF (Figure 2) and IPCS PF (10 m; $p=0.45$) (Figure 3), suggesting that players with greater maximal sprint capacities exhibited greater match-induced NMF. It is feasible that faster players might rely more heavily on hamstring-driven sprint mechanics, which could make them more susceptible to fatigue in muscle groups with a high proportion of Type II fibres. This relationship may also reflect other innate and training-induced differences between faster and slower players. For example, faster players are likely to possess a greater distribution of Type IIa and Type IIb muscle fibres, which, while advantageous for high-force and high-velocity match actions, might be more susceptible to muscle damage and fatigue^{22,28}. Consequently, it is possible that match-induced NMF is more pronounced in readily fatigable muscle groups associated with faster-, compared to slower- players. However, caution should be exercised when interpreting these exploratory findings.

Limitations

Consistent with similar hypothesis generating case reports,^{6,7,22} our results are underpowered and based on a limited sample size and number of player-match observations. This relates to the well-established challenges of conducting research in elite-level professional football environments owing to factors including player injury, illness, suspension, and team selection, described previously.²² Consequently, the results herein should be considered with caution and as hypothesis generating only. Indeed, we stress the need for further research drawing upon larger data sets to both confirm the hypotheses proposed and examine their underpinning mechanisms. We acknowledge the presence of individual data points across Figures 1–3 that may appear as outliers in our analyses. Given our limited sample size, these data points likely influenced the observed relationships. While we did not exclude individual data points due to the exploratory nature of this study, future research with larger datasets should investigate whether these findings are consistent when accounting for potential outliers and individual variability. We also

acknowledge that other potentially important confounding variables that might relate to the development of NMF (i.e., variability of prior training load, recovery strategies and the effect of playing position) were not considered in our analysis and stress that these limitations should be considered when interpreting our hypothesis-generating results.

Conclusion

Our exploratory findings generate important hypotheses relating to the beneficial moderating roles that maximal strength, reactive strength, and endurance capacities might serve in protecting young football players against match-induced NMF. Additionally, we observed a *small* positive relationship between 1 km TT performance and match-induced IPCS changes and that greater maximal sprint speeds were associated with increased perturbations to adductor and posterior chain performance, which might be explained by greater fatiguability in faster players. This finding suggests that ‘fast’ players might benefit from improving aspects of endurance capacity and by virtue of this, fatigue resistance, a hypothesis that warrants further exploration. For example, it is possible that enhanced anaerobic capacity might allow faster players to sustain higher concentric power outputs for longer periods of match play, which could mitigate reliance on inefficient eccentric braking strategies that might otherwise contribute to the development of muscle damage and NMF. Evidently, establishing how endurance characteristics might interact with the interplay between speed and NMF in football is important. Consequently, future research should examine the balance between speed and endurance development in fast players to maximize performance and minimise NMF decrements. Finally, the diversity of relationships observed in the individual data points (Figures 1–3) confirms the necessity for individual player monitoring in practice and the potential for CMJ JH and DJ RSI (in particular) to identify players who might be at risk of developing NMF.

The data that support the findings of this study are available from the corresponding author upon reasonable request.



Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

This research was supported by a UEFA Research Grant Programme (UEFA RGP) award.

ORCID iDs

Matthew Springham  <https://orcid.org/0000-0002-0206-3676>
Jamie Salter  <https://orcid.org/0000-0002-7375-1476>

References

1. Bittencourt NFN, Meeuwisse WH, Mendonca LD, et al. Complex systems approach for sports injuries: moving from risk factor identification to injury pattern recognition-narrative review and new concept. *Br J Sports Med* 2016; 50: 1309–1314.
2. Materne O, Chamari K, Farooq A, et al. Injury incidence and burden in a youth elite football academy: a four-season prospective study of 551 players aged from under 9 to under 19 years. *Br J Sports Med* 2021; 55: 493–500.
3. Schwiete C, Roth C, Skutschik C, et al. Effects of muscle fatigue on exercise-induced hamstring muscle damage: a three-armed randomized controlled trial. *Eur J Appl Physiol* 2023; 123: 2545–2561.
4. Meeusen R, Duclos M, Foster C, et al. Prevention, diagnosis, and treatment of the overtraining syndrome: joint consensus statement of the European college of sport science and the American college of sports medicine. *Med Sci Sports Exerc* 2013; 45: 186–205.
5. Soligard T, Schwellnus M, Alonso JM, et al. How much is too much? (part 1) international Olympic committee consensus statement on load in sport and risk of injury. *Br J Sports Med* 2016; 50: 1030–1041.
6. Johnston RD, Gabbett TJ and Jenkins DG. Influence of playing standard and physical fitness on activity profiles and post-match fatigue during intensified junior rugby league competition. *Sports Med Open* 2015; 1: 18.
7. Johnston RD, Gabbett TJ, Jenkins DG, et al. Influence of physical qualities on post-match fatigue in rugby league players. *J Sci Med Sport* 2015; 18: 209–213.
8. Constantine E, Taberner M, Richter C, et al. Isometric posterior chain peak force recovery response following match-play in elite youth soccer players: Associations with relative posterior chain strength. *Sports (Basel)* 2019; 7: 218–230.
9. Stolen T, Chamari K, Castagna C, et al. Physiology of soccer: an update. *Sports Med* 2005; 35: 501–536.
10. Springham M, Singh N, Stewart P, et al. Acute neuromuscular and perceptual responses to U-18 English Premier League academy football match play. *Eur J Sport Sci* 2024; 24: 1405–1413.
11. McKay AKA, Stellingwerff T, Smith ES, et al. Defining training and performance caliber: a participant classification framework. *Int J Sports Physiol Perform* 2022; 17: 317–331.
12. Springham M, Singh N, Stewart P, et al. Countermovement jump and isometric strength test-retest reliability in English premier league academy football players. *ACCEPTED IN: Int J Sports Physiol Perform* 2024; 12: 1444–1454.
13. Harper DJ, Carling C and Kiely J. High-intensity acceleration and deceleration demands in elite team sports competitive match play: a systematic review and meta-analysis of observational studies. *Sports Med* 2019; 49: 1923–1947.
14. Nicol C, Avela J and Komi PV. The stretch-shortening cycle : a model to study naturally occurring neuromuscular fatigue. *Sports Med* 2006; 36: 977–999.
15. Hopkins WG. Measures of reliability in sports medicine and science. *Sports Med* 2000; 30: 1–15.
16. Maloney SJ, Fletcher IM and Richards J. A comparison of methods to determine bilateral asymmetries in vertical leg stiffness. *J Sports Sci* 2016; 34: 829–835.
17. Bishop C, Turner A, Jordan M, et al. A framework to guide practitioners for selecting metrics during the countermovement and drop jump tests. *Strength Cond J* 2022; 44: 95–103.
18. Comfort P, Dos’Santos T, Beckham GK, et al. Standardization and methodological considerations for the isometric midhigh pull. *Strength Condition J* 2019; 41: 57–79.
19. Rosenblatt B, Evans M and Cain E. *Physical Profiling Technical Handbook*. London, UK: The Football Association, 2019.
20. Kelly VG and Wood A. The correlation between the 30-15 intermittent fitness test and a novel test of running performance. *J Austr Strength Condition* 2013; 21: 91–93.
21. Hopkins WG, Marshall SW, Batterham AM, et al. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc* 2009; 41: 3–13.
22. Owen A, Dunlop G, Rouissi M, et al. The relationship between lower-limb strength and match-related muscle damage in elite level professional European soccer players. *J Sports Sci* 2015; 33: 2100–2105.
23. Raastad T, Owe SG, Paulsen G, et al. Changes in calpain activity, muscle structure, and function after eccentric exercise. *Med Sci Sports Exerc* 2010; 42: 86–95.
24. Wilson JM, Loenneke JP, Jo E, et al. The effects of endurance, strength, and power training on muscle fiber type shifting. *J Strength Cond Res* 2012; 26: 1724–1729.
25. Kubo K, Kanehisa H and Fukunaga T. Effects of resistance and stretching training programmes on the viscoelastic properties of human tendon structures in vivo. *J Physiol* 2002; 538: 219–226.
26. Seynnes OR, Erskine RM, Maganaris CN, et al. Training-induced changes in structural and mechanical properties of the patellar tendon are related to muscle hypertrophy but not to strength gains. *J Appl Physiol (1985)* 2009; 107: 523–530.
27. Aagaard P, Simonsen EB, Andersen JL, et al. Neural adaptation to resistance training: changes in evoked V-wave and H-reflex responses. *J Appl Physiol (1985)* 2002; 92: 2309–2318.
28. Higashihara A, Nagano Y, Ono T, et al. Differences in activation properties of the hamstring muscles during overground sprinting. *Gait Posture* 2015; 42: 360–364.
29. Bojsen-Moller J, Magnusson SP, Rasmussen LR, et al. Muscle performance during maximal isometric and dynamic contractions is influenced by the stiffness of the tendinous structures. *J Appl Physiol (1985)* 2005; 99: 986–994.
30. Ishikawa M and Komi PV. Muscle fascicle and tendon behavior during human locomotion revisited. *Exerc Sport Sci Rev* 2008; 36: 193–199.
31. Komi PV. Stretch-shortening cycle: a powerful model to study normal and fatigued muscle. *J Biomech* 2000; 33: 1197–1206.
32. Taube W, Leukel C and Gollhofer A. How neurons make us jump: the neural control of stretch-shortening cycle movements. *Exerc Sport Sci Rev* 2012; 40: 106–115.