

Est.
1841

YORK
ST JOHN
UNIVERSITY

Madigan, Daniel J. ORCID logoORCID:
<https://orcid.org/0000-0002-9937-1818>, Stoeber, Joachim and Passfield, Louis (2017) Perfectionism and training distress in junior athletes: a longitudinal investigation. *Journal of Sports Sciences*, 35 (5). pp. 470-475.

Downloaded from: <https://ray.yorks.ac.uk/id/eprint/1721/>

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version:

<http://dx.doi.org/10.1080/02640414.2016.1172726>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at ray@yorks.ac.uk

Madigan, D. J., Stoeber, J., & Passfield, L. (in press). Perfectionism and training distress in junior athletes: A longitudinal investigation. *Journal of Sports Sciences*.

Perfectionism and Training Distress in Junior Athletes:

A Longitudinal Investigation

Daniel J. Madigan^a

Joachim Stoeber^b

Louis Passfield^a

Author Note

^aSchool of Sport & Exercise Sciences, University of Kent, Chatham Maritime, Kent ME4 4AG, United Kingdom.

^bSchool of Psychology, University of Kent, Canterbury, Kent CT2 7NP, United Kingdom.
Correspondence concerning this article should be addressed to Daniel Madigan, School of Sport & Exercise Sciences, University of Kent, Chatham Maritime, Kent ME4 4AG, United Kingdom.
Phone: +44-1634-888903; e-mail: dm412@kent.ac.uk

Abstract

1 Perfectionistic athletes may train harder and for longer than non-perfectionistic athletes, leaving
2 them susceptible to elevated levels of training distress. So far, however, no study has
3 investigated the relationships between perfectionism and training distress, a key indicator of
4 overtraining syndrome. Furthermore, no study has determined psychological predictors of
5 overtraining syndrome. Using a two-wave design, the present study examined perfectionistic
6 strivings, perfectionistic concerns, and training distress in 141 junior athletes (mean age 17.3
7 years, range 16-19 years) over 3 months of active training. Multiple regression analyses were
8 employed to test cross-sectional and longitudinal relationships between perfectionism and
9 training distress. In all analyses, perfectionism emerged as a significant predictor, but strivings
10 and concerns showed differential relationships. When the cross-sectional relationships were
11 regarded, perfectionistic concerns positively predicted training distress ($p < .01$), whereas
12 perfectionistic strivings negatively predicted training distress ($p < .001$). When the longitudinal
13 relationships were regarded, only perfectionistic concerns predicted increases in training distress
14 ($p < .05$), whereas perfectionistic strivings did not ($p > .05$). The findings suggest that sports
15 scientists who wish to identify athletes at risk of overtraining syndrome may monitor athletes'
16 perfectionistic concerns as a possible risk factor.

17 *Keywords:* perfectionistic strivings, perfectionistic concerns, training distress,
18 overtraining, junior athletes, longitudinal study

19

1 Perfectionism and Training Distress in Junior Athletes:

2 A Longitudinal Investigation

3 To succeed in competitive sports, athletes are required to participate in intensive training
4 regimes. However, excessive training accompanied by inadequate rest can result in overtraining
5 syndrome (Meeusen et al., 2013). Sports scientists have investigated ways to monitor athletes'
6 training responses with the aim of identifying at-risk athletes and intervening to prevent
7 overtraining syndrome. One psychological marker of overtraining syndrome that sport scientists
8 have identified is training distress (Kenttä, Hassmén, & Raglin, 2001; Meeusen et al., 2013;
9 Raglin & Morgan, 1994) which focuses on training-related mood disturbance. Consequently,
10 researchers have sought to determine factors that may predispose athletes to training distress.
11 One such factor may be perfectionism, as perfectionistic athletes may train harder and for longer
12 than non-perfectionistic athletes (Flett & Hewitt, 2014). In support of this suggestion, case
13 studies have shown that athletes who overtrained were characterised by exhibiting a high level
14 of perfectionism (Gould, Tuffey, Udry, & Loehr, 1997; Krane, Greenleaf, & Snow, 1997). In
15 addition, there is evidence that perfectionism is related to associated syndromes such as athlete
16 burnout and compulsive exercise (Hill & Curran, in press; Hill, Robson, & Stamp, 2015;
17 Madigan, Stoeber, & Passfield, 2015). So far, however, the relationships between perfectionism
18 in athletes and training distress have not been investigated. Furthermore, previous research has
19 yet to identify any psychological predictors of training distress. Therefore, the aim of the present
20 study was to provide a first investigation of perfectionism and training distress in junior athletes.

21 **Perfectionism**

22 Perfectionism is a personality disposition characterised by striving for flawlessness and
23 setting exceedingly high standards of performance accompanied by tendencies for overly critical

1 evaluations of one's behaviour (Flett & Hewitt, 2002). However, perfectionism has various
2 aspects, and there are different dimensions of perfectionism with different characteristics.
3 Consequently, perfectionism is best conceptualized as a multidimensional disposition (Frost,
4 Marten, Lahart, & Rosenblate, 1990; Hewitt & Flett, 1991; see Enns & Cox, 2002, for a review).
5 Factor analyses comparing various measures of multidimensional perfectionism have provided
6 support for two higher-order dimensions: *perfectionistic strivings* capturing perfectionist
7 personal standards and a self-oriented striving for perfection and *perfectionistic concerns*
8 capturing concerns about making mistakes, feelings of discrepancy between one's standards and
9 performance, and fears of negative evaluation and rejection by others if one fails to be perfect
10 (see Stoeber & Otto, 2006, for a review).

11 Differentiating between perfectionistic strivings and perfectionistic concerns is important
12 when investigating perfectionism in sports because the two dimensions show different, and often
13 opposite, patterns of relationships with psychological processes and outcomes. Perfectionistic
14 concerns are consistently associated with negative processes and outcomes (e.g., maladaptive
15 coping, negative affect), whereas perfectionistic strivings are often associated with positive
16 processes and outcomes (e.g., adaptive coping, positive affect) or inversely with negative
17 processes and outcomes. The latter is particularly evident when the overlap between
18 perfectionistic strivings and concerns is controlled for (Gotwals, Stoeber, Dunn, & Stoll, 2012;
19 Stoeber, 2011). In this case "pure perfectionistic strivings" are identified (i.e., as perfectionistic
20 strivings with the negative influence of perfectionistic concerns partialled out; Hill & Curran, in
21 press). Pure perfectionistic strivings are usually more adaptive than perfectionistic strivings
22 because they lack those aspects common to both perfectionistic strivings and concerns (e.g., self-
23 criticism, conditional self-acceptance; Hill, 2014).

1 **Training Distress**

2 Excessive training accompanied by inadequate recovery (and possible non-training
3 stressors) can result in an overtraining syndrome which is characterised by a sport-specific
4 decrease in performance that can persist for weeks and sometimes months (Meeusen et al.,
5 2013). Whereas there is no single diagnostic tool to identify athletes suffering from overtraining
6 syndrome, monitoring training responses allows for early identification of at-risk athletes and
7 may give practitioners a chance to reduce the negative consequences of excessive training. There
8 are numerous indicators of training responses associated with overtraining syndrome including
9 biochemical, physiological, immunological, and psychological indicators which all have
10 limitations (Meeusen et al., 2013). A recent systematic review, however, suggests that
11 psychological indicators capturing athletes' subjective responses to training can help identify
12 athletes at risk of overtraining syndrome and do so more effectively than physiological
13 indicators (Saw, Main, & Gastin, in press).

14 In particular, measures of training distress have shown promise in capturing athletes'
15 subjective responses to training (Meeusen et al., 2013). One such measure is the Training
16 Distress Scale (TDS; Raglin & Morgan, 1994). The TDS is a widely used mood-based measure
17 of training distress and is derived from the Profile of Mood States (POMS; McNair, Lorr, &
18 Droppleman, 1971), which has itself been found to be an effective tool in assessing training
19 stress and overtraining syndrome risk. Moreover, depression is one of the more serious outcomes
20 of overtraining syndrome and training distress is derived largely from POMS depression items
21 (Armstrong & VanHeest, 2002). Although the POMS has shown a dose-response relationship
22 with training load (Raglin & Wilson, 2000), the TDS has shown to be more accurate in
23 identifying overtrained athletes (Kenttä et al., 2001; Raglin & Morgan, 1984). Still, despite

1 attempts (e.g., Raglin & Wilson, 2000), no study has identified psychological factors that
2 longitudinally predispose athletes to greater risk of developing an overtraining syndrome, and
3 this information would provide a very useful (and currently missing) diagnostic tool for
4 preventing the development of the overtraining syndrome in athletes.

5 **The Present Study**

6 Against this background, the aim of the present study was to provide a first investigation
7 of the relationships between perfectionism and training distress in athletes examining cross-
8 sectional and longitudinal relationships between perfectionistic strivings, perfectionistic
9 concerns, and training distress. Based on previous theory and empirical evidence from cross-
10 sectional studies on perfectionism and compulsive exercise (e.g., Hill et al., 2015), we
11 hypothesised that perfectionism would predict training distress. In this, however, we expected
12 only perfectionistic concerns to be a positive predictor, whereas we expected perfectionistic
13 strivings to be either a negative predictor or to show nonsignificant relationships. Based on
14 previous theory and empirical evidence from a longitudinal study on perfectionism and burnout
15 (Madigan et al., 2015), we further expected perfectionism to predict longitudinal changes in
16 training distress, but expected only perfectionistic concerns to be a positive predictor.

17 **Method**

18 **Participants**

19 A sample of 141 junior athletes (125 male, 16 female) was recruited at two sports
20 academies (92 from one academy, 49 from the other) to participate in the present study. Sports
21 academies are part of the United Kingdom's further education system. Their main purpose is to
22 recruit and develop promising junior athletes by providing them with a professional coaching
23 environment while they study alongside their sporting commitments. Academy athletes are

1 selected based on their ability (competitive performance in trials to enter the academy) and
2 regularly compete at a regional, national, or international level. Participants' mean age was 17.3
3 years ($SD = 0.8$; range = 16-19 years). Participants were involved in a range of sports (60 in
4 soccer, 36 in rugby, 18 in basketball, 14 in athletics, and 13 in other sports [e.g., cycling,
5 squash]) and trained on average 9.6 hours per week ($SD = 5.6$).

6 **Procedure**

7 The study was approved by the university's ethics committee. Informed consent was
8 obtained from all participants. In addition, parental consent was obtained from participants
9 below the age of 18 (as per the ethics committee's recommendation). Questionnaires were
10 distributed during training in the presence of the first author, or athletes completed an online
11 version of the questionnaire. Participants were administered all measures twice separated by
12 three months, once in October (Time 1) and then again in January (Time 2). During this period,
13 all participants were in regular seasonal training and competition with the exception of those
14 involved in athletics who were in pre-seasonal training. Furthermore, a three-month period has
15 been found sufficient in longitudinal research on perfectionism and athlete burnout (Madigan et
16 al., 2015).

17 **Measures**

18 **Perfectionism.** To measure perfectionism, we followed a multi-measure approach
19 (Stoeber & Madigan, in press) and used four subscales from two multidimensional measures of
20 perfectionism in sport: the Sport Multidimensional Perfectionism Scale (Dunn et al., 2006) and
21 the Multidimensional Inventory of Perfectionism in Sport (Stoeber, Otto, Pescheck, Becker, &
22 Stoll, 2007). To measure perfectionistic strivings, we used two indicators: the 7-item Sport
23 Multidimensional Perfectionism Scale subscale capturing personal standards (e.g. "I have

1 extremely high goals for myself in my sport”; $M = 3.35$, $SD = 0.71$) and the 5-item
2 Multidimensional Inventory of Perfectionism in Sport subscale capturing striving for perfection
3 (“I strive to be as perfect as possible”; $M = 3.21$, $SD = 0.79$), and then standardised the scale
4 scores before combining them to measure perfectionistic strivings (cf. Dunkley, Zuroff, &
5 Blankstein, 2003). To measure perfectionistic concerns, we also used two indicators, the 8-item
6 Sport Multidimensional Perfectionism Scale subscale capturing concerns over mistakes (“People
7 will probably think less of me if I make mistakes in competition”; $M = 2.89$, $SD = 0.77$) and the
8 5-item Multidimensional Inventory of Perfectionism in Sport subscale capturing negative
9 reactions to imperfection (“I feel extremely stressed if everything does not go perfectly”; $M =$
10 2.89 , $SD = 0.83$), and again standardised the scale scores before combining them to measure
11 perfectionistic concerns. The four subscales have demonstrated reliability and validity in
12 previous studies (e.g., Madigan, Stoeber, & Passfield, in press; Stoeber, Stoll, Salmi, & Tiikkaja,
13 2009). Moreover, both are reliable and valid indicators of perfectionistic strivings and
14 perfectionistic concerns (e.g., Gotwals et al., 2012; Stoeber & Madigan, in press). Participants
15 were asked to indicate to what degree each statement characterised their attitudes in their sport
16 responding on a scale from 1 (*strongly disagree*) to 5 (*strongly agree*).

17 **Training Distress.** To measure training distress, we used the Training Distress Scale
18 (TDS; Raglin & Morgan, 1994). The TDS is comprised of ten items, seven items capturing
19 training distress (e.g., “worthless”, “miserable”, “bad tempered”) and three filler items (e.g.,
20 “helpful”) which are ignored when calculating TDS scores. The TDS has demonstrated
21 reliability and validity in numerous studies (e.g., Kenttä et al., 2001; Raglin & Morgan, 1994).
22 Participants were asked to indicate how often within the last week (“During training last week, I
23 felt...”) they had been feeling as described in each item responding on a scale from 1 (*not been*

1 *feeling this way*) to 5 (*been feeling extremely like this*).

2 **Data Screening**

3 First, we inspected the data for missing values. Because very few item responses were
4 missing ($i = 12$), missing responses were replaced with the mean of the item responses of the
5 corresponding scale (ipsatised item replacement; Graham, Cumsille, & Elek-Fisk, 2003). Next
6 we computed Cronbach's alphas for our variables which were all satisfactory (see Table 1).
7 Following recommendations by Tabachnick and Fidell (2007), data were screened for
8 multivariate outliers. One participant showed a Mahalanobis distance larger than the critical
9 value of $\chi^2(4) = 18.47, p < .001$, and was excluded from further analyses. Finally, we conducted
10 two Box's M tests to examine if the variance-covariance matrices showed any differences
11 between academies or gender. Both tests were nonsignificant with $F_s < 1.14, p_s > .21$ despite the
12 test being so sensitive to minor differences that the recommended significance level for this test
13 is $p < .001$ (Tabachnick & Fidell, 2007). Therefore, all further analyses were collapsed across
14 academies and gender. Because 35 participants did not complete the measures on both
15 occasions, the final cross-sectional sample size was $N = 140$ (124 male, 16 female) and the final
16 longitudinal sample size was $N = 106$ (90 male, 16 female).

17 **Results**

18 **Bivariate Correlations**

19 Next, we inspected the bivariate correlations between all variables (see Table 1). As in
20 previous research (e.g., Madigan et al., in press), the dimensions of perfectionism showed a
21 significant positive correlation with each other. Furthermore, training distress at Time 1 showed
22 a significant positive correlation with training distress at Time 2. Perfectionistic concerns
23 showed significant positive correlations with training distress at Time 1 and Time 2, whereas

1 perfectionistic strivings showed no significant correlations with training distress at either time
2 point.

3 **Multiple Regression Analyses**

4 We then conducted two multiple regression analyses (see Table 2). The first regression
5 analysis investigated the cross-sectional relationships between perfectionism and training
6 distress to examine the unique relationships of the two perfectionism dimensions by controlling
7 for their overlap. For this, we entered perfectionistic strivings and perfectionistic concerns
8 simultaneously into the regression. Results showed that the model explained 11% of the variance
9 in training distress ($R^2 = .114, p < .001$) and the perfectionism dimensions showed opposite
10 relationships with training distress: Perfectionistic concerns positively predicted training distress
11 ($\beta = .39, p < .001$), whereas perfectionistic strivings negatively predicted training distress ($\beta = -$
12 $.28, p < .01$).

13 The second regression analysis investigated the longitudinal relationship between
14 perfectionism and training distress. First, we controlled for baseline levels of training distress by
15 entering training distress at Time 1 in Step 1. We then entered the two perfectionism dimensions
16 simultaneously in Step 2. Results showed that perfectionistic concerns predicted residual
17 increases in training distress over time, whereas perfectionistic strivings emerged as a
18 nonsignificant predictor.

19 **Discussion**

20 The aim of the present study was to investigate the relationships between perfectionism in
21 athletes and training distress differentiating perfectionistic strivings and perfectionistic concerns.
22 Providing a first investigation of both cross-sectional and longitudinal relationships, we found
23 perfectionism to be significantly related to training distress, but the two dimensions of

1 perfectionism showed different relationships with training distress. When the cross-sectional
2 relationships were regarded, perfectionistic concerns positively predicted training distress,
3 whereas perfectionistic strivings negatively predicted training distress. When the longitudinal
4 relationships were regarded, only perfectionistic concerns positively predicted residual increases
5 in training distress, whereas perfectionistic strivings was as a nonsignificant predictor.

6 This is the first study to show that perfectionism is related to training distress in athletes.
7 The finding that perfectionistic concerns in athletes show positive cross-sectional and
8 longitudinal relationships with training distress is in agreement with case studies indicating that
9 athletes who overtrain are characterised by high levels of perfectionism (Gould et al., 1997;
10 Krane et al., 1997). They are also in agreement with findings from research on perfectionism and
11 compulsive exercise, showing perfectionistic athletes to have higher levels of compulsion to
12 exercise (Hill et al., 2015). More importantly, the present findings suggest that perfectionism
13 may be a factor contributing to the development of training distress in athletes. As training
14 distress is a psychological marker of overtraining syndrome, perfectionistic athletes may be
15 susceptible to the negative consequences of this syndrome. However, only perfectionistic
16 concerns appear to be a risk factor, not perfectionistic strivings.

17 Differently from perfectionistic concerns, perfectionistic strivings showed a negative
18 cross-sectional relationship with training distress. This dovetails with previous research on
19 perfectionism in sport suggesting that the strivings dimension of perfectionism often shows
20 positive relationships with processes and outcomes that can be considered adaptive or, as in the
21 present study, negative relationships with processes and outcomes that can be considered
22 maladaptive (for details, see Gotwals et al., 2012; Stoeber, 2011). Note, however, that
23 perfectionistic strivings showed a negative relationship with training distress only in the cross-

1 sectional analyses, but not in the longitudinal analyses. This finding suggests that perfectionistic
2 strivings may not have a protective effect for athletes in regard to training distress, and it
3 highlights the importance of using longitudinal designs when investigating the relationships of
4 perfectionism in sport. Note, however, that differences in findings could be explained by the
5 larger sample size and consequently improved statistical power for our cross-sectional analyses
6 enabling us to detect smaller effects. Furthermore, we note that the negative cross-sectional
7 relationship with training distress only emerged after controlling for the overlap with
8 perfectionistic concerns, suggesting that the relationship only holds for “pure perfectionistic
9 strivings,” that is, perfectionistic strivings with the negative influence of perfectionistic concerns
10 partialled out (Hill & Curran, in press).

11 Previously, no study has identified any psychological predictors of overtraining syndrome.
12 What may explain why perfectionistic concerns are such a predictor? One explanation may be
13 differences in training load. Athletes high in perfectionistic concerns may have trained more
14 excessively than athletes low in perfectionistic concerns leading to increased training distress. If
15 this suggestion is correct, this effect of perfectionistic concerns could be countered through
16 targeted monitoring and better management of training load by the coach and/or support staff
17 (Meeusen et al., 2013). Another explanation may be that athletes high in perfectionistic concerns
18 experienced more non-training stressors than athletes low in perfectionistic concerns. The sport
19 environment can be highly stressful for athletes, and athletes differ in how they cope with stress.
20 Research has shown that perfectionistic concerns are associated with maladaptive coping in
21 sports (Hill, Hall, & Appleton, 2010). Consequently, athletes high in perfectionistic concerns
22 may have coped less well with the stress associated with high training demands and experienced
23 greater training distress. An effect of perfectionistic concerns contributing to training distress

1 could be attenuated by helping athletes to better cope with stress (cf. Meeusen et al., 2013; see
2 also Antony & Swinson, 2009).

3 **Limitations and Future Research**

4 The present study had a number of limitations. First, our study focused on a sample
5 comprised exclusively of junior athletes therefore the generalizability of our findings may be
6 limited. However, previous research has shown that junior athletes experience lifetime rates of
7 overtraining syndrome equivalent to adult non-elite athletes. Furthermore, previous research has
8 shown that experiencing overtraining syndrome at a young age may predispose athletes to an
9 increased lifetime risk of developing overtraining syndrome (Raglin, Sawamura, Alexiou,
10 Hassmén, & Kenttä, 2000; Meeusen et al., 2013). Second, the study did not include any
11 mediators, that is, variables that may explain why perfectionistic concerns predicted increases in
12 training distress. Future longitudinal studies on perfectionism may therefore consider designs
13 that include mediators such as training load and coping in addition to perfectionism and training
14 distress (cf. Cole & Maxwell, 2003). Third, we may have found larger effects if we had
15 investigated only those sports that involve high levels of physical conditioning as the risk of
16 overtraining syndrome may be higher in these sports (Kenttä et al., 2001). Finally, the study only
17 examined training distress. Whereas training distress is a key indicator of overtraining
18 syndrome, future research would benefit from including further indicators (cf. Meeusen et al.,
19 2013) to explore whether the relationships we found between perfectionism and training distress
20 replicate with a wider range of indicators for overtraining syndrome.

21 **Conclusion**

22 The present study makes an important contribution to our understanding of the
23 relationships between perfectionism in sport and training distress, being the first to identify both

1 cross-sectional and longitudinal relationships in a large sample of athletes. Even though the
2 effects we found were only small- to medium-sized (Cohen, 1992) and perfectionism explained
3 only a modest percentage of variance in training distress, the present study is the first to identify
4 a psychological predictor of increased training distress. Moreover, even small-sized effects
5 matter as they may accumulate over time (Prentice & Miller, 1992). Consequently, sports
6 scientists monitoring athletes' training responses to identify athletes at risk of overtraining
7 syndrome may want to monitor athletes' perfectionistic concerns as a factor predisposing
8 athletes to experience higher levels of training distress that may further increase over time
9 putting athletes at risk of developing overtraining symptoms.

10

References

- 1
2 Antony, M. M., & Swinson, R. P. (2009). *When perfect isn't good enough: Strategies for coping*
3 *with perfectionism* (2nd ed.). Oakland, CA: New Harbinger.
- 4 Armstrong, L. E., & VanHeest, J. L. (2002). The unknown mechanism of the overtraining
5 syndrome: Clues from depression and psychoneuroimmunology. *Sports Medicine*, 32, 185-
6 209.
- 7 Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112, 155-159.
- 8 Cole, D. A., & Maxwell, S. E. (2003). Testing mediational models with longitudinal data:
9 Questions and tips in the use of structural equation modeling. *Journal of Abnormal*
10 *Psychology*, 112, 558-577.
- 11 Dunkley, D. M., Zuroff, D. C., & Blankstein, K. R. (2003). Self-critical perfectionism and daily
12 affect: Dispositional and situational influences on stress and coping. *Journal of Personality*
13 *and Social Psychology*, 84, 234-252.
- 14 Dunn, J. G. H., Causgrove Dunn, J., Gotwals, J. K., Vallance, J. K. H., Craft, J. M., & Syrotuik,
15 D. G. (2006). Establishing construct validity evidence for the Sport Multidimensional
16 Perfectionism Scale. *Psychology of Sport and Exercise*, 7, 57-79.
- 17 Enns, M. W., & Cox, B. J. (2002). The nature and assessment of perfectionism: A critical
18 analysis. In G. L. Flett & P. L. Hewitt (Eds.), *Perfectionism* (pp. 33-62). Washington, DC:
19 American Psychological Association.
- 20 Flett, G. L., & Hewitt, P. L. (2002). Perfectionism and maladjustment: An overview of
21 theoretical, definitional, and treatment issues. In P. L. Hewitt & G. L. Flett (Eds.),
22 *Perfectionism* (pp. 5-31). Washington, DC: American Psychological Association.
- 23 Flett, G. L., & Hewitt, P. L. (2014). "The perils of perfectionism in sports" revisited: Toward a

- 1 broader understanding of the pressure to be perfect and its impact on athletes and dancers.
2 *International Journal of Sport Psychology*, 45, 395-407.
- 3 Frost, R. O., Marten, P., Lahart, C., & Rosenblate, R. (1990). The dimensions of perfectionism.
4 *Cognitive Therapy and Research*, 14, 449-468.
- 5 Gotwals, J. K., Stoeber, J., Dunn, J. G. H., & Stoll, O. (2012). Are perfectionistic strivings in
6 sport adaptive? A systematic review of confirmatory, contradictory, and mixed evidence.
7 *Canadian Psychology*, 53, 263-279.
- 8 Gould, D., Tuffey, S., Udry, E., & Loehr, J. (1997). Burnout in competitive junior tennis players:
9 III. Individual differences in the burnout experience. *The Sport Psychologist*, 11, 257-276.
- 10 Graham, J. W., Cumsille, P. E., & Elek-Fisk, E. (2003). Methods for handling missing data. In J.
11 A. Schinka & W. F. Velicer (Eds.), *Handbook of psychology: Research methods in*
12 *psychology* (Vol. 2, pp. 87-114). New York: Wiley.
- 13 Hewitt, P. L., & Flett, G. L. (1991). Perfectionism in the self and social contexts:
14 Conceptualization, assessment, and association with psychopathology. *Journal of*
15 *Personality and Social Psychology*, 60, 456-470.
- 16 Hill, A. P. (2014). Perfectionistic strivings and the perils of partialling. *International Journal of*
17 *Sport and Exercise Psychology*, 12, 302-315.
- 18 Hill, A. P., & Curran, T. (in press). Multidimensional perfectionism and burnout A meta-
19 analysis. *Personality and Social Psychology Review*.
- 20 Hill, A. P., Hall, H. K., & Appleton, P. R. (2010). Perfectionism and athlete burnout in junior
21 elite athletes: The mediating influence of coping tendencies. *Anxiety, Stress, & Coping*, 23,
22 415-430.
- 23 Hill, A. P., Robson, S. J., & Stamp, G. M. (2015). The predictive ability of perfectionistic traits

- 1 and self-presentational styles in relation to exercise dependence. *Personality and Individual*
2 *Differences, 86*, 176-183.
- 3 Kenttä, G., Hassmén, P., & Raglin, J. S. (2001). Training practices and overtraining syndrome in
4 Swedish age-group athletes. *International journal of sports medicine, 22*, 460-465.
- 5 Krane, V., Greenleaf, C. A., & Snow, J. (1997). Reaching for gold and the price of glory: A
6 motivational case study of an elite gymnast. *The Sport Psychologist, 11*, 53-71.
- 7 Madigan, D. J., Stoeber, J., & Passfield, L. (2015). Perfectionism and burnout in junior athletes:
8 A three-month longitudinal study. *Journal of Sport & Exercise Psychology, 37*, 305-315.
- 9 Madigan, D. J., Stoeber, J., & Passfield, L. (in press). Perfectionism and attitudes towards doping
10 in junior athletes. *Journal of Sports Sciences*.
- 11 McNair, D. M., Lorr, M., & Droppleman, L. F. (1971). *Manual for the Profile of Mood States*.
12 San Diego, CA: Educational and Industrial Testing Services.
- 13 Meeusen, R., Duclos, M., Foster, C., Fry, A., Gleeson, M., Nieman, D., ... & Urhausen, A.
14 (2013). Prevention, diagnosis and treatment of the overtraining syndrome: Joint consensus
15 statement of the European College of Sport Science (ECSS) and the American College of
16 Sports Medicine (ACSM). *European Journal of Sport Science, 13*, 1-24.
- 17 Prentice, D. A., & Miller, D. T. (1992). When small effects are impressive. *Psychological*
18 *Bulletin, 112*, 160-164.
- 19 Raglin, J. S., & Morgan, W. P. (1994). Development of a scale for use in monitoring training-
20 induced distress in athletes. *International Journal of Sports Medicine, 15*, 84-88.
- 21 Raglin, J. S., Sawamura, S., Alexiou, S., Hassmén, P., & Kenttä, G. (2000). Training practices
22 and staleness in 13-18 year old swimmers: A cross-cultural study. *Pediatric Sports*
23 *Medicine, 12*, 61-70.

- 1 Raglin, J. S., & Wilson, G. S. (2000). Overtraining in athletes. In Y. L. Hanin (Ed.), *Emotions in*
2 *sport* (pp. 191-207). Champaign, IL: Human Kinetics.
- 3 Saw, A. E., Main, L. C., & Gustin, P. B. (in press). Monitoring the athlete training response:
4 Subjective self-reported measures trump commonly used objective measures: A systematic
5 review. *British Journal of Sports Medicine*.
- 6 Stoeber, J. (2011). The dual nature of perfectionism in sports: Relationships with emotion,
7 motivation, and performance. *International Review of Sport and Exercise Psychology*, 4,
8 128-145.
- 9 Stoeber, J., & Madigan, D. J. (in press). Measuring perfectionism in sport, dance, and exercise:
10 Review, critique, recommendations. In A. P. Hill (Ed.), *The psychology of perfectionism in*
11 *sport, dance, and exercise*. London: Routledge.
- 12 Stoeber, J., & Otto, K. (2006). Positive conceptions of perfectionism: Approaches, evidence,
13 challenges. *Personality and Social Psychology Review*, 10, 295-319.
- 14 Stoeber, J., Otto, K., Pescheck, E., Becker, C., & Stoll, O. (2007). Perfectionism and competitive
15 anxiety in athletes: Differentiating striving for perfection and negative reactions to
16 imperfection. *Personality and Individual Differences*, 42, 959-969.
- 17 Stoeber, J., Stoll, O., Salmi, O., & Tiikkaja, J. (2009). Perfectionism and achievement goals in
18 young Finnish ice-hockey players aspiring to make the Under-16 national team. *Journal of*
19 *Sports Sciences*, 27, 85-94.
- 20 Tabachnick, B.G., & Fidell, L.S. (2007). *Using multivariate statistics* (5th ed.). Boston, MA:
21 Pearson.

Table 1

Descriptive Statistics, Cronbach's Alphas, and Bivariate Correlations

Variable	1	2	3	4
1. Perfectionistic strivings				
2. Perfectionistic concerns	.54***			
3. Training distress Time 1	-.07	.24**		
4. Training distress Time 2	.09	.33**	.56***	
<i>M</i>	0.00	0.00	1.89	2.07
<i>SD</i>	0.91	0.93	0.92	0.71
Cronbach's alpha	.79	.85	.90	.77

Note. $N = 140$ for Time 1. $N = 106$ for Time 2. Time 2 = 3 months after Time 1.

** $p < .01$. *** $p < .001$.

Table 2

Multiple Regression Predicting Longitudinal Changes in Training Distress

	Training distress Time 2	
	ΔR^2	β
Step 1: Training distress Time 1	.319***	
Training distress Time 1		.56***
Step 2: Perfectionism	.046*	
Training distress Time 1		.51***
Perfectionistic strivings		-.02
Perfectionistic concerns		.23*

Note. $N = 106$. Time 2 = 3 months after Time 1.

* $p < .05$. *** $p < .001$.