

Est.  
1841

YORK  
ST JOHN  
UNIVERSITY

Brislane, Aine ORCID logoORCID:

<https://orcid.org/0000-0002-3214-6544>, Jones, Helen, Low, David, Holder, Sophie and Hopkins, Nicola, D. (2018) Regular moderate intensity aerobic exercise does not affect vascular outcomes during pregnancy: A pilot study. In: Institute for Health Research Conference, 1st May 2018, Liverpool, United Kingdom. (Unpublished)


Downloaded from: <https://ray.yorks.ac.uk/id/eprint/3748/>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

# RaY

Research at the University of York St John

For more information please contact RaY at [ray@yorks.ac.uk](mailto:ray@yorks.ac.uk)



INSTITUTE FOR HEALTH RESEARCH  
 CONFERENCE 1<sup>ST</sup> MAY 2018  
 Royal Court, Liverpool

Register for a free ticket at <https://goo.gl/UPZftS>

**Abstract Submission Form for Poster presentation**

Complete all text within the clear boxes provided in 12pt Arial.

Name of person submitting the abstract:					
First	Aine	Last	Brislane	Title	Ms
Affiliation(s) Liverpool John Moores University					
Names of additional author(s) and their affiliations	1. Jones, H. Liverpool John Moores University				
	2. Low, D. Liverpool John Moores University				
	3. Holder, S. Liverpool John Moores University				
	4. Hopkins, N.D. Liverpool John Moores University				
Title	Regular moderate intensity aerobic exercise does not affect vascular outcomes during pregnancy: A pilot study				
<b>Abstract.</b> (Maximum of 250 words)					
<p><b>Introduction:</b> Cardiovascular diseases (CVD) are the leading cause of morbidity and mortality during pregnancy attributed to progressive vascular impairments. Regular exercise has been shown to improve vascular outcomes. The aim of this pilot study was to determine the impact of a 6-month exercise intervention on vascular outcomes in previously inactive pregnant women.</p> <p><b>Methods:</b> Ten healthy pregnant women were recruited to a control (CONT; N=6 33±0.5 years, BMI 22.4kg/m<sup>2</sup>) or exercise group (EX; N=4 31.5± 0.6 years, BMI 23.8±2.1kg/m<sup>2</sup>) at 13-15 weeks gestation. Ultrasound was used to assess brachial and femoral flow mediated dilation (bFMD; fFMD respectively) at the end of trimester 1(T1), 2(T2) and 3(T3). Aerobic capacity was estimated using the Astrand submaximal cycling protocol. Physical activity (PA) was measured over 7 days using accelerometry. The exercise intervention consisted of 3x15 minute weekly exercise sessions in trimester-2 (T2), progressing to 4x30 minutes in trimester-3 (T3). Data were analyzed for main effects of group and time using repeated measures ANOVA.</p>					

**Results:** There were no time, group or interaction effects for BFMD (T1: 9.2±2.1%, T2:6.7±0.9%, T3: 9±5%, P=0.76), FFMD (T1, 6.5±3.4%, T2, 6.2±2%, T3, 2.4±1.9%, P=0.18). No differences in aerobic capacity were evident (T1, 47.4±5ml/kg, T2, 43.4±3.3ml/kg, T3, 39.1±3.5ml/kg, p=0.22). No differences were observed for physical activity (T1, 330±87mins/d; T2, 296±52mins/d; T3, 271±16mins/d).

**Conclusion:** Our findings suggest that vascular outcomes are not impacted upon exercising during trimesters 2 and 3 of pregnancy. It is plausible that the hormonally induced hemodynamic adaptations which occur during pregnancy cannot be overridden by moderate intensity exercise.

**References**  
(maximum of 3)

**IHR Interest Group** Cardiovascular Health and Care

The deadline for submitting abstracts is **5pm Friday 23<sup>rd</sup> March 2018**.

Please submit completed forms to [S.L.Schofield@ljmu.ac.uk](mailto:S.L.Schofield@ljmu.ac.uk)