

Est.
1841

YORK
ST JOHN
UNIVERSITY

Lundkvist, Erik, Gustafsson, Henrik, Gerber, Markus, Lundqvist, Carolina, Ivarsson, Andreas and Madigan, Daniel J. ORCID: <https://orcid.org/0000-0002-9937-1818> (2019) Commentary: Early risk detection of burnout: Development of the Burnout Prevention Questionnaire for coaches. *Frontiers in Psychology*.

Downloaded from: <http://ray.yorks.ac.uk/id/eprint/4173/>

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version: <https://www.frontiersin.org/articles/10.3389/fpsyg.2019.02721/full>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at ray@yorks.ac.uk

Lundkvist, E., Gustafsson, H., Gerber, M., Lundqvist, C., Ivarsson, A., & Madigan, D. J. (in press). Commentary: Early risk detection of burnout: Development of the Burnout Prevention Questionnaire for coaches. *Frontiers in Psychology*.

Commentary: Early Risk Detection of Burnout: Development of the Burnout Prevention Questionnaire for Coaches

Erik Lundkvist^{1*}, Henrik Gustafsson^{2,3}, Markus Gerber⁴, Carolina Lundqvist⁵, Andreas Ivarsson⁶, Daniel J. Madigan⁷

Author note

¹Performance and Training Unit, The Swedish School of Sport and Health Sciences, Stockholm, Sweden

²Faculty of Arts and Social Sciences, Karlstad University, Karlstad, Sweden

³Norwegian School of Sport Sciences, Oslo, Norway

⁴SportScience Section, Department of Sport, Exercise and Health, University of Basel, Basel, Switzerland

⁵Division of Psychology, Department of Behavioural Sciences and Learning, Linköping University, Linköping, Sweden

⁶Centre of Research on Welfare, Health and Sport, Halmstad University, Halmstad, Sweden

⁷School of Sport, York St John University, York, United Kingdom

* Corresponding Author: erik.lundkvist@gih.se

1 Introduction

In a recent volume of this journal, Schaffran and colleagues (2019) introduced the Burnout Prevention Questionnaire for Coaches (BPQ-C). Although we recognize the worthwhile efforts of Schaffran et al., we believe that there are several issues associated with this instrument. This commentary aims to expand on why we think the BPQ-C should not currently be used by practitioners and researchers to screen for burnout.

2.1 Lack of theoretical foundations

With regards to theory, we have four concerns. First, Schaffran et al.'s definition (and subsequent measurement) of burnout is inadequate. Although several definitions of burnout exist, most researchers agree that burnout is a multidimensional construct and models share – at a minimum – an exhaustion component (Maslach, Jackson, & Leiter, 1996; Raedeke & Smith, 2001; Shirom, 2005). This was recently highlighted by the World Health Organization (2018). While not everyone agrees with one exact definition, no well-recognized definition includes general stress, social stress, and amotivation (Maslach et al., 1996; World Health Organization, 2018; Raedeke & Smith, 2001).

Second, Schaffran et al. (2019) seem to conflate antecedents, outcomes, and the construct of burnout itself. This point is exemplified by the fact that chronic job stress has historically been defined as an essential burnout antecedent and not a defining characteristic of burnout (Maslach et al., 1996; Shirom, 2005). Therefore, using stress as a symptom of burnout and not as an antecedent is conceptually incorrect. Similarly, the dimension labeled “pre-burnout” includes fatigue. This is confusing because of the evident conceptual overlap with exhaustion, which is known as a defining characteristic of burnout (Grossi, Perski, Osika, & Savic, 2015).

Third, an underlying assumption for the BPQ-C seems to be that there exist sequential events in the burnout process. It is therefore peculiar that the authors relied exclusively on cross-sectional data when they created and tested their instrument (Hendrix, Acevedo, & Hebert, 2000; Leiter & Maslach, 1999, 2003). Especially since the complexity regarding temporality between burnout dimensions has been discussed before both in sport and more generally (Martinent, Louvet, & Decret, 2016; Shirom & Melamed, 2006).

Our final conceptual concern is the reliance on statistical considerations (from analysis of one sample), instead of also considering theoretical aspects, when deciding on the inclusion/exclusion of certain BPQ-C dimensions (Study 2). This exploratory approach has been criticized in the burnout literature previously and heightens risks for conceptual overlap with other constructs (see Shirom & Melamed, 2006), as well as misunderstanding of temporal relations with emotional exhaustion (Moore, 2000). The problem with this procedure becomes particularly apparent in relation to the resources dimension which defines sleep as a resource that may buffer potential burnout. However, with the same logic, poor sleep quality could be seen as a part of “pre-burnout” since poor sleep quality is a well-known antecedent of burnout symptoms (Söderström, Jeding, Ekstedt, Perski, & Åkerstedt, 2012). Thus, although the authors briefly acknowledge the complexity of sleep in relation to burnout in their discussion, they do not provide any guidelines based on sound empirical work and theory to explain how sleep scores in the BPQ-C should be interpreted.

2.2 Utility as a screening tool

The authors claim that the BPQ-C has the potential to be used as a screening tool in practical settings to detect early signs of burnout, and to guide prevention strategies before a coach develops more

serious burnout symptoms. Although a commendable goal, one well-known problem is that there is often no clear distinction between normal functioning and psychological illness. Evaluations of symptoms should be related to the context in which they appear as well as to contemporary values and economic forces present in society and politics (Kinderman, Allsopp, & Cooke, 2017). Furthermore, Schaffran and colleagues do not provide evidence that the BPQ-C can be used to adequately screen for clinical burnout. To do so, analyses such as receiver operating characteristic analysis (ROC), which tests the ability of measures to discriminate between individuals with and without a certain clinical characteristic, is necessary (Kraemer, 1992). Comparisons between a clinical and a healthy sample were not a part of the validation of the BPQ-C. In fact, no descriptive statistics of burnout are reported. This makes it even harder to determine variation and representability of the samples that were used. In addition, even when a clinical sample is compared with a healthy sample, discriminating between healthy and ill individuals is challenging as they show substantial overlap (Lundgren-Nilsson, Jonsdottir, Pallant, & Ahlborg, 2012). Therefore, using screening tools without clinical interviews creates substantial risks of both false positives and negatives, a matter that has been neglected in the sport psychology literature and is not considered by Schaffran et al. in their article.

2.3 Questionnaire development

Our last point of criticism concerns the development and validation of the BPQ-C. Several guides on how to develop a psychological instrument are available. Although they differ, there are some common grounds when it comes to item development, scale development, and scale evaluation (Boateng, Neilands, Frongillo, Melgar-Quinonez, & Young, 2018; MacKenzie, Podsakoff, & Podsakoff, 2011). Schaffran et al., however, only communicate a few of these suggested steps. When such a step is communicated, like confirmatory factor analysis (CFA), the high degree of similarity in subscale content suggests a redundancy of these factors. This issue is compounded by the fact that parallel analysis suggested a one-factor model (see Sample 1). A stepwise evaluation would have easily eliminated this in the item developing phase. Lastly, in the three-factor CFA, several residuals are correlated (see Figure 1). The reasons for this needs to be communicated since correlating residuals comes with several risks (with capitalization of chance being the most prominent one), making replication in another sample unlikely (Landis, Edwards, & Cortina, 2009).

3 Discussion

This commentary provides several arguments as to why, in our opinion, the BPQ-C fails to meet several theoretical, clinical, and psychometric standards. Substantial work is therefore needed before the BPQ-C can be considered a valid measure. Instead, we recommend the use of existing validated measures of burnout. One example of such a measure is the Shirom Melamed Burnout Questionnaire (SMBQ; Kushnir & Melamed, 1992), which is based on Hobfoll's Conservation of Resources (COR) theory (Hobfoll, 1989). The SMBQ has been validated in both clinical and healthy samples (Lundgren-Nilsson, Jonsdottir, Pallant, & Ahlborg, 2012). Another alternative would be the Maslach Burnout Inventory (Maslach, Jackson, & Leiter, 1996), which follows the theoretical underpinnings provided by Maslach and colleagues (Maslach, Jackson, & Leiter, 1996). The MBI has shown reasonable psychometric properties in previous research, and for which at least some evidence exists that the instrument is able to discriminate between clinical and healthy individuals. However, the latter only applies to the exhaustion subscale (Kleijweg, Verbraak, & Van Dijk, 2013). Here, we note, however, that using self-report measures as the only source for clinical screening may not be successful because of the overlap between clinical and healthy individuals. We therefore recommend

that, wherever possible, self-report screening should be combined with clinical interviews (Kleijweg, Verbraak, & Van Dijk, 2013; Lundgren-Nilsson, Jonsdottir, Pallant, & Ahlborg, 2012) and that the interpretation of self-reported burnout measures with clinical cut-offs should be made with caution.

4 References

- Boateng, G. O., Neilands, T. B., Frongillo, E. A., Melgar-Quiñonez, H. R., & Young, S. L. (2018). Best Practices for Developing and Validating Scales for Health, Social, and Behavioral Research: A Primer. *Frontiers in Public Health*, 6(149). doi:10.3389/fpubh.2018.00149
- Grossi, G., Perski, A., Osika, W., & Savic, I. (2015). Stress-related exhaustion disorder—clinical manifestation of burnout? A review of assessment methods, sleep impairments, cognitive disturbances, and neuro-biological and physiological changes in clinical burnout. *Scandinavian Journal of Psychology*, 56(6), 626-636.
- Hendrix, A. E., Acevedo, E. O., & Hebert, E. (2000). An examination of stress and burnout in certified athletic trainers at Division IA universities. *Journal of athletic training*, 35(2), 139.
- Hobfoll, S. E. (1989). *Conservation of resources: A new attempt at conceptualizing stress*. *American Psychologist*, 44(3), 513.
- Kinderman, P., Allsopp, K., & Cooke, A. (2017). Responses to the publication of the American Psychiatric Association's DSM-5. *Journal of Humanistic Psychology*, 57(6), 625-649. doi:10.1177/0022167817698262
- Kleijweg, J. H., Verbraak, M. J., & Van Dijk, M. K. (2013). *The clinical utility of the Maslach Burnout Inventory in a clinical population*. *Psychological Assessment*, 25(2), 435.
- Kraemer, H. C. (1992). *Evaluating medical tests: Objective and quantitative guidelines* (Vol. 26): Sage publications Newbury Park, CA:.
- Kushnir, T., & Melamed, S. (1992). *The Gulf War and its impact on burnout and well-being of working civilians*. *Psychological Medicine*, 22(4), 987-995.
- Landis, R., Edwards, B., & Cortina, J. (2009). Correlated residuals among items in the estimation of measurement models. *Statistical and methodological myths and urban legends: Doctrine, verity, and fable in the organizational and social sciences*, 195-214.
- Leiter, M. P., & Maslach, C. (1999). Six areas of worklife: a model of the organizational context of burnout. *Journal of health and Human Services administration*, 472-489.
- Leiter, M. P., & Maslach, C. (2003). Areas of worklife: A structured approach to organizational predictors of job burnout *Emotional and physiological processes and positive intervention strategies* (pp. 91-134): Emerald Group Publishing Limited.
- Lundgren-Nilsson, Å., Jonsdottir, I. H., Pallant, J., & Ahlberg, G. (2012). Internal construct validity of the Shirom-Melamed Burnout questionnaire (SMBQ). *BMC Public Health*, 12(1), 1.
- MacKenzie, S. B., Podsakoff, P. M., & Podsakoff, N. P. (2011). Construct measurement and validation procedures in MIS and behavioral research: Integrating new and existing techniques. *MIS quarterly*, 35(2), 293-334.
- Martinent, G., Louvet, B., & Decret, J.-C. (2016). Longitudinal trajectories of athlete burnout among young table tennis players: A 3-wave study. *Journal of Sport and Health Science*.
- Maslach, C., Jackson, S. E., & Leiter, M. P. (1996). *Maslach burnout inventory manual* (Vol. 4): Consulting psychologists press Palo Alto, CA.
- Moore, J. E. (2000). Why is this happening? A causal attribution approach to work exhaustion consequences. *Academy of Management Review*, 25(2), 335-349.
- World Health Organization. (2018). *ICD-11 International classification of diseases for mortality and morbidity statistics (11th Revision)*. Retrieved from <https://icd.who.int/browse11/l-m/en>
- Raedeke, T. D., & Smith, A. L. (2001). Development and preliminary validation of an athlete burnout measure. *Journal of sport and exercise psychology*, 23(4), 281-306.
- Schaffran, P., Kleinert, J., Altfeld, S., Zepp, C., Kallus, K. W., & Kellmann, M. (2019). Early Risk Detection of Burnout: Development of the Burnout Prevention Questionnaire for Coaches. *Frontiers in Psychology*, 10. doi:10.3389/fpsyg.2019.00714
- Shirom, A. (2005). Reflections on the study of burnout. *Work & Stress*, 19(3), 263-270.

- Shirom, A., & Melamed, S. (2006). A comparison of the construct validity of two burnout measures in two groups of professionals. *International journal of stress management*, 13(2), 176.
- Söderström, M., Jeding, K., Ekstedt, M., Perski, A., & Åkerstedt, T. (2012). Insufficient sleep predicts clinical burnout. *Journal of occupational health psychology*, 17(2), 175.