

Est.
1841

YORK
ST JOHN
UNIVERSITY

Salter, Jamie ORCID:

<https://orcid.org/0000-0002-7375-1476>, De Ste Croix, Mark De Ste Croix and Hughes, Jonathan (2020) The moderating impact of maturation on acute neuromuscular and psycho-physiological responses to simulated soccer activity in academy soccer players. European Journal of Sport Science.

Downloaded from: <http://ray.yorks.ac.uk/id/eprint/4806/>

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version:

<https://www.tandfonline.com/doi/full/10.1080/17461391.2020.1851775>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at ray@yorks.ac.uk

1 **The moderating impact of maturation on acute neuromuscular and**
2 **psycho-physiological responses to simulated soccer activity in academy**
3 **soccer players**

4
5 Jamie Salter^{1,2}, Mark B.A. De Ste Croix² and Jonathan D. Hughes²

6
7 ¹*School of Sport, York St John University, York, UK*

8 ²*School of Sport and Exercise, University of Gloucestershire, Gloucester, UK*

9
10 Submission Type: *Original Investigation*

11 Running Head: *Moderating impact of maturation on acute responses*

12 Corresponding Author:

13 Jamie Salter

14 School of Sport

15 York St John University

16 Lord Mayors Walk

17 York, YO31 7EX,

18 United Kingdom

19 j.salter@yorks.ac.uk

20 @jay_salter

21 ORCID: 0000-0002-7375-147

22
23 Abstract Word Count: 248

24
25 Main Text Word Count: 3997

26
27 Tables: 3

28
29 Figures: 1

30

31

32

33

34

35

36

37

38

39

40 **The moderating impact of maturation on acute neuromuscular and**
41 **psycho-physiological responses to simulated soccer activity in academy**
42 **soccer players**
43

44 **Abstract**

45 Resource constraints complicate load monitoring practices in some academies, which is problematic based on load-injury
46 associations surrounding periods of rapid non-linear growth. Limited research has explored relationships between maturation
47 and perceived psycho-physiological response to activity and associated neuromuscular performance changes. This study
48 aimed to quantify neuromuscular and psycho-physiological responses to standardised activity and analyse whether dose-
49 responses were moderated by maturation. Fifty-seven male soccer players (age: 14.1 ± 0.9 years; stature: 165 ± 10 cm; body
50 mass, 57 ± 9 kg; percentage of predicted adult height $92.7 \pm 5\%$) from two Elite Player Performance Plan (EPPP) academies
51 completed the youth soccer-specific aerobic fitness test (Y-SAFT⁶⁰). Countermovement jump (CMJ), reactive strength index
52 (RSI), absolute (ABS) and relative leg stiffness (REL) were measured pre-post the Y-SAFT⁶⁰ with playerload (PL), heart
53 rate (HR), total distance (TD^{ist}) and differential ratings of perceived exertion (dRPE) used as markers of load and intensity.
54 A moderation model was employed to analyse interactions of maturation as a continuous variable. Analysis indicated no
55 significant interaction ($p < 0.05$) between maturation and neuromuscular performance but RPE-Technical demonstrated
56 significant interactions ($p = 0.01$). Slope analysis indicated four variables (PL, RSI, ABS and REL) that demonstrated
57 significance at various stages of maturation, most notably aligning with peak height velocity (~87-96% PAH). Tentatively,
58 we propose that maturational developments in the neuromuscular system offer some mechanistic explanation to the varied
59 dose-responses observed. It is therefore important that maturation is habitually considered within prescription of training
60 programmes and that further empirical studies are completed to determine maturity specific dose-responses.

61
62 **Key words:** *Adolescence, injury & prevention, neuromuscular, team sports, training load*
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79

80 **Introduction**

81 Technological advances have proliferated research around dose-response relationships between load, fatigue and training
82 prescription to incorporate various internal and external training load metrics (i.e. total distance covered, high-speed running
83 and player load). As a result, quantifying physical demands has become commonplace in elite adult populations, with dose-
84 response markers used to influence athletic outcomes (Hader et al., 2019). Recently soccer academies indicated that resource
85 and logistical constraints sometimes prevented implementation of best-practice monitoring strategies (Salter et al., 2020).
86 This is problematic for practitioners working with adolescent populations as the temporal, dynamic and non-linear changes
87 in biological development (e.g. hormonal, neural, bone and muscle) likely complicate dose-responses (Bergeron et al., 2015).
88 Barriers to accurate quantification of individual loads make effective manipulation of the training prescription complex and
89 likely inappropriate, particularly for early or late maturing players (approx. 16-40% of academy populations) (Hill et al.,
90 2020; A. Johnson et al., 2017).

91
92 Injuries are multifactorial, but evidence suggests that inappropriate loading may compound already relatively high
93 incidences during adolescent soccer (D. Johnson et al., 2020; Rommers et al., 2020). Associations between load and injury
94 are noteworthy since the introduction of the Elite Player Performance Plan (EPPP) presented a substantial linear,
95 chronological-age correlated increase in coaching hours from approximately 3,760 to 8,500 (accrued annually between the
96 ages of 8-21) (Premier League, 2011). Subsequent trends suggests increased growth and overuse related injuries particularly
97 in U13-U14 age groups, with Osgood-Schlatter's and Severs Disease widespread (Read et al., 2018). Additionally, audit
98 methods likely underestimate incidences of overuse injuries as they typically adopt a 'time-loss' definition, when often
99 players do not miss training but need modification (Whalan et al., 2020). Between 46-72% of EPPP injuries are non-contact
100 and 30-43% are moderate in nature with approximately 50% injuries occurring during training sessions (Read et al., 2018).
101 It is largely considered that non-contact injuries are preventable whereas traumatic contact injuries are unavoidable (Read
102 et al., 2016). Therefore, better appreciation of dose-responses within these age groups may better mitigate risk and
103 individualise training based on the physical capabilities of individuals.

104
105 Rating of Perceived Exertion (RPE) has been extensively applied within soccer and suggested to have acceptable re-call bias
106 and interchangeability between validated scales (CR10 and CR100) (Fanchini et al., 2017; Wright et al., 2020). This
107 versatility and intuitive application proposed by Foster (1998) (RPE multiplied by session duration) make this a robust and
108 accessible method to document training volume and intensity and subsequently inform training prescription. However,
109 despite being common within academy settings, only one empirical study has explored maturity specific responses to
110 subjective psycho-physiological perceptions of intensity (De Ste Croix et al., 2019) and only three studies have observed the
111 age related training response using RPE derived methods in adolescent team sports (Brink et al., 2010; Wright et al., 2020;
112 Wrigley et al., 2012). Instead, most maturity related research has focussed on performance markers such as speed, endurance
113 or match performance (Beyer et al., 2020; Buchheit & Mendez-Villanueva, 2014). These studies offer mechanistic
114 explanations regarding changes in performance during the major growth period but offer limited insight as to how individuals
115 perceive dose-controlled activity during this period.

116
117 Academy squads are comprised of players within chronological parameters but often present significant variations in
118 physical characteristics including body mass (~50%), stature (~29cm), percentages of predicted adult height (10-15%) and
119 fat free mass (3-8.6kg) (Figueiredo et al., 2010; Hannon et al., 2020; van der Sluis et al., 2015). These maturation specific

120 components likely influence performance (Buchheit & Mendez-Villanueva, 2014), but limited studies using standardised
121 activity profiles have directly observed this influence (Lehnert et al., 2018). Relationships between maturation related dose-
122 responses have also been performed observing neuromuscular performance around competitive fixtures and/or training (De
123 Ste Croix et al., 2019; Oliver et al., 2015; Read et al., 2016). Radnor et al. (2018) suggests that stretch-shortening cycle
124 (SSC) performance increases with age across various neuromuscular tasks such as sprinting, hopping and jumping, therefore
125 it is intuitive to assume maturation influences acute responses in neuromuscular performance. Ambiguously however,
126 findings indicate both significant and non-significant relationships with neuromuscular performance, such as
127 countermovement jump (CMJ), reactive strength index (RSI) and leg stiffness (De Ste Croix et al., 2015, 2019; Lehnert et
128 al., 2017). Most recently De Ste Croix and colleagues (2019) concluded that maturation did not influence responses to match-
129 play in youth soccer despite large changes in RSI, a finding reinforced by Lehnert et al. (2017). Interestingly, despite sharing
130 many of the same SSC characteristics as RSI, leg stiffness response is less predictable, whereby some individuals
131 experienced acute reductions in leg stiffness with others a state of potentiation with improved stiffness scores post-exercise
132 (De Ste Croix et al., 2019; Oliver et al., 2015). Yet mechanistic theories suggest an inhibited stretch-reflex and reduced rate-
133 of-force development and golgi-tendon organ (GTO) involvement surrounding peak height velocity (PHV), which likely
134 heightens injury threat and reduces mechanical efficiency (Lehnert et al., 2017).

135
136 The relative ambiguity and apparent paucity of evidence observing the influence of maturation on neuromuscular
137 performance and psycho-physiological response provides a rationale for further work in this area. These player monitoring
138 methods (i.e. RPE, CMJ and RSI) are commonly used within academy settings to routinely assess development and can
139 therefore provide meaningful data to practitioners regarding maturity related dose-responses. By better understanding this
140 relationship throughout maturation, we can develop approaches to training prescription and recovery. This may, in turn
141 reduce the possible detrimental impact of loading that is associated with non-contact injury within this population (Oliver et
142 al., 2015). Therefore, the aim of this study was to quantify the neuromuscular performance and psycho-physiological
143 responses to a simulated soccer-specific activity profile and analyse whether this dose-response was moderated by
144 maturation in EPPP academy players.

145 146 **Materials and Methods**

147 *Participants*

148 Fifty-seven male soccer players (age: 14.1 ± 0.9 years; stature: 165 ± 10 cm; body mass, 57 ± 9 kg; predicted adult height
149 $92.7 \pm 5\%$) from Youth Development Phase (YDP) age groups at two EPPP academies took part in this study. Participants
150 were included if they were part of the YDP (U13-U16 age groups), free from injury and available for all scheduled training
151 sessions and matches for the two-weeks prior to data collection. Participants usually complete three training sessions per
152 week and compete in matches at weekends for approximately forty weeks a year. Participants completed written informed
153 consent and parental assent in line with the declaration of Helsinki following ethical approval from the University ethics
154 committee.

155 156 *Procedures*

157 Maturity status was expressed as a percentage of predicted adult height (PAH%) determined by measurement of somatic
158 markers (standing stature, body mass) alongside self-reported parent stature corrected for overestimation (Epstein et al.,
159 1995; Khamis & Roche, 1994). This common approach to estimating maturity status in EPPP academies appears to more

160 accurately predict timing of peak height velocity (PHV) than maturity offset methods (Parr et al., 2020). It is common within
161 maturity research to dichotomise the variable into pre- or post-PHV, or even trichotomise into pre-, circa- and post-PHV for
162 analysis (Radnor et al., 2020; van der Sluis et al., 2015). Although it is possible to do this using the PAH% approach (PHV
163 occurs at approximately between 88-93% PAH) (Cumming et al., 2017), categorisation of continuous variables limits
164 sensitivity (by consuming variability) of data dramatically reducing statistical power, particularly in smaller sample studies
165 (Altman & Royston, 2006). For example, two individuals at a similar percentage of adult height that fall either side of the
166 cut-off threshold for pre-post PHV will be reported as very different opposed to very similar. Therefore, maturity status
167 (PAH%) was regressed as a continuous variable to protect true variations in the outcomes and maintain statistical power
168 (Altman & Royston, 2006).

169
170 To simulate soccer match-play an adapted soccer-specific aerobic field test (SAFT⁹⁰) was used, termed Youth-SAFT (Barrett
171 et al., 2013). This 15-minute fixed-intensity activity profile is based on global-positioning satellite (GPS) locomotion data
172 from male academy competition and includes acceleration, deceleration, lateral shuffling, jogging, back-peddalling and
173 sprinting to replicate demands of match-play. The intensity and locomotor activity is controlled by audio cueing which
174 adopts the multidirectional model of the original SAFT⁹⁰ (Small et al. 2009). In groups of four to minimise variation in
175 neuromuscular sampling time, the 15-minute activity profile was repeated a total of four times (Y-SAFT⁶⁰), interspersed
176 with a 10-minute half-time interval to approximately replicate adolescent competition duration between U13 and U16
177 (Premier League, 2011). All Y-SAFT⁶⁰ simulations were completed on 3G pitch whilst participants wore their normal
178 training attire in order to reduce the variance associated with different surfaces and maintain ecological validity.

179
180 To quantify external load (total distance [T^{Dist}] and playerload [PL]) each player wore a microtechnology Global Positioning
181 Satellite (GPS) unit sampling at 10Hz (Catapult, Optimeye S5, firmware version 7.27, Melbourne, Australia) including a tri-
182 axial accelerometer, gyroscope and magnetometer and sampled at a rate of 100Hz. Devices were located in commercially
183 available vests and sized appropriately for the individual to prevent movement artifact. The mean number of satellites (13.1)
184 and horizontal dilution of precision (HDOP; 0.57) was assessed across all testing sessions to validate data fidelity. Internal
185 load was measured using a T31 Heart Rate (HR) belt (Polar, Finland) and sessional rating of perceived exertion (sRPE)
186 using the centiMax scale (CR100®) obtained within 15-minutes post Y-SAFT⁶⁰. However, this single gestalt sRPE score
187 may oversimplify the psycho-physiological dose-response which is comprised of various physiological and psychological
188 components (i.e. cardiovascular, neuromuscular, cognitive) (McLaren et al., 2017). Therefore, ratings for breathlessness
189 (RPE-B), leg muscle exertion (RPE-L) and cognitive/technical (RPE-T) demands were added, utilising the same reliable
190 scale (ICC = 0.99; TEM = 4%) (McLaren et al., 2017; Wright et al., 2020) and obtained post Y-SAFT⁶⁰ as outlined above.
191 Players were habituated during training sessions and used a numerically blinded, touch-screen tablet (Acer Iconia One 8 B1-
192 850, Taipei, Taiwan; Acer Inc) to record perceptions of intensity using a customised application to provide confidential
193 responses free from conformation bias (Wright et al., 2020).

194
195 Neuromuscular performance was measured with countermovement jump (CMJ), absolute and relative leg stiffness and
196 reactive strength index (RSI) before, at half-time and after the Y-SAFT⁶⁰. Participants had opportunity to familiarise
197 themselves with the protocol after a standardised 5-minute dynamic warm-up consisting of bodyweight activities (e.g.
198 skipping, lunges, squats, hopping) designed to mobilise and activate appropriate muscles. After sufficient rest (3-5 mins)

199 participants completed two attempts of each protocol using the Optojump photocell system (Microgate, Bolzano, Italy) with
200 the best result taken for analysis. CMJ and RSI were calculated from five consecutive, maximal bilateral jumps. Participants
201 started in an upright, standing position with hands on hips and then squatted to a self-selected depth and without pausing
202 jumped maximally five times. Participants were encouraged to perform the eccentric phase as quickly as possible to
203 maximise jump height and minimise ground contact time. Jump height (cm) was calculated from the first jump using flight
204 time from the following equation (equation 1) (Oliver et al., 2015). Jump height and ground contact time were averaged
205 across the five rebound maximal bilateral jumps to calculate RSI using the following equation (equation 2).
206

207 *Equation 1*

$$\text{Jump height} = (\text{Flight time}^2 * \text{gravity}) / 8$$

209 *Equation 2*

$$\text{RSI} = \text{jump height (m)} / \text{ground contact time (s)}$$

211

212 Absolute and relative leg stiffness was measured from contact times and flight times during 20 consecutive bilateral sub-
213 maximal hops at a frequency of 2.5 Hz. This tempo has the highest reliability of leg stiffness measured in adolescent
214 populations (CV 7.2%) (De Ste Croix et al., 2015). Participants were asked to place hands on their hips to minimise upper
215 body interference; rebound for height and land within the photocell gates; landing with legs fully extended and looking
216 forwards. Absolute leg stiffness (K_{leg}) was calculated using equation 3 where K_{leg} refers to leg stiffness, M is total body
217 mass, T_c refers to ground contact time and T_f is equal to flight time. To account for the influence of mass on leg stiffness
218 and leg length on mechanical properties of locomotion between participants, absolute values of leg stiffness were divided
219 by body mass and leg length to provide a dimensionless value of relative leg stiffness (De Ste Croix et al., 2017).
220

221 *Equation 3: Absolute Leg Stiffness*

$$K_{\text{leg}} = [M * \pi (T_f + T_c)] / T_c [T_f + T_c / \pi] - (T_c / 4)$$

223

224 *Data Analysis*

225 To investigate whether PAH% influenced responses in neuromuscular performance, a within-participant repeated measures
226 moderation model was employed. Relationships between the focal predictor (Y_1) and the outcome (Y_2) were examined by
227 testing interactions between these and the stable moderating variable (W_1) utilising a simple moderation model (Montoya,
228 2019)¹:

$$Y_{i2} - Y_{i1} = b_{01} + (b_{12} - b_{11}) W_i + (\varepsilon_{i2} - \varepsilon_{i1})$$

$$Y_{Di} = b_0 + b_1 W_i + \varepsilon_i$$

231

232 Ordinary least squares (OLS) regression equations for pre-post changes in neuromuscular performance were conducted using
233 the SPSS macro MEMORE v2.1 (Montoya, 2019). The moderator (PAH%) and focal predictor (e.g. baseline CMJ) were
234 mean centred to eliminate the effects of multicollinearity between variables and to allow the intercept to equal the overall
235 treatment effect (Wu & Zumbo, 2008).
236

¹ where i = individual participant; i or 2 denotes instance; b_0 intercept; b_1 = regression weight; ε = normally distributed residuals with Intercept variance

237 Analysis to explore whether PAH% moderated interactions between mean centred predictor variable (i.e. total distance) and
 238 outcome variable (e.g. player load) was conducted using a simple moderator model in the SPSS macro PROCESS v3.4
 239 (Hayes & Rockwood, 2017). Due to the continuous nature of all variables, interactions for both analysis methods were
 240 further analysed using the Johnson-Neyman procedure to identify points of transition (i.e. boundaries of significance) across
 241 the whole data range, rather than across arbitrarily selected points (Montoya, 2019). If the 95% confidence intervals (CI) did
 242 not contain zero, the test was considered significant at the $p < .05$ level.

243
 244 **Results**

245 The descriptive changes in neuromuscular performance are shown in table 1 and the pooled descriptive internal and external
 246 load metrics derived from the Y-SAFT⁶⁰ are shown in table 2.

Table 1. Pooled changes (Mean ± SD) in CMJ, RSI and Stiffness (Absolute and Relative) throughout the Y-SAFT⁶⁰

	Pre	HT	% Change	Pre	FT	% Change	HT	FT	% Change
CMJ	25.10 ± 5.0	24.90 ± 6	-0.8%	25.10 ± 5	23.90 ± 6	-4.7%	24.90 ± 5.5	23.90 ± 6.0	-4.1%
RSI	0.55 ± 0.20	0.60 ± 0.2	9.1%	0.55 ± 0.2	0.57 ± 0.2	3.6%	0.60 ± 0.22	0.57 ± 0.2	5.0%
Absolute Stiffness	36.60 ± 11.0	37.60 ± 10	2.7%	36.60 ± 11	35.20 ± 11	-3.8%	37.60 ± 10	35.20 ± 12.0	-6.3%
Relative Stiffness	57.30 ± 17.0	62.20 ± 16	8.5%	57.30 ± 17	58.90 ± 17	2.7%	62.20 ± 16	58.90 ± 17.0	-5.3%

CMJ, countermovement jump; RSI, reactive strength index; Pre, before SAFT⁶⁰; HT, half-time; FT, full-time

247

Table 2. Pooled internal and external load metrics (mean ± SD) for the Y-SAFT⁶⁰

	Mean ± SD
<i>Internal Load</i>	
sRPE (AU)	65 ± 14
RPE-B (AU)	55 ± 19
RPE-L (AU)	67 ± 17
RPE-T (AU)	43 ± 24
Heart Rate (bpm)	166 ± 10
<i>External Load</i>	
Total Distance (m)	5139 ± 226
Metres per minute (m.min ⁻¹)	85 ± 4
Player Load (Au)	651 ± 69

sRPE; sessional rating of perceived exertion; RPE-B, breathlessness; RPE-L, leg muscle exertion; RPE-T, technical exertion

248 Within-participant moderation analysis indicated that there
 249 were no significant interactions between percentage of
 250 PAH% and neuromuscular response post Y-SAFT⁶⁰.
 251 However, both absolute and relative stiffness indicated that
 252 PAH% explained approximately 27-28% of the variance
 253 from the model respectively (Table 3). Although there was
 254 a clear decline in countermovement jump (-4.7%) and a
 255 slight improvement in overall RSI performance (3.6%)
 256 these were not considered significant. A similar outcome
 257 emerged when analysing the interactions between PAH%
 258 and load metrics, whereby only RPE-T produced a
 259 significant interaction, indicating that approximately 10%
 260 ($R^2 = 0.10$) of the variation explained by PAH% (Table 3).

261
 262
 263
 264
 265
 266
 267

Table 3. Regression model characteristics of training load metrics moderated by percentage of predicted adult height

Estimate	R ²	Coefficient	SE	t value	p-value	95% CI
<i>sRPE ~ T^{Dist}</i>	0.06	0.01	0.01	1.19	0.23	-0.01 to 0.02
<i>sRPE ~ PAH</i>		-0.51	0.39	-1.31	0.19	-1.29 to 0.27
<i>sRPE ~ T^{Dist} x PAH</i>	0.01	-0.01	0.01	-0.72	0.46	-0.01 to 0.01
<i>RPE-B ~ T^{Dist}</i>	0.05	-0.00	0.01	-0.07	0.94	-0.02 to 0.12
<i>RPE-B ~ PAH</i>		-0.78	0.53	-1.48	0.14	-1.84 to 0.27
<i>RPE-B ~ T^{Dist} x PAH</i>	0.01	-0.01	0.01	-0.96	0.33	-0.01 to 0.01
<i>RPE-L ~ T^{Dist}</i>	0.07	0.01	0.01	1.65	0.10	-0.01 to 0.03
<i>RPE-L ~ PAH</i>		-0.46	0.46	-1.00	0.32	-1.39 to 0.46
<i>RPE-L ~ T^{Dist} x PAH</i>	0.01	-0.01	0.01	-0.41	0.68	-0.01 to 0.01
<i>RPE-T ~ T^{Dist}</i>	0.13	-0.01	0.01	0.37	0.70	-0.02 to 0.03
<i>RPE-T ~ PAH</i>		-1.73	0.62	-2.77	0.00*	-2.99 to -0.47
<i>RPE-T ~ T^{Dist} x PAH</i>	0.10	-0.01	0.01	-2.50	0.01*	-0.01 to -0.01
<i>HR ~ T^{Dist}</i>	0.02	0.01	0.01	1.18	0.23	-0.01 to 0.02
<i>HR ~ PAH</i>		-0.05	0.29	-0.20	0.84	-0.64 to 0.2
<i>HR ~ T^{Dist} x PAH</i>	0.01	-0.01	0.01	-0.08	0.93	-0.01 to 0.01
<i>PL ~ T^{Dist}</i>	0.23	0.11	0.04	2.93	0.04*	0.03 to 0.19
<i>PL ~ PAH</i>		-4.58	1.7	-2.69	0.00*	-8.00 to -1.17
<i>PL ~ T^{Dist} x PAH</i>	0.01	-0.00	0.00	-2.59	0.12	-0.01 to -0.00
<i>CMJ</i>	0.01	-0.02	0.36	-0.79	0.42	-0.10 to 0.43
<i>RSI</i>	0.01	0.01	0.01	0.23	0.81	-0.04 to .01
<i>Absolute Stiffness</i>	0.27	-0.11	0.09	-1.22	0.22	-0.30 to 0.07
<i>Relative Stiffness</i>	0.28	-0.20	0.16	-1.25	0.27	-0.52 to 0.12

sRPE, sessional rating of perceived exertion; RPE-B, breathlessness; RPE-L, leg muscle exertion; RPE-T, cognitive/technical demand; HR, heart rate;

PL, player load; T^{Dist}, total distance; PAH, predicted adult height; CMJ, countermovement jump; RSI, reactive strength index

SE, standard error; CI, confidence interval



268 Johnson-Neyman slope analysis of data identified that specific ranges of four variables (PL, RSI, absolute and relative
 269 stiffness) showed some evidence of moderation by PAH% (Figure 1). RSI demonstrated a significant slope from 86.3% of
 270 PAH% with less mature individuals experiencing smaller changes in RSI over the course of the Y-SAFT⁶⁰. Similarly, smaller
 271 changes in absolute leg stiffness were evident in less mature individuals, with slope significance observed from 89.3%
 272 PAH% (Figure 1). In contrast, this trend reversed for relative stiffness, where there was evidence of slope significance across
 273 the PAH% spectrum to ~100%. Additionally, it appears that there is evidence of slope significance between 87.9-96.1 PAH%
 274 for PL, but not for other load variables (Figure 1).

275



276 Discussion

277 This study aimed to quantify neuromuscular performance and psycho-physiological responses to a soccer-specific activity
 278 profile and the influence of maturation in EPPP academy players. Firstly, T^{Dist}, HR and sRPE during the Y-SAFT⁶⁰ were
 279 comparable with data previously reported from U12 to U16 age-competition, however the intensity of 85 m.min⁻¹ was below
 280 data previously reported (98-118 m.min⁻¹) (Harley et al., 2010; Wrigley et al., 2012). Despite this slight reduced intensity,
 281 the Y-SAFT⁶⁰ provided a locomotive stimulus comparable to previously reported activity (Table 2). Primary findings from
 282 these data illustrate an interaction between perceived psycho-physiological load (RPE-T) and maturation, with absolute
 283 stiffness, relative stiffness and PL showing slope significance across various stages of maturation. These interactions suggest
 284 that psycho-physiological dose responses are influenced by maturation and should be considered for training prescription
 285 purposes.



286





287





288



289



290



291

292

Figure 1. Johnson-Neyman slope analysis illustrating responses of neuromuscular performance and training load variables with predicted adult height (%) as the moderator, with thin vertical lines illustrating regions of significance.

293 Mean pooled data (Table 1) illustrates RSI performance improved by >3% during the activity, which is likely a potentiation
294 effect and one that opposes previous findings (De Ste Croix et al., 2019; Lehnert et al., 2018). These studies utilised actual
295 game-play (De Ste Croix et al., 2019) and the SAFT⁹⁰ (Lehnert et al., 2018) which provided greater stimulus duration,
296 possibly explaining the differences. However, slope analysis indicated a linear trend whereby more mature individuals
297 experienced greater potentiation. Radnor et al. (2018) explained that SSC function develops during maturation through
298 increased elastic energy utilisation, neural potentiation and more efficient stretch-reflex combined with a higher force
299 production propensity. The improved tendon stiffness, motor unit-recruitment and electro-mechanical delay may result in
300 more mature individuals coping better with activity demands and therefore demonstrating more positive responses to the
301 same stimulus. Less mature individuals may experience greater inhibition of SSC function, in turn elevating stress on
302 contractile components and resisting limb deformation during the spring-mass model. This places a greater emphasis on
303 strength characteristics of muscle (compliance) leading to longer propulsive contact times (De Ste Croix et al., 2019; Lehnert
304 et al., 2017). Therefore, with tendon cross-sectional area (CSA) having large influences on its function and CSA increasing
305 with growth, it is unsurprising that more mature individuals demonstrate efficient SSC function (Radnor et al., 2018) and,
306 therefore, more favourable RSI responses from the standardised activity.

307
308 This inhibition of SSC function in less mature individuals can partly explain the responses to CMJ, absolute and relative leg
309 stiffness. Pooled means indicate reduced CMJ of 4.1% and leg stiffness of 5.3-6.3% respectively with evidence of a
310 significant slope from ~89% PAH for absolute stiffness and across maturation for relative stiffness (Figure 1). Mechanically
311 this translates to increased yielding in the less mature individuals, with greater ground contact times, centre of mass (CoM)
312 displacement and reduced movement efficiency (Radnor et al., 2018). Yielding is caused by the tendon being stiffer than the
313 contractile element of the musculotendinous unit reducing the ability to utilise the elastic storage potential of the tendon.
314 Therefore, less mature individuals fail to generate sufficient muscle stiffness through adequate recruitment of motor units
315 and correct activation strategies (Radnor et al., 2018). This scenario may increase injury risk in the less mature individuals,
316 via increased relative biomechanical load due to increased absorption of shear stress and greater reliance on the knee
317 compared to more mature counterparts (Lloyd et al., 2012). This notion is similar to that reported in other sports, namely
318 ballet where characteristically training loads associate closely with chronological age and where later maturing individuals
319 experience crucial training periods in conjunction with PHV, often resulting in negative injury outcomes (Bradshaw et al.,
320 2014; Mitchell et al., 2020). Findings relating to stiffness are similar to those proposed by De Ste Croix et al. (2017) in
321 adolescent female soccer, whereby the negative impact of soccer activity on leg stiffness reduced as the maturation increased,
322 with prepubescent individuals having the most substantial performance decrements and heightened risk of knee injury. In
323 general, the lack of model significance supports assertions by Oliver et al. (2014) and De Ste Croix et al. (2019) in that leg
324 stiffness responses are highly individualised and improvements in one neuromuscular component are not necessarily
325 mirrored in others, despite sharing many SSC characteristics (Lloyd et al., 2012).

326
327 Variations in locomotive movement patterns may be linked to interactions observed between PL and PAH% highlighted in
328 Figure 1. Although PL appeared consistent through the PAH% range, slope analysis suggests there is greater dose-response
329 variation at the tails of maturation (considered pre- or post-PHV) with a significant slope identified almost exactly aligned
330 with PHV (87.9-96.1% PAH; Figure 1). Tentatively, this may suggest that load-response patterns are influenced around
331 PHV from a single exercise bout. Admittedly, the likely larger variation in anthropometric and physical properties at the
332 tails of maturation could have also influenced the variation observed either side of the significant slope (Figueiredo et al.,

2010). PL is a measure of the instantaneous rate of change in acceleration in each of the three vectors and can infer biomechanical load, though its accuracy for assessing load-response pathways at a structural level is yet to be determined (Verheul et al., 2020). However, a similar response that aligned with PHV (88-93% PAH) was observed in both absolute and relative stiffness. Therefore, in light of a fairly small sample size, interpretations should be cautious and treated as an informed possibility that whole-body load-response pathways are influenced by PAH% with further empirical investigation required. More frequent use of accelerometry within academy soccer will permit this work on which has the potential to provide valuable insights for injury prevention within youth soccer and more widely youth sport.

In general, internal load metrics were not moderated by PAH%, but analysis indicated an interaction with RPE-T and Figure 1 illustrates a progressive reduction in perceived RPE-T as PAH% increases. In line with other research (McLaren et al., 2017) RPE-T was perceived to be lower than other RPE metrics, likely because the Y-SAFT⁶⁰ is a repetitive audio-controlled activity profile requiring minimal cognitive strain in comparison to training and match activities. Why less mature individuals perceive the Y-SAFT⁶⁰ more technically demanding remains unclear, although one theory is linked to the relative physical shortcomings explained previously. Lloyd et al. (2012) suggested that changes in SSC during growth are associated with improvements in anaerobic performance such as sprinting and change of direction, which may have indirectly influenced perceptions of technical effort. For example, Y-SAFT⁶⁰ audio cues such as 'stride' or 'sprint' require maximal effort. A highly intense period during the Y-SAFT⁶⁰ could be perceived more cognitively demanding for less mature individuals endeavouring to maintain pace with the audio. An alternative theory is that less mature individuals experienced greater training monotony during the repetitive activity profile or non-linear differences in physical and cognitive maturation could have influenced perceptions (Brink et al., 2010; Wright et al., 2020). Despite no interaction, other dimensions of intensity (sRPE, RPE-B and HR) followed a similar trend whereby more mature individuals found the activity less intense (Figure 1). This would confirm the intuitive hypotheses, but lack of statistical significance prevents this conclusion; again, suggesting the need for more work in this area. It would also appear that perceptions of exertion during adolescence are influenced by the specific protocol and training mode which may have contributed to these findings (McLaren et al., 2017).

358 *Limitations*

359 Whilst the standardised Y-SAFT⁶⁰ facilitated a dose-response comparison, it is conceded that it disregards many reactive soccer-specific actions (e.g. responding to opponents, jumping and kicking). Consequently, although comparative from a locomotive perspective, the Y-SAFT⁶⁰ may underrepresent true anatomical, physiological and biomechanical stresses placed on adolescent players during actual training and/or competition. This may explain why some individuals experienced a level of potentiation in neuromuscular performance as the demands were insufficient to elicit detrimental performance changes. However, the large positional and individual variability of external loads from competitive matches and training activities make more ecologically valid dose-response protocols highly complex for research. In addition, it is appreciated that baseline markers of physical fitness (e.g. aerobic fitness, strength, speed) may influence individual responses to the standardised activity profile and therefore should not be discounted when considering the moderating impact of PAH%.

369 This acute study suggests that a single soccer specific activity bout can elicit maturation specific dose-responses (RSI, RPE-T, Stiffness), which are likely magnified when combined with repeated activities each week. Therefore, practitioners are urged to consider the maturational load-response variation in an attempt to reduce injury incidence from inappropriate levels of physical and cognitive stress (Brink et al., 2010). Based on tentative findings, it would be irresponsible to make robust

373 recommendations regarding how to change training practices, but the authors would encourage practitioners to be mindful
374 of dose-response variations across maturation in the following ways. Firstly, activities high in biomechanical load
375 characterised by short, intense intervals with frequent accelerations, decelerations and changes of direction, often over small
376 to moderate pitch-dimensions may accentuate variations in response. Therefore, it may be prudent to restrict the frequency
377 and volume of exposure to such load across the training week, particularly in those approaching peak-height velocity.
378 Secondly, practitioners are encouraged to adapt their prescription processes and routinely consider maturation specific load
379 management strategies whenever possible, for example utilising existing maturity-categorised strategies such as bio-banded
380 training or competitions (Cumming et al., 2018). Although this does not automatically reduce the load on those deemed
381 vulnerable, it provides opportunity for the coach to better manage the frequency and volume of exposure to stressful activities
382 on a group level and therefore reduce large variations in inter-individual response attributed to maturity. Finally, this would
383 facilitate the inclusion of activities that required less unpredictable, reactive strength (typified by deceleration or COD)
384 actions in favour of more structured ‘technical’ practices that built on key movement competencies such as acceleration,
385 deceleration, change of direction and strength development as suggested by the Youth Physical Development (YPD) model
386 (Lloyd & Oliver, 2012). It is not recommended that practitioners reduce training exposure per se or remove any particular
387 activities completely, but more that the emphasis of training shifts to allow smart prescription of loads that are directed at
388 the individual appropriately.

389

390 *Conclusion*

391 Collectively, these findings provide evidence of the moderating effect of maturation to perceptions and responses to
392 standardised activity. Therefore, the findings suggest that failure to consider maturity-related responses may elevate the
393 predisposition to injury of some athletes. Both psycho- and physiological responses are altered during maturation which
394 must be considered by practitioners within their training prescription to avoid negative consequences on their players.
395 Strategies already exist within research and practice to facilitate maturity specific training interventions which may help
396 coaches manage the individual dose-response smartly and practically within a group environment. More work is needed to
397 corroborate these findings and provide exact mechanisms of the causes; however, tentative inferences suggest that repeated
398 high-velocity biomechanical loads may contribute to the observed differences in neuromuscular response across maturation
399 and should be managed considerately.

400

401 **Acknowledgements**

402 The author team would like to thank the clubs involved with the study for facilitating the data collection.

403

404 **Declarations of interest**

405 The authors report no conflicts of interest.

406

407

408

409

410

411

412

413 **References**

- 414 Altman, D. G., & Royston, P. (2006). The cost of dichotomising continuous variables. *British Medical Journal*, *6*(332),
415 1080. <https://doi.org/10.1136/bmj.332.7549.1080>
- 416 Barrett, S., Guard, A., & Lovell, R. (2013). *SAFT90 simulates the internal and external loads of competitive soccer match-*
417 *play* (pp. 95–100).
- 418 Bergeron, M. F., Mountjoy, M., Armstrong, N., Chia, M., Côté, J., Emery, C. A., Faigenbaum, A., Hall, G., Kriemler, S.,
419 Léglise, M., Malina, R. M., Pensgaard, A. M., Sanchez, A., Soligard, T., Sundgot-Borgen, J., van Mechelen, W.,
420 Weissensteiner, J. R., & Engebretsen, L. (2015). International Olympic Committee consensus statement on youth
421 athletic development. *British Journal of Sports Medicine*, *49*(13), 843–851. [https://doi.org/10.1136/bjsports-2015-](https://doi.org/10.1136/bjsports-2015-094962)
422 [094962](https://doi.org/10.1136/bjsports-2015-094962)
- 423 Beyer, K. S., Stout, J. R., Redd, M. J., Baker, K. M., Church, D. D., Bergstrom, H. C., Hoffman, J. R., & Fukuda, D. H.
424 (2020). Effect of somatic maturity on the aerobic and anaerobic adaptations to sprint interval training.
425 *Physiological Reports*, *8*(9). <https://doi.org/10.14814/phy2.14426>
- 426 Bradshaw, E., Karin, J., Bowermann, E., & Hume, P. (2014). *Growth and injury patterns of aspiring classical ballet*
427 *dancers*.
- 428 Brink, M. S., Visscher, C., Arends, S., Zwerver, J., Post, W. J., & Lemmink, K. A. (2010). Monitoring stress and recovery:
429 New insights for the prevention of injuries and illnesses in elite youth soccer players. *British Journal of Sports*
430 *Medicine*, *44*(11), 809–815. <https://doi.org/10.1136/bjism.2009.069476>
- 431 Buchheit, M., & Mendez-Villanueva, A. (2014). Effects of age, maturity and body dimensions on match running
432 performance in highly trained under-15 soccer players. *Journal of Sports Sciences*, *32*(13), 1271–1278.
433 <https://doi.org/10.1080/02640414.2014.884721>
- 434 Cumming, S. P., Brown, D. J., Mitchell, S., Bunce, J., Hunt, D., Hedges, C., Crane, G., Gross, A., Scott, S., Franklin, E.,
435 Breakspear, D., Dennison, L., White, P., Cain, A., Eisenmann, J. C., & Malina, R. M. (2018). Premier League
436 academy soccer players' experiences of competing in a tournament bio-banded for biological maturation. *Journal*
437 *of Sports Sciences*, *36*(7), 757–765. <https://doi.org/10.1080/02640414.2017.1340656>
- 438 Cumming, S. P., Lloyd, R. S., Oliver, J. L., Eisenmann, J. C., & Malina, R. M. (2017). Bio-banding in Sport: Applications
439 to Competition, Talent Identification, and Strength and Conditioning of Youth Athletes. *Strength and*
440 *Conditioning Journal*, *39*(2), 34–47. <https://doi.org/10.1519/SSC.0000000000000281>
- 441 De Ste Croix, M., Hughes, J. D., Lloyd, R. S., Oliver, J. L., & Read, P. J. (2017). Leg Stiffness in Female Soccer Players:
442 Intersession Reliability and the Fatiguing Effects of Soccer-Specific Exercise. *Journal of Strength and*
443 *Conditioning Research*, *31*(11), 3052–3058. <https://doi.org/10.1519/JSC.0000000000001715>

- 444 De Ste Croix, M., Lehnert, M., Maixnerova, E., Zaatari, A., Svoboda, Z., Botek, M., Varekova, R., & Stastny, P. (2019).
445 Does maturation influence neuromuscular performance and muscle damage after competitive match-play in youth
446 male soccer players? *European Journal of Sport Science*, *19*(8), 1130–1139.
447 <https://doi.org/10.1080/17461391.2019.1575913>
- 448 De Ste Croix, M., Priestley, A. M., Lloyd, R. S., & Oliver, J. L. (2015). ACL injury risk in elite female youth soccer:
449 Changes in neuromuscular control of the knee following soccer-specific fatigue: Female youth soccer and fatigue.
450 *Scandinavian Journal of Medicine & Science in Sports*, *25*(5), e531–e538. <https://doi.org/10.1111/sms.12355>
- 451 Epstein, L. H., Valoski, A. M., Kalarchian, M. A., & McCurley, J. (1995). Do Children Lose and Maintain Weight Easier
452 Than Adults: A Comparison of Child and Parent Weight Changes From Six Months to Ten Years. *Obesity*
453 *Research*, *3*(5), 411–417. <https://doi.org/10.1002/j.1550-8528.1995.tb00170.x>
- 454 Fanchini, M., Ferraresi, I., Petruolo, A., Azzalin, A., Ghielmetti, R., Schena, F., & Impellizzeri, F. M. (2017). Is a
455 retrospective RPE appropriate in soccer? Response shift and recall bias. *Science and Medicine in Football*, *1*(1),
456 53–59. <https://doi.org/10.1080/02640414.2016.1231411>
- 457 Figueiredo, A. J., Silva, M. J. C. e, Cumming, S. P., & Malina, R. M. (2010). Size and Maturity Mismatch in Youth Soccer
458 Players 11- to 14-Years-Old. *Pediatric Exercise Science*, *22*(4), 596–612. <https://doi.org/10.1123/pes.22.4.596>
- 459 Foster, C. (1998). Monitoring training in athletes with reference to overtraining syndrome: *Medicine & Science in Sports*
460 *& Exercise*, *30*(7), 1164–1168. <https://doi.org/10.1097/00005768-199807000-00023>
- 461 Hader, K., Rumpf, M. C., Hertzog, M., Kilduff, L. P., Girard, O., & Silva, J. R. (2019). Monitoring the Athlete Match
462 Response: Can External Load Variables Predict Post-match Acute and Residual Fatigue in Soccer? A Systematic
463 Review with Meta-analysis. *Sports Medicine - Open*, *5*(1), 48. <https://doi.org/10.1186/s40798-019-0219-7>
- 464 Hannon, M. P., Carney, D. J., Floyd, S., Parker, L. J. F., McKeown, J., Drust, B., Unnithan, V. B., Close, G. L., & Morton,
465 J. P. (2020). Cross-sectional comparison of body composition and resting metabolic rate in Premier League
466 academy soccer players: Implications for growth and maturation. *Journal of Sports Sciences*, 1–9.
467 <https://doi.org/10.1080/02640414.2020.1717286>
- 468 Harley, J. A., Barnes, C. A., Portas, M., Lovell, R., Barrett, S., Paul, D., & Weston, M. (2010). Motion analysis of match-
469 play in elite U12 to U16 age-group soccer players. *Journal of Sports Sciences*, *28*(13), 1391–1397.
470 <https://doi.org/10.1080/02640414.2010.510142>
- 471 Hayes, A. F., & Rockwood, N. J. (2017). Regression-based statistical mediation and moderation analysis in clinical
472 research: Observations, recommendations, and implementation. *Behaviour Research and Therapy*, *98*, 39–57.
473 <https://doi.org/10.1016/j.brat.2016.11.001>

474 Hill, M., Scott, S., Malina, R. M., McGee, D., & Cumming, S. P. (2020). Relative age and maturation selection biases in
475 academy football. *Journal of Sports Sciences*, 38(11–12), 1359–1367.
476 <https://doi.org/10.1080/02640414.2019.1649524>

477 Johnson, A., Farooq, A., & Whiteley, R. (2017). Skeletal maturation status is more strongly associated with academy
478 selection than birth quarter. *Science and Medicine in Football*, 1(2), 157–163.
479 <https://doi.org/10.1080/24733938.2017.1283434>

480 Johnson, D., Williams, S., Bradley, B., Sayer, S., Fisher, J. M., & Cumming, S. (2020). Growing pains: Maturity
481 associated variation in injury risk in academy football. *European Journal of Sport Science*, 20(4), 544–552.
482 <https://doi.org/10.1080/17461391.2019.1633416>

483 Khamis, H. J., & Roche, A. F. (1994). Predicting Adult Stature Without Using Skeletal Age: The Khamis-Roche Method.
484 *Pediatrics*, 94(4), 504–507.

485 Lehnert, M., Croix, M. D. S., Xaverova, Z., Botek, M., Varekova, R., Zaatar, A., Lastovicka, O., & Stastny, P. (2018).
486 Changes in Injury Risk Mechanisms After Soccer-Specific Fatigue in Male Youth Soccer Players. *Journal of*
487 *Human Kinetics*, 62(1), 33–42. <https://doi.org/10.1515/hukin-2017-0157>

488 Lehnert, M., De Ste Croix, M., Zaatar, A., Hughes, J., Varekova, R., & Lastovicka, O. (2017). Muscular and
489 neuromuscular control following soccer-specific exercise in male youth: Changes in injury risk mechanisms.
490 *Scandinavian Journal of Medicine & Science in Sports*, 27(9), 975–982. <https://doi.org/10.1111/sms.12705>

491 Lloyd, R. S., & Oliver, J. L. (2012). The Youth Physical Development Model: A New Approach to Long-Term Athletic
492 Development. *Strength & Conditioning Journal*, 34(3), 61–72. <https://doi.org/10.1519/SSC.0b013e31825760ea>

493 Lloyd, R. S., Oliver, J. L., Hughes, M. G., & Williams, C. A. (2012). Age-related differences in the neural regulation of
494 stretch–shortening cycle activities in male youths during maximal and sub-maximal hopping. *Journal of*
495 *Electromyography and Kinesiology*, 22(1), 37–43. <https://doi.org/10.1016/j.jelekin.2011.09.008>

496 Lovell, R. (n.d.). *Physiological responses to SAFT90: A new soccer-specific match simulation*. 1.

497 McLaren, S. J., Smith, A., Spears, I. R., & Weston, M. (2017). A detailed quantification of differential ratings of perceived
498 exertion during team-sport training. *Journal of Science and Medicine in Sport*, 20(3), 290–295.
499 <https://doi.org/10.1016/j.jsams.2016.06.011>

500 Mitchell, S. B., Haase, A. M., & Cumming, S. P. (2020). Experiences of delayed maturation in female vocational ballet
501 students: An interpretative phenomenological analysis. *Journal of Adolescence*, 80, 233–241.
502 <https://doi.org/10.1016/j.adolescence.2020.03.005>

503 Montoya, A. K. (2019). Moderation analysis in two-instance repeated measures designs: Probing methods and multiple
504 moderator models. *Behavior Research Methods*, 51(1), 61–82. <https://doi.org/10.3758/s13428-018-1088-6>

505 Oliver, J. L., De Ste Croix, M. B. A., Lloyd, R. S., & Williams, C. A. (2014). Altered neuromuscular control of leg
506 stiffness following soccer-specific exercise. *European Journal of Applied Physiology*, 114(11), 2241–2249.
507 <https://doi.org/10.1007/s00421-014-2949-z>

508 Oliver, J. L., Lloyd, R. S., & Whitney, A. (2015). Monitoring of in-season neuromuscular and perceptual fatigue in youth
509 rugby players. *European Journal of Sport Science*, 15(6), 514–522.
510 <https://doi.org/10.1080/17461391.2015.1063700>

511 Parr, J., Winwood, K., Hodson-Tole, E., Deconinck, F. J. A., Parry, L., Hill, J. P., Malina, R. M., & Cumming, S. P.
512 (2020). Predicting the timing of the peak of the pubertal growth spurt in elite youth soccer players: Evaluation of
513 methods. *Annals of Human Biology*, 0(ja), 1–23. <https://doi.org/10.1080/03014460.2020.1782989>

514 Premier League. (2011). *The Elite Player Performance Plan*. English Premier League.

515 Radnor, J. M., Oliver, J. L., Waugh, C. M., Myer, G. D., & Lloyd, R. S. (2020). The Influence of Maturity Status on
516 Muscle Architecture in School-Aged Boys. *Pediatric Exercise Science*, 32(2), 89–96.
517 <https://doi.org/10.1123/pes.2019-0201>

518 Radnor, J. M., Oliver, J. L., Waugh, C. M., Myer, G. D., Moore, I. S., & Lloyd, R. S. (2018). The Influence of Growth and
519 Maturation on Stretch-Shortening Cycle Function in Youth. *Sports Medicine*, 48(1), 57–71.
520 <https://doi.org/10.1007/s40279-017-0785-0>

521 Read, P. J., Oliver, J. L., De Ste Croix, M. B. A., Myer, G. D., & Lloyd, R. S. (2016). Neuromuscular Risk Factors for
522 Knee and Ankle Ligament Injuries in Male Youth Soccer Players. *Sports Medicine*, 46(8), 1059–1066.
523 <https://doi.org/10.1007/s40279-016-0479-z>

524 Read, P. J., Oliver, J. L., De Ste Croix, M. B. A., Myer, G. D., & Lloyd, R. S. (2018). An audit of injuries in six english
525 professional soccer academies. *Journal of Sports Sciences*, 36(13), 1542–1548.
526 <https://doi.org/10.1080/02640414.2017.1402535>

527 Rommers, N., Rössler, R., Goossens, L., Vaeyens, R., Lenoir, M., Witvrouw, E., & D’Hondt, E. (2020). Risk of acute and
528 overuse injuries in youth elite soccer players: Body size and growth matter. *Journal of Science and Medicine in*
529 *Sport*, 23(3), 246–251. <https://doi.org/10.1016/j.jsams.2019.10.001>

530 Salter, J., De Ste Croix, M., Hughes, J., Weston, M., & Towlson, C. (2020). Monitoring practices of training load and
531 biological maturity in UK soccer academies. *International Journal of Sports Physiology and Performance*, 28.

- 532 Small, K., McNaughton, L. R., Greig, M., Lohkamp, M., & Lovell, R. (2009). Soccer Fatigue, Sprinting and Hamstring
533 Injury Risk. *International Journal of Sports Medicine*, 30(08), 573–578. <https://doi.org/10.1055/s-0029-1202822>
- 534 van der Sluis, A., Elferink-Gemser, M., Brink, M., & Visscher, C. (2015). Importance of Peak Height Velocity Timing in
535 Terms of Injuries in Talented Soccer Players. *International Journal of Sports Medicine*, 36(04), 327–332.
536 <https://doi.org/10.1055/s-0034-1385879>
- 537 Verheul, J., Nedergaard, N. J., Vanrenterghem, J., & Robinson, M. A. (2020). Measuring biomechanical loads in team
538 sports – from lab to field. *Science and Medicine in Football*, 1–7.
539 <https://doi.org/10.1080/24733938.2019.1709654>
- 540 Whalan, M., Lovell, R., & Sampson, J. A. (2020). Do Niggles Matter? - Increased injury risk following physical
541 complaints in football (soccer). *Science and Medicine in Football*, 4(3), 216–224.
542 <https://doi.org/10.1080/24733938.2019.1705996>
- 543 Wright, M. D., Songane, F., Emmonds, S., Chesterton, P., Weston, M., & McLaren, S. J. (2020). *Differential ratings of*
544 *perceived match and training exertion in youth female soccer: Differential RPE in girls soccer.*
- 545 Wrigley, R., Drust, B., Stratton, G., Scott, M., & Gregson, W. (2012). Quantification of the typical weekly in-season
546 training load in elite junior soccer players. *Journal of Sports Sciences*, 30(15), 1573–1580.
547 <https://doi.org/10.1080/02640414.2012.709265>
- 548 Wu, A. D., & Zumbo, B. D. (2008). Understanding and Using Mediators and Moderators. *Social Indicators Research*,
549 87(3), 367–392. <https://doi.org/10.1007/s11205-007-9143-1>