



Mair, Jacqueline L., Hayes, Lawrence D., Campbell, Amy ORCID logo ORCID: <https://orcid.org/0000-0003-3711-3896> and Sculthorpe, Nicholas (2021) Should We Use Activity Tracker Data From Smartphones and Wearables to Understand Population Physical Activity Patterns? *Journal for the Measurement of Physical Behaviour*, 5 (1). pp. 3-7.

Downloaded from: <https://ray.yorks.ac.uk/id/eprint/6280/>

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version:
<http://dx.doi.org/10.1123/jmpb.2021-0012>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repositories Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at
ray@yorks.ac.uk

1 **Should we use consumer-grade activity tracker data to understand population**
2 **physical activity patterns?**

3

4 Submitted 24 February 2021

5

6 Revised 28 June 2021

7

8 **ABSTRACT**

9

10 Researchers, practitioners, and public health organisations from around the world are
11 becoming increasingly interested in using data from wearable activity trackers, from
12 companies such as Fitbit Inc, Garmin Ltd, Xiaomi, and Apple Inc, to measure physical activity.
13 Indeed, large-scale, easily accessible, and autonomous data collection concerning physical
14 activity as well as other health behaviours is becoming ever more attractive. There are several
15 benefits of using wearable activity trackers to collect physical activity data, including the
16 ability to obtain big data, retrospectively as well as prospectively, to understand individual-
17 level physical activity patterns over time and in response to natural events. However, there
18 are challenges related to representativeness, data access, and proprietary algorithms that, at
19 present, limit the utility of this data in understanding population-level physical activity. In this
20 brief report we aim to highlight the benefits, as well as the limitations, of using existing data
21 from wearable activity trackers to understand large-scale physical activity patterns and
22 stimulate discussion amongst the scientific community on what the future holds with respect
23 to physical activity measurement and surveillance.

24

25 **KEY WORDS**

26 m-Health; quantified-self; big data; surveillance; wearables

INTRODUCTION

Physical activity (PA) and exercise have pronounced positive effects on physical, mental, and social health and wellbeing and, according to recent estimates, prevent 3.9 million premature deaths worldwide annually (Strain et al., 2020). Accordingly, global PA guidelines recommend all adults to undertake 150–300 min of moderate-intensity, or 75–150 min of vigorous-intensity PA, or some equivalent combination per week (Bull et al., 2020). Such guidelines rely on population level surveillance methods to regularly monitor PA indicators and inform public health policy, and the most common approach in this regard is to assess PA using self-report methods. Self-report remains an accepted method of large-scale data collection due to its cost effectiveness, unobtrusiveness and adaptability to different country contexts (Troiano et al., 2020). This is despite the accepted limitations of self-reporting with respect to accuracy, recall bias and social-desirability (Brenner & DeLamater, 2014; Prince et al., 2008). Advances in technology over the last two decades have, however, created new possibilities for PA measurement, not only for population-level surveillance but at an individual-level in terms of cohort studies, intervention research, and the evaluation of public health promotion programs. Research-grade devices such as wrist, hip, and thigh worn accelerometers have been used widely in such studies as they remove the biases associated with self-reporting and are able to provide a more granular quantification of PA. Nevertheless, research-grade accelerometers are costly to use at scale and cannot assess the domain or context in which PA takes place. Furthermore, accelerometers, as with self-report methods, only offer a ‘snapshot in time’ to infer usual PA behaviour (typically a 7-day period) meaning assessment of longer-term dynamic patterns of PA, particularly in response to natural events, is either not possible or not feasible. Over the last 15 years, the emergence of consumer-grade devices, such as smartphones and wearable activity trackers, has opened new doors in the

field of PA measurement. These devices gather rich activity data continuously in a free-living setting thus providing large-scale and low-cost datasets that could advance our understanding of PA patterns in a way that was never possible before. While this seems an exciting prospect, as with any other PA measurement tool, the use of wearable activity tracker data should be carefully considered before being used in PA research.

Comparing and contrasting all the available PA measurement methods available to researchers, practitioners, and public health professionals is beyond the scope of this brief report. Instead, we focus on the emerging opportunities offered by consumer grade devices, including smartphones and wearable activity trackers, and how these may be utilised in the fields of PA surveillance, cohort studies, intervention research, and evaluation of public health promotion programs.

Consumer-grade devices: too good to ignore?

Technological advancements over the last decade, allied to the rapid proliferation of smartphone use in both developed and developing regions globally (Deloitte, 2017), have provided new possibilities in monitoring, understanding, and influencing human movement at scale. Compared to traditional approaches, objective, real-world PA data sets with very large sample sizes have become relatively low cost for researchers to collect or access. Consequently, we are now beginning to witness the emergence of big data on PA in the literature. For example, Althoff and colleagues (2017) recently used minute-by-minute step count data, collected from the smartphone's inbuilt inertial unit, from over 700,000 individuals across 111 countries, to identify variability in PA levels across the world. With this

data they revealed city walkability as a factor associated with PA levels as well as associations between PA inequalities and obesity.

The questions that could be addressed, and the new insights afforded, by large-scale PA data from smartphones is an exciting prospect. However, concerns remain over the quality of the PA data that can be obtained from smartphones, including the validity and reliability of step detection, the restriction to only ambulatory activity, and their reliance on the individual carrying the smartphone (Brodie et al., 2018). Wearable activity trackers, although currently less prevalent than smartphones, are growing in popularity (Deloitte, 2017; Thompson, 2019) and can address some of these pitfalls. Activity trackers have progressed beyond simple pedometers and can now provide data on pulse rate, distance covered, moderate-to-vigorous PA minutes, stair flights climbed, energy expenditure, and sleep. Unlike research grade devices such as accelerometers, they also blend attractive design with invisible and effort-free data capture. This combination often results in high adherence (in terms of daily wear time) for extended periods of time. Additionally, while synchronisation of the activity tracker to a smartphone application displays summary information to users, these summary data are calculated from extensive intra-day data gathered at high frequency (e.g., 1 Hz), which are also stored. High adherence alongside high frequency data capture means individuals accumulate an extensive data resource that could be utilised to answer important PA questions.

Big data analyses from smartphones and wearable activity trackers have, thus far, been cross-sectional which limits our understanding of PA to a single point in time. However, longitudinal data from smartphones and wearable activity tracker could also be analysed prospectively or

retrospectively, given its perpetual collection and long-term storage. Of note is the unique opportunity offered by smartphones and wearable activity trackers to analyse data retrospectively and in response to natural events without the need for foresight. This has had obvious implications during the coronavirus pandemic when objective accelerometry was not possible, or feasible, due to the speed and variability with which restrictions were imposed across the world. As a result, the need for remote and scalable means to both measure and support PA has become more prominent since the coronavirus pandemic. Although the use of wearable activity trackers is not without inherent limitations (discussed further below), we feel they are unique in their ability to be utilised in retrospective cohort study designs (i.e., when the start of the study is only known after the event).

Interestingly, to date, there has been almost no independent large-scale reporting of existing data from wearable activity trackers. This might be due to the complexity of large-scale data access and processing from commercial wearables. A cross-sectional study of pulse rate data from over 8 million Fitbit users was recently published by the Fitbit Research team (Natarajan et al., 2020), reporting a positive relationship between heart rate variability and step count. Regardless of the study findings, it seems that collecting, processing, and interpreting this volume of data is possible, but requires an interdisciplinary team including data scientists, database analysts, and cardiovascular and behavioural scientists, which has so far been limited to large proprietary companies such as Fitbit. Furthermore, accessing this volume of data is, so far, only feasible for companies such as Fitbit because they require all users to give them permission to use the data collected by the device. For independent researchers it is possible to request access to the data from the user directly. But this would be on an

individual basis, therefore, to amass a dataset of 8 million users would require 8 million individual access requests. The issue of data access is discussed on more detail below.

Balancing feasibility against validity, reliability, and sensitivity

When choosing a PA data collection tool or methodology, researchers must balance validity, reliability, and sensitivity of the approach with the costs and feasibility of its deployment in the target population. Despite their known limitations (Brenner & DeLamater, 2014; Prince et al., 2008), self-report methods remain an accepted means by which to collect large-scale and population-level PA data, particularly where cost and sample size make accelerometry an unfeasible approach. However, the volume and detail of information that can be obtained from self-report surveys can be limited, preventing more nuanced analysis of PA patterns. Device-based methods, such as accelerometry, provide a more valid and reliable estimate of PA than do self-report measures (Dowd et al., 2018), but also have several limitations. Not only are accelerometers costly, but their data also must be extracted from each device individually, making them unfeasible for large-scale use. Data from wearable activity trackers on the other hand should be considered feasible for large-scale use. Suitable activity trackers are generally cheaper than accelerometers and their attractive design should translate into greater wear time. Like accelerometers, they provide continuous data capture, but with the additional advantage of these data being stored on a central server meaning data retrieval and analysis can occur remotely and at scale. Thus, in principle, it is possible to analyse the PA data of thousands of participants worldwide in a manner that is simply not possible with current research grade accelerometers. Research has shown wearable activity trackers to have high interdevice reliability for measuring steps, energy expenditure, and sleep (Evenson et al., 2015), and despite ongoing concerns, the accuracy of wearable activity trackers also

continues to improve. In a recent systematic review of 67 studies, Fitbit devices were found to provide a relatively accurate measure of free-living steps (within $\pm 10\%$, 50% of the time) when compared to research-grade accelerometers (Feehan et al., 2018). Garmin activity trackers are also reported to have good-to-excellent correlation coefficients and acceptable ($<10\%$) mean absolute percentage errors with respect to step count (Evenson & Spade, 2020). While the accuracy of wearable activity trackers in measuring step count in free-living settings is considered to be acceptable for normal walking pace (Evenson & Spade, 2020; Feehan et al., 2018; Fokkema et al., 2017), they do not yet provide a valid measure of moderate-to-vigorous PA (Redenius et al., 2019) or walking at very slow or very fast speeds (Fokkema et al., 2017). However, considering this evidence is based on devices manufactured up to 2015, refined algorithms over the past 5 years have likely further improved accuracy.

For intervention research the responsiveness, or sensitivity, to change in PA over time may be a more important consideration than the validity of the tool. When examining the effectiveness of an intervention in changing PA it is paramount that the measurement tool employed is capable of detecting change. Research has shown the responsiveness indices for self-report and device-based methods to vary not just by tool, or device, but by PA variable measured. Reeves et al. (2010) compared the responsiveness of the Community Health Activities Model Program for Seniors (CHAMPS) questionnaire, the Active Australia Questionnaire (AAQ), and two items on exercise from the US National Health Interview Survey (USNHIS), and reported responsiveness indices ranging from 0.15 (AAQ) to 0.27 (USNHIS) for walking duration and 0.25 (AAQ) to 0.32 (CHAMPS) for moderate to vigorous intensity PA duration per week. Swartz et al. (2014) compared two research-grade accelerometers, the Actigraph GTX3 (ActiGraph LLC, Pensacola, Florida, USA) and the activPAL (PAL Technologies

Ltd, Glasgow, Scotland, UK) and found both to have comparable responsiveness to change across a range of free living physical activity and sedentary behaviour variables (standardised response mean values between 0.159 – 0.436). Donnachie and colleagues (2020) compared a self-report PA measure (the International Physical Activity Questionnaire; IPAQ) and an accelerometer (activPAL), and found both to have comparable and moderate standardised response mean values of 0.54 (activPAL) and 0.59 (IPAQ) for total PA duration per day. There appears to be no evidence on the responsiveness to change of wearable activity trackers. This surprisingly under-researched topic warrants further attention by the PA research community.

We have an array of options to measure elements of PA (such as duration, intensity, type, domain, context, and quality), but no single tool can fully capture the complexity of PA behaviour. Consumer-grade devices offer new opportunities for combining PA data collection methods. For example, passive sensing of movement using a smartphone or wearable activity tracker, combined with synchronised ‘smart’ self-report techniques, such as ecological momentary assessment, could address many of the issues outlined previously. With further evidence to support the validity, reliability and sensitivity of such methods, this approach could provide powerful insights into PA patterns and help us better understand PA behaviour.

The issue of data harmonisation

Another issue researchers must consider when evaluating device-based PA measurement tools is the harmonisation or comparability between devices from different manufacturers. Data harmonisation is an essential step if researchers wish to conduct analyses on data derived from different sources (Pearce et al., 2020). While all activity tracking devices gather

193 raw uni- or tri-axial accelerations, each manufacturer applies different algorithms to process
194 the data into its summary form thereby influencing the comparability of the data gathered.
195 Therefore, researchers who wish to use data from multiple devices/manufacturers to
196 increase sample representativeness and reach will need to consider data harmonisation using
197 statistical models derived from validation studies (Pearce et al., 2020). This could be
198 problematic when algorithms change, and validation data are no longer available.
199 Manufacturers of research-grade devices publish open source algorithms allowing
200 researchers to evaluate the impact of changes on measurement properties (Evenson et al.,
201 2015), however consumer-grade device manufacturers keep this information proprietary.
202 The use of different proprietary algorithms by each consumer-grade device manufacturer is
203 undoubtedly an issue for harmonisation too. In the longer term, this would be solved by
204 manufacturers making raw data counts available or at least allowing researchers to apply to
205 access this information. However, due to the proprietary nature of data processing, it is
206 unclear if raw data or only processed data are available. In the short-term however,
207 comparative validation between devices should enable statistical techniques that allow for
208 between device data pooling without compromising data quality. Finally, it is also worth
209 noting that there is a small but growing sector of ‘hackable’ wearables. These devices are
210 usually based on small form factor processing boards (e.g., small Raspberry Pi or Arduino
211 boards) which include tri-axial accelerometers, heart rate measurement, WIFI and Bluetooth.
212 These devices also support the remote storage of raw data signals, which would overcome
213 the limitations of unknown and proprietary algorithms. Although useful for research studies,
214 it seems unlikely that such devices will achieve the market penetration of larger
215 manufacturers.

216

The issue of representativeness

Given the widespread use of smartphones and the growing use of activity trackers, we should not ignore the possibility that in the near future wearable activity tracker data could also be used as a population PA surveillance tool. However, at present the primary challenge relating to such data is that it likely over-represents individuals who are more physically active and more proactive in setting and meeting activity goals relative to the general population who may not be tracking their activity level (Omura et al., 2017; Strain et al., 2019). Therefore, any cohort or surveillance research exclusively involving participants who own, and wear, activity trackers will introduce selection bias. The issue of representativeness is, however, not necessarily limited to wearable activity trackers. Selection bias might also occur in data derived from public calls to self-report PA or participate in cohort studies involving self-report or device-based measures of PA. Indeed, it has previously been suggested that selection bias is a significant issue in many cohort studies including those with objective assessments (Barreto et al., 2013; Folley et al., 2018; Stamatakis et al., 2021). Nevertheless, in such cohort and surveillance studies it is possible to use weighting to adjust for non-responders. This is not currently possible for data from wearable activity tracker and future research should focus on statistical approaches to estimate the population effect, and the effect in those with trackers, to help overcome this limitation.

While activity tracker sales and usage are increasing, the demographic reach appears, so far, to be constrained to young adults from more affluent backgrounds (Omura et al., 2017; Strain et al., 2019). Nevertheless, the cost of activity trackers has decreased significantly in recent years making them more affordable and accessible. This, combined with the increasing interest in activity trackers as behaviour change tools, may reduce this constrained

demographic reach over time. For example, recent initiatives to provide activity trackers as part of health care (NHS England, 2019), health improvement (Yao et al., 2020) or health insurance (Buckle et al., 2020) may serve to increase the breadth of the population using the devices. The more initiatives and interventions utilising activity trackers, the more they could be adopted by individuals from underserved populations, such as older adults and those with lower incomes.

The issue of data access

Finally, it is worth noting some of the challenges inherent in accessing data from consumer-grade activity trackers. To access data, researchers can establish an industry agreement with a relevant company (e.g. Fitbit or Garmin) whose terms of service for collecting research data are different from those governing commercial access (Hicks et al., 2019). While the specific manufacturers control access to the data repositories, the data remains the property of the individual user, therefore to access any data collected by the device, each individual user must consent and agree to share the data. Managing thousands, and possibly tens of thousands, of data sharing requests to individual users, and subsequently also having to manage their authorisation and access details, brings its own logistic challenges. The most effective approach is for participants to be directed to a project website which manages participant information, consent, and authorisation requests via the specific manufacturers API. Following successful authorisation, access codes for each user can be securely sent to the research team for subsequent processing. It is worth noting that, even with successful authorisation, there remain additional challenges. Authorisation is usually limited to a maximum of 12 months before the user must re-approve access, which may limit follow-up assessments in very large cohorts where direct contact with participants is limited. In

addition, it is users, not researchers, who define the scope of the data that can be accessed; therefore, users may allow access to all or only some of their data (e.g. only pulse rate, or step count, or some combination thereof), resulting in incomplete data sets. Additionally, most devices allow users to manually add activity to account for any activity not passively detected by the device (e.g. swimming or cycling). At present, it is unclear if such self-reported estimates affect validity. Most databases separate device collected (passive) data from user added (self-reported) data, meaning the research team have to make a decision regarding which should be regarded as the 'canonical' source of users' PA.

Clearly, these challenges are not trivial, and future research teams will require multi-disciplinary skills, including specialists in behavioural science, PA, data science, and software and web development to successfully manage such projects. Nevertheless, if accessed and interpreted appropriately, these data may allow understanding of PA behaviour at a scale previously unimaginable. We are in the process of using this method at a national level to understand the impact of coronavirus, but future research using this technique could examine worldwide PA patterns, both prospectively and retrospectively, using multi-site and multi-lingual research teams.

CONCLUSIONS

As with other device-based and self-report methods, we propose that consumer-grade activity tracker data be considered with their limitations in mind rather than dismissed as a flawed approach, particularly when the feasibility of large-scale accelerometry is prohibitive. Given the rising popularity of wearable activity trackers, the volume of data collected, and the possibilities in analysing data retrospectively, we believe data from wearable activity

trackers should be considered a viable PA measurement tool. To be clear, we are not advocating that other tools, particularly self-report methods, should be consigned to history or replaced by wearable activity tracker 'big data'. Quite the contrary, despite their limitations self-report methods have provided critical insights into PA behaviour and are likely to remain important in the future. Rather, our view is that if physical activity researchers, practitioners, and public health professionals can use and interpret self-report data in light of their limitations, the same should be possible for activity tracker data.

LIST OF ABBREVIATIONS

AAQ	Active Australia Questionnaire
API	Application Programming Interface
CHAMPS	Community Health Activities Model Program for Seniors
IPAQ	International Physical Activity Questionnaire
PA	Physical Activity
USNHIS	US National Health Interview Survey

DECLARATIONS

Ethical Approval

Not Applicable

Consent for Publication

Not Applicable

Competing Interests

None to declare.

Funding

The authors' research referred to in this paper is funded by [REDACTED]

[REDACTED].

Authors' Contributions

All authors contributed equally to the writing of this manuscript.

Acknowledgements

Not Applicable

REFERENCES

Althoff, T., Sosič, R., Hicks, J. L., King, A. C., Delp, S. L., & Leskovec, J. (2017). Large-scale physical activity data reveal worldwide activity inequality. *Nature*, 547(7663), 336–339. <https://doi.org/10.1038/nature23018>

Barreto, P. de S., Ferrandez, A.-M., & Saliba-Serre, B. (2013). Are Older Adults Who Volunteer to Participate in an Exercise Study Fitter and Healthier Than Nonvolunteers? The Participation Bias of the Study Population. *Journal of Physical Activity and Health*, 10(3), 359–367. <https://doi.org/10.1123/jpah.10.3.359>

Brenner, P. S., & DeLamater, J. D. (2014). Social Desirability Bias in Self-reports of Physical Activity: Is an Exercise Identity the Culprit? *Social Indicators Research*, 117(2), 489–504. <https://doi.org/10.1007/s11205-013-0359-y>

337 Brodie, M. A., Pliner, E. M., Ho, A., Li, K., Chen, Z., Gandevia, S. C., & Lord, S. R. (2018). Big
338 data vs accurate data in health research: Large-scale physical activity monitoring,
339 smartphones, wearable devices and risk of unconscious bias. *Medical Hypotheses*,
340 119, 32–36. <https://doi.org/10.1016/j.mehy.2018.07.015>

341 Buckle, J., Hayward, T., Singhal, N., & Desai, K. (2020). *The role of wearables in private*
342 *medical insurance* (p. 24). Milliman.

343 Bull, F. C., Al-Ansari, S. S., Biddle, S., Borodulin, K., Buman, M. P., Cardon, G., Carty, C.,
344 Chaput, J.-P., Chastin, S., Chou, R., Dempsey, P. C., DiPietro, L., Ekelund, U., Firth, J.,
345 Friedenreich, C. M., Garcia, L., Gichu, M., Jago, R., Katzmarzyk, P. T., ... Willumsen, J.
346 F. (2020). World Health Organization 2020 guidelines on physical activity and
347 sedentary behaviour. *British Journal of Sports Medicine*, 54(24), 1451–1462.
348 <https://doi.org/10.1136/bjsports-2020-102955>

349 Deloitte. (2017). *Global mobile consumer trends, 2nd edition*.
350 [https://www2.deloitte.com/content/dam/Deloitte/us/Documents/technology-](https://www2.deloitte.com/content/dam/Deloitte/us/Documents/technology-media-telecommunications/us-global-mobile-consumer-survey-second-edition.pdf)
351 [media-telecommunications/us-global-mobile-consumer-survey-second-edition.pdf](https://www2.deloitte.com/content/dam/Deloitte/us/Documents/technology-media-telecommunications/us-global-mobile-consumer-survey-second-edition.pdf)

352 Donnachie, C., Hunt, K., Mutrie, N., Gill, J. M. R., & Kelly, P. (2020). Responsiveness of
353 Device-Based and Self-Report Measures of Physical Activity to Detect Behavior
354 Change in Men Taking Part in the Football Fans in Training (FFIT) Program. *Journal*
355 *for the Measurement of Physical Behaviour*, 3(1), 67–77.
356 <https://doi.org/10.1123/jmpb.2019-0018>

357 Dowd, K. P., Szeklicki, R., Minetto, M. A., Murphy, M. H., Polito, A., Ghigo, E., van der Ploeg,
358 H., Ekelund, U., Maciaszek, J., Stemplewski, R., Tomczak, M., & Donnelly, A. E. (2018).
359 A systematic literature review of reviews on techniques for physical activity
360 measurement in adults: A DEDIPAC study. *International Journal of Behavioral*

361 *Nutrition and Physical Activity*, 15(1), 15. [https://doi.org/10.1186/s12966-017-0636-](https://doi.org/10.1186/s12966-017-0636-2)

362 2

363 Evenson, K. R., Goto, M. M., & Furberg, R. D. (2015). Systematic review of the validity and

364 reliability of consumer-wearable activity trackers. *International Journal of Behavioral*

365 *Nutrition and Physical Activity*, 12(1), 159. [https://doi.org/10.1186/s12966-015-](https://doi.org/10.1186/s12966-015-0314-1)

366 0314-1

367 Evenson, K. R., & Spade, C. L. (2020). Review of Validity and Reliability of Garmin Activity

368 Trackers. *Journal for the Measurement of Physical Behaviour*, 3(2), 170–185.

369 <https://doi.org/10.1123/jmpb.2019-0035>

370 Feehan, L., Geldman, J., Sayre, E., Park, C., Ezzat, A., Yoo, J., Hamilton, C., & Li, L. (2018).

371 Accuracy of Fitbit Devices: Systematic Review and Narrative Syntheses of

372 Quantitative Data. *JMIR Mhealth Uhealth*, 6(8), e10527.

373 <https://doi.org/10.2196/10527>

374 Fokkema, T., Kooiman, T. J. M., Krijnen, W. P., Van Der Schans, C. P., & De Groot, M. (2017).

375 Reliability and Validity of Ten Consumer Activity Trackers Depend on Walking Speed.

376 *Medicine & Science in Sports & Exercise*, 49(4), 793–800.

377 <https://doi.org/10.1249/MSS.0000000000001146>

378 Folley, S., Zhou, A., & Hyppönen, E. (2018). Information bias in measures of self-reported

379 physical activity. *International Journal of Obesity*, 42(12), 2062–2063.

380 <https://doi.org/10.1038/s41366-018-0223-x>

381 Hicks, J. L., Althoff, T., Sosič, R., Kuhar, P., Bostjancic, B., King, A. C., Leskovec, J., & Delp, S. L.

382 (2019). Best practices for analyzing large-scale health data from wearables and

383 smartphone apps. *Npj Digital Medicine*, 2. [https://doi.org/10.1038/s41746-019-](https://doi.org/10.1038/s41746-019-0121-1)

384 0121-1

385 Natarajan, A., Pantelopoulos, A., Emir-Farinas, H., & Natarajan, P. (2020). Heart rate
386 variability with photoplethysmography in 8 million individuals: A cross-sectional
387 study. *The Lancet Digital Health*, 2(12), e650–e657. [https://doi.org/10.1016/S2589-](https://doi.org/10.1016/S2589-7500(20)30246-6)
388 7500(20)30246-6

389 NHS England. (2019). *Digital diabetes prevention rolled out as part of NHS Long Term Plan*.
390 [https://www.england.nhs.uk/2019/08/digital-diabetes-prevention-rolled-out-as-](https://www.england.nhs.uk/2019/08/digital-diabetes-prevention-rolled-out-as-part-of-nhs-long-term-plan/)
391 [part-of-nhs-long-term-plan/](https://www.england.nhs.uk/2019/08/digital-diabetes-prevention-rolled-out-as-part-of-nhs-long-term-plan/)

392 Omura, J. D., Carlson, S. A., Paul, P., Watson, K. B., & Fulton, J. E. (2017). National physical
393 activity surveillance: Users of wearable activity monitors as a potential data source.
394 *Preventive Medicine Reports*, 5, 124–126.
395 <https://doi.org/10.1016/j.pmedr.2016.10.014>

396 Pearce, M., Bishop, T. R. P., Sharp, S., Westgate, K., Venables, M., Wareham, N. J., & Brage,
397 S. (2020). Network Harmonization of Physical Activity Variables Through Indirect
398 Validation. *Journal for the Measurement of Physical Behaviour*, 3(1), 8–18.
399 <https://doi.org/10.1123/jmpb.2019-0001>

400 Prince, S. A., Adamo, K. B., Hamel, M. E., Hardt, J., Gorber, S. C., & Tremblay, M. (2008). A
401 comparison of direct versus self-report measures for assessing physical activity in
402 adults: A systematic review. *International Journal of Behavioral Nutrition and*
403 *Physical Activity*, 5(1), 56. <https://doi.org/10.1186/1479-5868-5-56>

404 Redenius, N., Youngwon, K., & Wonwoo, B. (2019). Concurrent validity of the Fitbit for
405 assessing sedentary behavior and moderate-to-vigorous physical activity. *BMC*
406 *Medical Research Methodology*, 19, 29. <https://doi.org/10.1186/s12874-019-0668-1>

407 Reeves, M. M., Marshall, A. L., Owen, N., Winkler, E. A. H., & Eakin, E. G. (2010). Measuring
 408 Physical Activity Change in Broad-Reach Intervention Trials. *Journal of Physical*
 409 *Activity and Health*, 7(2), 194–202. <https://doi.org/10.1123/jpah.7.2.194>
 410 Stamatakis, E., Owen, K. B., Shepherd, L., Drayton, B., Hamer, M., & Bauman, A. E. (2021). Is
 411 Cohort Representativeness Passé? Poststratified Associations of Lifestyle Risk Factors
 412 with Mortality in the UK Biobank. *Epidemiology (Cambridge, Mass.)*, 32(2), 179–188.
 413 <https://doi.org/10.1097/EDE.0000000000001316>
 414 Strain, T., Brage, S., Sharp, S. J., Richards, J., Tainio, M., Ding, D., Benichou, J., & Kelly, P.
 415 (2020). Use of the prevented fraction for the population to determine deaths
 416 averted by existing prevalence of physical activity: A descriptive study. *The Lancet*
 417 *Global Health*, 8(7), e920–e930. [https://doi.org/10.1016/S2214-109X\(20\)30211-4](https://doi.org/10.1016/S2214-109X(20)30211-4)
 418 Strain, T., Wijndaele, K., & Brage, S. (2019). Physical Activity Surveillance Through
 419 Smartphone Apps and Wearable Trackers: Examining the UK Potential for Nationally
 420 Representative Sampling. *JMIR MHealth and UHealth*, 7(1), e11898.
 421 <https://doi.org/10.2196/11898>
 422 Swartz, A. M., Rote, A. E., Cho, Y. I., Welch, W. A., & Strath, S. J. (2014). Responsiveness of
 423 motion sensors to detect change in sedentary and physical activity behaviour. *British*
 424 *Journal of Sports Medicine*, 48(13), 1043–1047. [https://doi.org/10.1136/bjsports-](https://doi.org/10.1136/bjsports-2014-093520)
 425 [2014-093520](https://doi.org/10.1136/bjsports-2014-093520)
 426 Thompson, W. R. (2019). WORLDWIDE SURVEY OF FITNESS TRENDS FOR 2020. *ACSM's*
 427 *Health & Fitness Journal*, 23(6), 10–18.
 428 <https://doi.org/10.1249/FIT.0000000000000526>

429 Troiano, R. P., Stamatakis, E., & Bull, F. C. (2020). How can global physical activity
430 surveillance adapt to evolving physical activity guidelines? Needs, challenges and
431 future directions. *British Journal of Sports Medicine*, 54(24), 1468–1473.

432 Yao, J., Tan, C. S., Chen, C., Tan, J., Lim, N., & Müller-Riemenschneider, F. (2020). Bright
433 spots, physical activity investments that work: National Steps Challenge, Singapore: a
434 nationwide mHealth physical activity programme. *British Journal of Sports Medicine*,
435 54(17), 1047–1048. <https://doi.org/10.1136/bjsports-2019-101662>

436