



Bose, Rahul, Bhattacharya, Ekta, Pramanik, Arindam, Hughes, Thomas A. ORCID logoORCID: <https://orcid.org/0000-0003-1169-3386> and Biswas, Suparna Mandal (2021) Potential oil resources from underutilized seeds of *Sterculia foetida*, L. - Quality assessment and chemical profiling with other edible vegetable oils based on fatty acid composition, oxidative stability, antioxidant activity and cytotoxicity. *Biocatalysis and Agricultural Biotechnology*, 33 (102002).

Downloaded from: <https://ray.yorks.ac.uk/id/eprint/8729/>

The version presented here may differ from the published version or version of record. If you intend to cite from the work you are advised to consult the publisher's version:
<http://dx.doi.org/10.1016/j.bcab.2021.102002>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repositories Policy Statement](#)

RaY

Research at the University of York St John

For more information please contact RaY at
ray@yorks.ac.uk

Research Paper

Potential oil resources from underutilized seeds of *Sterculia foetida*, L. - quality assessment and chemical profiling with other edible vegetable oils based on fatty acid composition, oxidative stability, antioxidant activity and cytotoxicity

Short title: Potential oil resources from seeds of *Sterculia foetida*, L.

Rahul Bose^a, Ekta Bhattacharya^a, Arindam Pramanik^b, Thomas Hughes^b and Suparna Mandal Biswas*^a

^a Agricultural and Ecological Research Unit, Indian Statistical Institute,

203, B.T. Road, Kolkata 700108, INDIA

Tel: (+91)(033)25753225, Fax: (+91)(033)25753049

Email: suparna@isical.ac.in

^b School of Medicine, Wellcome Trust Brenner Building, St James's University Hospital,

University of Leeds, LS9 7TF, UK

Email: T.Hughes@leeds.ac.uk

1

¹ *Corresponding Author: Suparna Mandal Biswas, Agricultural and Ecological Research Unit, Indian Statistical Institute, 203, B. T. Road, Calcutta 700 108, INDIA. Tel: (+91) (033) 25753225, Fax: (+91) (033)25753049. Email: suparna@isical.ac.in

Abstract

Vegetable oils are integral part in production of manufactured food both in domestic and industrial scale. Vegetable oil market is increasing upward with a CAGR of 3.25% during the forecast period (2019-2024) all around the world. Therefore, it is necessary to find alternative sources of vegetable oil to fulfil the scarcity in the market. Seeds of *Sterculia foetida*, L. yielded considerable amount of oil (58.7 g/100g) as compared to other vegetable oils (such as sunflower, ground nut, mustard, soybean). Fatty acids composition of all the five tested oils showed that total fatty acid as well as unsaturated fatty acid percentage is higher in Sterculia seed oil. Proximate and mineral composition analysis suggested that Sterculia oil is a good source of protein, lipids, macro and micronutrients. Lowest TOTOX value (2.67) and higher iodine value (132-144) indicated its higher oxidative stability and presence of greater number unsaturated bands in the fatty acid moieties which is also beneficial for human health. Sterculia oil exhibited lower IC₅₀ values in DPPH (825.73 µg/ml), and NO (111.98 µg/ml) radical scavenging assays. In case of ABTS radical scavenging activity, no significant differences were observed in groundnut, mustard, sterculia and soybean except sunflower. Sterculia oil did not exhibit any cytotoxic effect on both normal and cancerous cell lines even at concentrations of 40µg/ml as evident from MTT assay. Thus, seed oil of *Sterculia foetida* may be a cost-effective and viable source of safe nutritious edible oils to combat the present market demand.

Key words: Edible vegetable oil; *Sterculia foetida*, L; Fatty acids; Oxidative stability; Cytotoxic activity

1. Introduction

With a rising standard of living as well as industrial growth, demand for vegetable oils (VO) is also soaring in global market. It is expected that in 2014, global vegetable oil market is

subjected to exceed 275 million metric tons due to the growing health consciousness of the consumers. Palm oil mainly dominated the world vegetable oil market by more than one-third of the total vegetable oil consumption and rest of the market occupied by soybean oil, canola oil and sunflower seed oil (Mielke, 2018). Though palm oil represents one potential source to meet this demand yet its consumption is associated with high risk of cardiovascular disease (WHO, 2003; Ismail et al, 2018) and also it is eco-destructive. In Asia-Pacific regions, vegetable oil market extended upward steadily with a CAGR of 5.4% over the analysis period due to the factors such as population growth, simultaneous growth of food commodities, changing dietary habits; rapid urbanization; improving living standards; increasing crop yields and oil production and growing biofuel production in countries such as Indonesia, Malaysia, Thailand, the Philippines, China and India. At present, there is a great demand of VO internationally, but alternative sources are very few. Therefore, it is necessary to find alternative sources of VO from underutilized oil resources to fulfil the global scarcity of VO applying non-agriculture land (Shi et al, 2019).

Sterculia foetida, L. commonly known as bastard poon, java olive, hazel sterculia, wild almond, is a large, straight, deciduous tree. It grows up to 40m in height and 3m in girth and its branches are arranged in whorls and spreading horizontally. This plant species is widely distributed in different geographical regions of India, south-east Asia and east coast of Africa. The lifespan of this tree is more than 100 years. Plants are readily propagated through seeds and need no special care (Staples and Herbst, 2005). It can be grown in non-cultivated or even in waste lands with minimum water and nutrient supply. The plant grows very fast and produces seeds within 2-3 years. The productivity of *S. foetida*, L. is reported to approx. 2000kg of seeds from one tree in one year.

This tree has immense potential for many therapeutic applications. Calcium content of *S. foetida* leaves is about 2.66% and also sufficient amount of protein and phosphorus are found in the leaves (Prakash & Kaviarasan, 2012). The leaves of this plant exhibit numerous medicinal properties such as laxative, carminative, anti-inflammatory, antioxidant, antimicrobial, cytotoxicity, anti-diabetic, anti-hyperlipidemic and insecticidal activities (Naik et al, 2004; Hussain et al, 2014; Mujumdar et al, 2000; Vital et al, 2010; Rani & Rajasekharreddy, 2010; Suganya et al, 2017). Methanol extract of *S. foetida* seed possess antioxidant activity (Galla, 2012). The de-oiled seed cake is rich in protein (28-89%) that can be used as animal and fish food supplements (Oliveira et al, 2000; Shamsundar & Paramjyothi, 2010).

Braga et al (2015) isolated and purified a new lectin with antibacterial and hemolytic activity from the seeds of *S. foetida*, L. Recently, a novel dye was extracted from the fruit shell waste of *S. foetida* which was applied on mulberry silk fabric to produce aesthetic coloration and wellness properties such as ultra-violet (UV) protection and antibacterial properties (Teli and Pandit, 2018).

The aim of our present study was to validate the edibility of *Sterculia* seed oil as an alternative safe vegetable oil. Therefore, in this study, in-depth characterization of fatty acid composition of *Sterculia* seed oil and four other commonly available vegetable oils along with their oxidative stability, radical scavenging activity, proximate and element content and cytotoxicity have been performed to validate its safe use as edible oil.

2. Materials and Methods

2.1. Collection of seed samples

S. foetida, L. seeds were collected from the campus of Indian Statistical Institute (ISI), Kolkata, India and nearby areas in the month of December to January from 2017 to 2018. Other commonly available edible oil seeds, namely sunflower, groundnut, mustard and soybean, were collected from the local market during the same time. Kolkata (22°33'N and 88°20'E) is located in the gangetic delta region of West Bengal and in the eastern part of India. It has a tropical wet-and-dry climate with an annual mean temperature of 26.8 °C (80 °F) and annual rainfall of 1,582 mm (62 in). This region has alluvial soil as it is near sea level, with the average elevation being 17 feet ([Weather Atlas, Kolkata, India, 2017](#)).

2.2. Extraction of oil

100 g of each ground seed samples were soaked in 500 ml of hexane in 1000 ml capacity extraction flasks ([Armah-Agyeman et al, 2016](#)). The mixture was then vortexed at 3000 rpm by Mechanical Stirrer (Model No. DC Stirrer NZ-1000s AC220V, EYELA) for 2 h and then filtered through sintered disc funnel. The recovered collected extract was concentrated in a rotary vacuum evaporator (EYELA, Model No. N1-NW) and oil was recovered in the concentrating flask.

2.3. Proximate Composition of the Seeds

Proximate composition of seeds was examined using the standard official methods of the Association of Official Analytical Chemists ([AOAC, 2006](#)).

2.4. Mineral Content of Seed Flour

Mineral content was determined by following the method of [Pinheiro et al. \(2010\)](#). Seed flour sample of 5.0 g was incinerated in a furnace at 550°C and the residues were dissolved in 50mL of 0.5M HNO₃ solution. The concentrations of K, Ca, Na, Mg, Zn, Cu, Mn, Pb and Fe were

measured by Atomic Emission Spectroscopy (AES) 4200 MP-AES SYSTEMS manufactured by AGILENT TECHNOLOGIES. A calibration curve was prepared by using standard metal solution.

2.5. Fatty Acid Composition of Oils

For determining the fatty acid composition of all the tested oils, fatty acid methyl ester (FAME) of oils were prepared according to the AOCS Official Method 996.06 (2001) with minor alterations described by [Symoniuk et al. \(2017\)](#).

FAME samples of all the five oils along with standard fatty acids were subjected to MDLC analysis (Model No. WATERS Technologies, 2695, LC System with 2487 inert Mass Selective Detector) for determination of the fatty acids compositions. MDLC analysis of the samples was done at Indian Institute of Chemical Biology, Jadavpur, Kolkata-700032, W.B. India. The gradient run was for 15 mins which was programmed as follows –First 0 min for column A it was 100% and for B it was 0%, next 2 mins it was 100% for A and for B it was 0%, next 2.5 mins it was 80% and 20% respectively for column A and B. For next 4 mins, 4.5 mins, 8 mins, 8.5 mins, 12 mins ,12.5 mins and 15 mins for column A and column B were 80% and 20%, 70 % and 30 % ,80% and 20% ,80% and 20%, 100% and 0% and 100% and 0% respectively. Total run was performed for 15 mins.

2.6. Antioxidant Activities

Different mechanisms such as free radical scavenging, reduction capacity, and metals chelation, are employed to evaluate the antioxidant potential of a substance or a complex mixture. In our study DPPH, ABTS and NO radical scavenging capacity assay were performed to detect the antioxidant activities of all the five tested oils namely sterculia, sunflower, groundnut, mustard and soybean.

2.6.1. DPPH radical scavenging assay

The DPPH radical scavenging activities of all the five oils were assessed in-vitro using 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical as described by Pavithra and Vadivukkarasi (2015).

2.6.2. ABTS scavenging capacity assay

ABTS radical scavenging activity of all the five oils were determined by ABTS radical cation decolorization assay (Re et al, 1999).

2.6.3. Nitric oxide scavenging capacity assay

Nitric oxide radical scavenging activity was measured spectrophotometrically according to the method described by (Jagetia et al 2004) with minor modification.

2.7. Determination of TOTOX value

TOTOX value i.e. overall oxidation state of oils is calculated by using the formula AV + 2PV. Better quality of the oil is determined by its lower TOTOX value (Maszewska et al, 2018).

Peroxide value (PV) was measured according to the AOCS Official Method 965.33 (AOAC, 1999), and the content of secondary oxidation products i.e. p-anisidine value (p-AV) was determined by the AOCS Official Method Cd 18-90 (AOCS, 2005).

2.8. Determination of Iodine value

Iodine value of oils is directly proportional to the degree of unsaturation of the product which indicates the oxidative stability of the oil and was determined following the method of Soares and Rocha (2018).

2.9. Cytotoxicity Assay

Cytotoxicity of *Sterculia* seed oil was measured by MTT assay on normal cell lines (MCF-10A and HB2 - breast non-cancerous cell lines) as well as cancerous cell lines (AU-565, BT-4T4 - breast cancer cell lines). These cell lines (MCF-10A, HB2, AU-565 and BT-4T4) were

procured from ATCC and maintained in Dulbecco's Modified Eagle Medium with 5% fetal calf serum and antibiotics in 37°C incubator. Cells were seeded in 24 well culture plates in DMEM growth medium at a density of 2.5×10^4 cells/well and incubated overnight in 37°C at 5% CO₂. After 18 h, cells were treated with different concentrations of *Sterculia* oil (0 – 160 µg/ml i.e. six set of experiments starting from concentrations 10 µg/ml , 20 µg/ml, 40 µg/ml, 80 µg/ml, 160 µg/ml including control) dissolved in DMSO, where the final concentration of DMSO was kept below 1%. Further, after 24 h cells were washed with 1× PBS and then incubated with 0.5 mg/ml of MTT solutions (in 1× PBS) for 2.5 h (Mosmann, 1983 and Stockert, et al, 2018). The Formazan crystals formed within the cells were dissolved using 400 µl of DMSO and absorbance of the solution was measured at 570 nm on a multi well plate reader (Biotech Instruments, USA).

2.10. Statistical Analysis

Values obtained from biochemical tests and antioxidant assays were analysed using independent sample T-test and the significance level was set at $p < 0.05$. All the experiments were replicated three times, and the data was represented as mean values and standard deviation of the same. SPSS Statistics 19 software for Windows had been used for all statistical analyses (SPSS 2009).

3. Results

3.1. Proximate Composition of Seeds

Proximate composition values of all the five tested oils namely *sterculia*, sunflower, groundnut, mustard and soybean are shown in Table 1. Seed kernel of *S. foetida* produces 58.7g of oil per 100g of seeds as compared to 47.5 g, 49.8 g, 31.4 g and 16.3 g from seeds of sunflower, ground nut, mustard and soybean respectively. Highest lipid content was found in

sterculia (58.4 g/100g) followed by sunflower (51 g), groundnut (50 g), mustard (36 g) and soybean (18.39g). Significant amount of protein was found in sterculia (38.43 g), groundnut (38.61 g) and soybean (37.69 g) per 100g of seeds. Soybean seeds contain higher amount of soluble sugar (16.24 g/100g). Sterculia seeds retain lowest amount of moisture (5.28 g/100g) in comparison to other tested oils. In the case of above mentioned oils, the total proximate compositions of the oilseeds did not differ significantly as the seeds of Sterculia contain the highest amount of oil and most minerals compared to other oils shown in [Table-1](#).

3.2. Mineral Composition of Seed Flour

Among the heavy metals, Sterculia seeds contained lowest amount of copper (16.67 mg/kg), lead (12.6 mg/kg), manganese (10.6mg/kg), iron (38.67 mg/kg) but high amount of Zinc (138.33 mg/kg) which is found to be higher in sterculia as compared to other tested seeds. Highest amount of lead (48.67 mg/kg) and iron (146.33 mg/kg) were detected in mustard seeds whereas highest amount of copper (52.67 mg/kg) and manganese (62.2 mg/kg) were recorded from groundnut seeds ([Table 1](#)). Among the alkaline earth metals only sodium (350.67 mg/kg), magnesium (3436 mg/kg) and potassium (19858 mg/kg) is found in highest quantity in the seeds of sterculia compared to other oils as shown in the ([Table 1](#)). Only calcium (5218.33 mg/kg) is present in highest quantity in the seeds of mustard compared to other oil seed.

3.3. Analyses of fatty acid composition

The MDLC analysis of FAME samples of all the five tested oils shows the fatty acid composition as shown in ([Table 2](#)).

In the five oils, the 6 fatty acids presents are linolenic acids (C18:3), linoleic acids (C18:2), palmitic acid (C16:0), myristic acid (C14:0), oleic acid (C18:1) and sterculic acid

(C19:1). Of these the linolenic and linoleic acids are the polyunsaturated fatty acids (PUFA), palmitic and myristic acid are the saturated fatty acids (SFA) and oleic acid is the monounsaturated fatty acids (MUFA) and only sterculic acid is the cyclopropenoid fatty acids (CFA). In the five oils, the carbon number in the fatty acids varies from C₁₄ to C₁₉. Total SFA was present higher in stercuria oil (1.29 mg/ml) and lowest in groundnut oil (0.08mg/ml) other than that rest other oils i.e sunflower oil, mustard oil and soybean oil consists of 0.23mg/ml, 0.77mg/ml and 0.77 mg/ml respectively. Total MUFA was present higher in sunflower oil (0.62mg/ml) and lowest in stercuria oil (0.016mg/ml) and mustard, soybean and groundnut oil consists of 0.18mg/ml, 0.14mg/ml and 0.02 mg/ml respectively. Finally total PUFA were present in higher quantity in stercuria oil (1.44 mg/ml) and lowest in sunflower oil (0.56 mg/ml). The remaining mustard, groundnut and soybean oil consists of (0.88mg/ml), (1.25mg/ml) and (0.72 mg/ml) fatty acids respectively. Calculating the total fatty acids it shows that stercuria oil consists of highest amount of fatty acids i.e 3.296 mg/ml and groundnut oil consists the lowest 1.35 mg/ml.

3.4. Determination of TOTOX Value

Oxidation of oil over time is measured by peroxide value (PV), para-anisidine value (AV) and TOTOX value as shown in [Table 3](#).

Highest para-anisidine (p-AV) value was revealed by groundnut oil (5.45) followed by mustard oil (5.09) whereas lowest p-AV value was detected in sunflower oil (2.99). Stercuria and soybean oil showed moderate p-AV value i.e. 3.73 and 3.77 respectively.

Peroxide value of all the five tested oils ranges from 0.018 – 0.025 meq./kg oil. Stercuria oil showed lowest TOTOX value (2.67) with higher oxidative stability followed by sunflower

(3.04) and soybean (3.81) oil. Higher TOTOX values were revealed by groundnut (5.48) and mustard oil (5.13).

3.5. Antioxidant activities

Low IC₅₀ value is inversely related to high antioxidant capacity of the extract (Rufino et al, 2009).

3.5.1. DPPH Radical Scavenging Activity

DPPH radical scavenging activity of all the five oils were expressed in IC₅₀ (µg/ml) with BHT as standard as shown in Figure 1. Highest antioxidant activity with lowest IC₅₀ value (797.98 µg/ml) was exhibited by soybean oil followed by groundnut (815.19 µg/ml) and sterculia (825.96 µg/ml) oil. Mustard and sunflower oil revealed least DPPH radical scavenging activity with IC₅₀ value of 1858.89 µg/ml and 1557.63 µg/ml respectively. In sterculia, groundnut and soybean oil, there is no significant differences whereas sunflower and mustard oil showed significant differences in DPPH activity.

3.5.2. ABTS Radical Scavenging Activity

ABTS radical scavenging activity of all the five oils were represented in IC₅₀ (µg/ml) as shown in Figure 1. Sunflower oil showed highest ABTS⁺ scavenging activity with significantly low IC₅₀ of 19.39 µg/ml. Soybean oil showed least antioxidant activity with highest IC₅₀ of 237 µg/ml, followed by sterculia (IC₅₀ of 220 µg/ml), mustard (IC₅₀ of 199 µg/ml) and groundnut (IC₅₀ of 169 µg/ml). No significant differences were observed in ABTS activity of groundnut, mustard, sterculia and soybean except sunflower.

3.5.3. NO Radical Scavenging Activity

NO radical scavenging activity of all the five oils expressed in IC₅₀ (µg/ml) is given in Figure 1. Sterculia oil exhibited highest NO radical scavenging activity with low IC₅₀ value of 114.98

µg/ml, followed by mustard (121.97 µg/ml), groundnut (123.14 µg/ml), soybean (230.24 µg/ml) and sunflower (266.64 µg/ml) oils. NO radical scavenging activity did not show any significant differences in all the five tested oils.

3.6. Cytotoxicity Assay

Figure 2 showed the result of cytotoxic activity of sterculia seed oil on both normal (MCF-10A and HB-2) and cancerous cell lines (BT-474 and AU-565) including control with DMSO on MCF-10A cell line. From MTT assay it was revealed that the sterculia oil did not show any cytotoxicity on both normal and cancer cell lines. On cancerous cell lines namely BT-474 and AU-565, sterculia oil at 10 µg/ml concentration exhibited 92.91% and 90.07% survivability of cells respectively, at 20 µg/ml concentration 92.38% and 85.12%, at 40 µg/ml concentration 89.47% and 80.64% and at 80 µg/ml concentration 82.46% and 74.01% survivability of cells were noticed. On normal cell lines namely MCF-10A and HB-2, sterculia oil at 10 µg/ml concentration showed 94.40% and 92.08% survivability of cells respectively whereas at 20 µg/ml concentration 92.62% and 85.35%, at 40 µg/ml concentration 86.61 and 84.42% and at 80 µg/ml concentration 80.56% and 76.89% survivability of cells were detected. The control experiment was done with DMSO on all cell lines in the same concentrations as the tested samples. Minor reduction in survivability of cells was observed due to the presence of DMSO as the solvent.

4. Discussion

Vegetable oil market in the world is expanding upward with a CAGR of 3.25% during forecast period from 2019 to 2024. Domestic consumption of edible oils in Asian Pacific countries like India, China, Indonesia, and Malaysia has increased substantially over the years due to upliftment of economic condition, urbanization, changing dietary habits and proclivity

of processed foods. The country's vegetable oil consumption was at 23 million tonnes in 2017 and it will be expanded by three per cent annually to exceed 34 million tonnes by 2030 according to the Rabo Research Report (PTI, Mumbai, June 25, 2018). Because of stagnant domestic vegetable oil supplies, over increasing demand will be filled by extending import volumes. Thus, it is high time to explore alternative rich natural source of vegetable oil and we show here that Sterculia seed oil could provide a viable source of nutritious edible oil.

Seeds of *S. foetida* yielded substantial amounts of oil (58.7 g per 100 g seed). The moisture content of Sterculia seeds was 5.28%, which is low as compared to other tested oils. Due to low moisture content, the seeds of *S. foetida* may have a prolonged shelf life. The seeds contained significant amounts of crude oil, protein, lipid and minerals that include heavy metals and alkaline earth metals. Heavy metals were present in very low amount except zinc which is essential for proper functioning of the immune system. Alkaline earth metals that are present includes magnesium, sodium and potassium present in higher quantity in sterculia oil compared to other oilseeds except calcium that is present in lower quantity. These earth metals potassium and sodium are electrolytes needed for the body to function normally and help maintain fluid and blood volume in the body, and magnesium is necessary for the formation of bone and teeth and for normal nerve and muscle function.

Fatty acids like palmitic acid, linoleic acid, linolenic acids, stearic acid and oleic acids were common in all the five oils. Results reveal that sterculia oil consists of highest amount of fatty acids, which includes the SFA, MUFA and PUFA. Highest percentage of polyunsaturated fatty acids such as linoleic acids and linolenic acids and monounsaturated fatty acids such as oleic acids along with sterculic acid (CFA) are present in sterculia oil that helps in increasing the high density lipoprotein (HDL) i.e the good cholesterol which assists in the removal of triacyl glycerols from the bloodstream (J.Lunn, & Theobald, 2006).

Unsaturated fats helps to reduce the risk of heart disease and lower the cholesterol as they replace saturated fats in the diet [EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) 2010]. Along with this the data also shows that PUFA are present in higher quantity than the MUFA in the sterculia oil. This is because the PUFA has more than one double bond in their structure than MUFA that help lower blood cholesterol and also contain omega-3 and omega-6 fatty acids that the body needs for proper brain function and cell growth. Omega-3 fats lower trygliceride levels and increases the HDL (good cholesterol) levels (Schriber, A. Medline Plus NIH US National Library of Medicine). Moreover, the unique cyclopropenoid fatty acid i.e. sterculic acid [namely 8-(2-Octacyclopropen-1-yl) octanoic acid] was found in the Sterculia seed oil (Kale et al, 2011; Vipunungeun and Chanida, 2009). Sterculic acid is a potent natural product to fight against obesity by suppressing a bodily enzyme associated with insulin resistance, which could indirectly help with reducing belly fat (Bao et al, 2003). It is also known to inhibit of SCD1 (Stearoyl-CoA desaturase-1), a major enzyme involved in the control of lipid metabolism and has emerged as a potential therapeutic target for reducing obesity and its associated metabolic complications including insulin resistance and hepatic steatosis (Ortinau, et al, 2013). This sterculic acid directly inhibits SCD activity, possibly by a turnover-dependent reaction, without affecting the processes required for adipocyte differentiation, *scd* gene expression or SCD protein translation (Gomez et al, 2003). So Sterculia oil has a promise to act to reduce some factors causing obesity.

Lowest TOTOX value and higher iodine value of Sterculic oil indicates its higher oxidative stability and presence of greater number of double bonds in the fatty acid moieties which further supports that this oil would be beneficial for edible purposes.

No significant differences in IC₅₀ values among the oils were observed as measured by NO radical scavenging activity. ABTS radical scavenging activity showed significant difference

only with sunflower oil whereas DPPH showed differences with sunflower and mustard oil. So Sterculia oil is comparable to other vegetable oil based on their radical scavenging activity.

Furthermore, sterculia oil did not reveal any cytotoxic effect even at 40µg/ml concentration against normal (MCF-10A and HB-2) and cancerous cell lines (BT-474 and AU-565). So, it can be suggested that this oil has no toxicity on human beings and safe for human consumption, however, further studies are to be performed in order to ensure the same. All the parameters of sterculia oil recommends that the seed oil of *S. foetida*, L. may be an alternative source of safe edible oil. As an additional fact, the seeds of *Sterculia apetala* are reported to be commonly used in some tropical areas in Mexico for human and animal nutrition (Herrera-Meza et al, 2014). Consumption of *S. apetala* seed oil in Zucker rats reduces anxiety-like behaviour and some behavioural alterations in locomotor activity tests (Herrera-Meza et al, 2017).

Because of the need of edible oil in the global market, seeds of this tree would be a viable resource of nutritious, non-toxic edible vegetable oil. This plant has a wide range of distribution in all around the world so it is not invasive nor eco-destructive. It can be cultivated in un-utilized lands or even in waste lands with very nominal water and nutrient supply which is very relevant to developing countries. Moreover, this tree can be used for backyard planting, coastal protection and stabilization, urban greening, shade tree, large road side tree, wild grafting (Orwa et al, 2009).

5. Conclusions

From our study it was revealed that *Sterculia foetida* seeds yield considerable amount of oil (58.7 g/ 100 g) among all the tested vegetable oil seed samples. Proximate and mineral composition analysis suggests that these seeds are rich and could be considered

as an alternative source of oil, protein, and micronutrients. Fatty acid composition of all the tested oils showed that seed oil of sterculia contains highest amount of total fatty acids that includes both PUFA and SFA and also CFA. So, it might be substituted with other edible vegetable oils. This oil has a very low TOTOX value and high iodine value as compared to other vegetable oils which indicates its higher oxidative stability and higher degree of unsaturation which might be beneficial for our health. Radical scavenging activity supported the similar nature of sterculia oil to the other edible oils. This oil does not have any toxicity on human health as revealed from MTT assay on both normal and cancerous cell lines. Based on all the tested parameters, we may conclude that this oil may serve as an alternative, viable source of safe edible oil.

Conflicts of Interest

The authors declare that they have no conflicts of interest regarding publication of this paper.

Authorship Contribution Statement

Ekta Bhattacharya & Rahul Bose – Investigation, collection of test data, formal analysis, drafting the article; Suparna Mandal Biswas – Writing original draft, funding acquisition, project administration, validation, supervision. Thomas Hughes and Arindam Pramanik – Design of the experiment, investigation, formal analysis and made the critical revision of the article.

Acknowledgements

We convey our sincere gratitude to our Director, Prof. Sanghamitra Bandhopadhyay for providing Laboratory facilities and financial support for performing this research work. We are thankful to Prof. Susmita Mukhopadhyay, Prof-in-Charge, Biological Sciences Division; Dr.

Sabyasachi Bhattacharya, Head, Agricultural and Ecological Research Unit, Indian Statistical Institute, for their encouragement, valuable advice and laboratory facilities. We thank to Dr. Ramdhan Majhi of Central Instrument Facility of Indian Institute of Chemical Biology, Kolkata, INDIA for carrying out the MDLC (multi-dimensional liquid chromatography) analysis of the plant oil samples.

References

AOAC, *Official Methods of Analysis of the Association of Official Analytical Chemists*, 2006. Arlington, Va, USA, 18th edition

AOAC, *Official Methods of Analysis of the Association of Official Analytical Chemists*. 1999. Washington, DC, USA, 16th edition.

AOCS, *Official Methods of Analysis of the Association of Official Analytical Chemists*. 2005. Official Methods of Analysis Cd 8b-90. Champaign, IL: AOCS Publishing

Armah-Agyeman, G., Gyamerah, M., Biney, P. O., & Woldeesenbet, S. 2016. Extraction and characterization of triglycerides from coffee weed and switch grass seeds as potential feed stocks for biodiesel production. *Journal of the Science of Food and Agriculture*, 96 (13), 4390-4397.

Bao, X., Thelen, J. J., Bonaventure, G., & Ohlrogge, J. B. 2003. Characterization of Cyclopropane Fatty-acid Synthase from *Sterculia foetida*. *Journal of Biological Chemistry*, 278 (15), 12846-12853.

Braga, A. A., e Lacerda, R. R., de Vasconcelos Medeiros, G. K. V., Gonçalves, G. F., Pessoa, H. D. L. F., Cardoso, J. D., de Almeida Gadelha, C.A., da Silva, B.A., and Santi-Gadelha, T.

2015. Antibacterial and hemolytic activity of a new lectin purified from the seeds of *Sterculia foetida*, L. *Applied biochemistry and biotechnology*, 175 (3), 1689-1699.
- EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) (2010) Scientific Opinion on Dietary Reference Values for fats, including saturated fatty acids, polyunsaturated fatty acids, © 2020 The Author. Nutrition Bulletin published by John Wiley & Sons Ltd on behalf of British Nutrition Foundation. Nutrition Bulletin, 45, 271–280 278 I. A. Brouwer monounsaturated fatty acids, trans fatty acids, and cholesterol. EFSA Journal 8 1461
- Galla, N. R. 2012. In vitro Antioxidants activity of *Sterculia foetida* Linn seed methanol extract. *American Journal of PharmTech Research*, 2 (6), 572-581.
- Gomez, F. E., Bauman, D. E., Ntambi, J. M., & Fox, B. G. 2003. Effects of sterculic acid on stearoyl-CoA desaturase in differentiating 3T3-L1 adipocytes. *Biochemical and biophysical research communications*, 300 (2), 316-326.
- Herrera-Meza, S., Martínez, A. J., Sánchez-Otero, M. G., Mendoza-López, M. R., García-Barradas, O., Ortiz-Viveros, G. R., & Oliart-Ros, R. M. 2014. Fatty acid composition and some physicochemical characteristics of *Sterculia apetala* seed oils. *Grasas y Aceites*, 65 (3), 039. ISSN-L: 0017-3495.doi: <http://dx.doi.org/10.3989/gya.0223141>
- Herrera-Meza, S., Rodríguez-Landa, J. F., Martínez, A. J., Herrera-Meza, G., Fernández-Demeneghi, R., Reyes-Saldana, K., & Oliart-Ros, R. M. 2017. Behavioral Effect of *Sterculia apetala* Seed Oil Consumption in Male Zucker Rats. *Journal of medicinal food*, 20 (11), 1133-1139.
- Hussain, S. S., Janarthan, M., Anusha, S. K., & Ranjani, M. 2014. Preclinical evaluation of anti-diabetic and antihyperlipidemic activity of methanolic extract of *Sterculia foetida* leaves by using wistar albino rats. *Indian Journal of Research in Pharmacy and Biotechnology*, 2 (6), 1430-1438.

391 Ismail, S. R., Maarof, S. K., Ali, S. S., & Ali, A. 2018. Systematic review of palm oil consumption
 392 and the risk of cardiovascular disease. *PLoS One*, 13 (2). e0193533.

393 J.Lunn, H.E. Theobald, 2006. The health effects of dietary unsaturated fatty acids. *Nutrition*
 394 *Bulletin*, British Nutrition Foundation.

395 Jagetia, G. C., Rao, S. K., Baliga, M. S., & S. Babu, K. 2004. The evaluation of nitric oxide
 396 scavenging activity of certain herbal formulations in vitro: a preliminary
 397 study. *Phytotherapy Research*, 18 (7), 561-565.

398 Kale, S. S., Darade, V., & Thakur, H. A. 2011. Analysis of fixed oil from *Sterculia foetida*
 399 Linn. *International Journal of Pharmaceutical Sciences and Research*, 2 (11), 2908-2914.

400 Maszewska, M., Florowska, A., Dłużewska, E., Wroniak, M., Marciniak-Lukasiak, K., &
 401 Żbikowska, A. 2018. Oxidative stability of selected edible oils. *Molecules*, 23 (7), 1746;
 402 doi: 10.3390/molecules23071746

403 MEDLINE PLUS, NIH-U.S. National library of medicine.

404 Mielke, T. 2018. World markets for vegetable oils and animal fats. In M. Kaltschmitt, & U.
 405 Neuling (Eds.). *Biokerosene: Status and prospects* (pp. 147–188). Berlin, Heidelberg:
 406 Springer Berlin Heidelberg.

407 Mosmann, T. 1983. Rapid colorimetric assay for cellular growth and survival: application to
 408 proliferation and cytotoxicity assays. *Journal of immunological methods*, 65 (1-2), 55-
 409 63.

410 Mujumdar, A. M., Naik, D. G., Waghole, R. J., Kulkarni, D. K., & Kumbhojkar, M. S. 2000.
 411 Pharmacological studies on *Sterculia foetida* leaves. *Pharmaceutical biology*, 38 (1), 13-
 412 17.

413 Naik, D. G., Mujumdar, A. M., Waghole, R. J., Misar, A. V., Bligh, S. A., Bashall, A., & Crowder,
 414 J. 2004. Taraxer-14-en-3 β -ol, an anti-inflammatory compound from *Sterculia foetida*
 415 L. *Planta medica*, 70 (01), 68-69.

416 Oliveira, J. T. A., Vasconcelos, I. M., Bezerra, L. C. N. M., Silveira, S. B., Monteiro, A. C. O., &
 417 Moreira, R. A. 2000. Composition and nutritional properties of seeds from *Pachira*
 418 *aquatica* Aubl, *Sterculia striata* St Hil et Naud and *Terminalia catappa* Linn. *Food*
 419 *Chemistry*, 70 (2), 185-191.

420 Ortainau, L. C., Nickelson, K. J., Stromsdorfer, K. L., Naik, C. Y., Pickering, R. T., Haynes, R. A.,
 421 Fritsche, K.L., and Perfield, J. W. 2013. Sterculic oil, a natural inhibitor of SCD1,
 422 improves the metabolic state of obese OLETF rats. *Obesity*, 21 (2), 344-352

423 Orwa, C., Mutua, A., Kindt, R., Jamnadass, R., and Simons, A. (2009). Agro forest tree
 424 database: a tree species reference and selection guide version 4.0. World Agroforestry
 425 Centre ICRAF, Nairobi, KE.

426 Pavithra, K., & Vadivukkarasi, S. 2015. Evaluation of free radical scavenging activity of various
 427 extracts of leaves from *Kedrostis foetidissima* (Jacq.) Cogn. *Food Science and Human*
 428 *Wellness*, 4 (1), 42-46.

429 Pinheiro, C., Baeta, J. P., Pereira, A. M., Domingues, H., & Ricardo, C. P. 2010. Diversity of seed
 430 mineral composition of *Phaseolus vulgaris* L. germplasm. *Journal of Food Composition*
 431 *and Analysis*, 23 (4), 319-325.

432 Prakash, Y. G., & Kaviarasan, L. G. V. 2012. Promising pharmaceutical prospective of java olive-
 433 *Sterculia foetida* Linn (Sterculiaceae). *International Journal of Pharmaceutical Sciences*
 434 *Review and Research*, 2, 93-96.

435 Rabo Research Report, 2018. Published in PTI Mumbai on 25th June, 2018.

436 [https://www.thehindubusinessline.com/economy/agri-business/indias-edible-oil-](https://www.thehindubusinessline.com/economy/agri-business/indias-edible-oil-consumption-to-exceed-34-million-tonnes-by-2030-report/article24254536.ece)

437 [consumption-to-exceed-34-million-tonnes-by-2030-report/article24254536.ece](https://www.thehindubusinessline.com/economy/agri-business/indias-edible-oil-consumption-to-exceed-34-million-tonnes-by-2030-report/article24254536.ece)

438 Rani, P. U., & Rajasekharreddy, P. 2010. Insecticidal activity of (2n-octylcycloprop-1-enyl)-

439 octanoic acid (I) against three Coleopteran stored product insects from *Sterculia*

440 *foetida* (L.). *Journal of pest science*, 83(3), 273-279.

441 Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., & Rice-Evans, C. 1999. Antioxidant

442 activity applying an improved ABTS radical cation decolorization assay. *Free radical*

443 *biology and medicine*, 26 (9-10), 1231-1237.

444 Rufino, M. S., Alves, R.E., Fernandes, F.A., and Brito, E.S. 2011. Free radical scavenging

445 behavior of ten exotic tropical fruits extracts. *Food Research International*, 44 (7), 2072-

446 2075

447 Schriber, A. Medline Plus NIH US National Library of Medicine. *Sleep disordersoverview*. ADAM

448 *Editorial team Accreditation Health Care Commission*. [Actualizado el 2 de febrero de

449 2015][Internet]. [Acceso el 3 de abril 2016] Disponible en: <https://www.nlm.nih.gov/medlineplus/ency/article/000800.htm>.

450

451 Shamsundar, S. G., & Paramjyothi, S. 2010. Preliminary pharmacognostical and phytochemical

452 investigation on *Sterculia foetida* Linn. seeds. *African Journal of Biotechnology*, 9 (13),

453 1978-1989.

454 Shi, L., Zheng, L., Liu, R., Chang, M., Huang, J., Zhao, C., Jin, Q. and Wang, X. 2019. Potential

455 underutilized oil resources from the fruit and seed of *Rhus chinensis* Mill. *Industrial*

456 *Crops and Products*, 129, 339–344.

457 Soares, S., & Rocha, F. R. 2018. Fast spectrophotometric determination of iodine value in

458 biodiesel and vegetable oils. *Journal of the Brazilian Chemical Society*, 29 (8), 1701-

459 1706.

460 SPSS Inc. Released. PASW Statistics for Windows, Version 18.0. Chicago: 2009.

461 Staples G, Herbst DR. Tropical garden flora. Bishop Museum Press, 2005.

462 Stockert, J. C., Horobin, R. W., Colombo, L. L., & Blázquez-Castro, A. 2018. Tetrazolium salts
 463 and formazan products in Cell Biology: Viability assessment, fluorescence imaging, and
 464 labeling perspectives. *Acta histochemica*, 120 (3), 159-167.

465 Suganya, J., Viswanathan, T., Radha, M., Rathisre, P. R., & Marimuthu, N. 2017. In vitro
 466 Antibacterial Activity of different crude leaves extracts of *Sterculia foetida*
 467 Linn. *Research Journal of Pharmacy and Technology*, 10 (7), 2013-2017.

468 Symoniuk, E., Ratusz, K., and Krygier, K. 2017. Comparison of the oxidative stability of cold-
 469 pressed rapeseed oil using Pressure Differential Scanning Calorimetry and Rancimat
 470 methods. *European Journal of Lipid Science and Technology*, 119 (5), 1600182.

471 Teli, M. D., & Pandit, P. 2018. Application of *Sterculia foetida* fruit shell waste biomolecules
 472 on silk for aesthetic and wellness properties. *Fibers and Polymers*, 19 (1), 41-54.

473 Vipunungeun, N., & Palanuvej, C. 2009. Fatty acids of *Sterculia foetida* seed oil. *Journal of*
 474 *Health Research*, 23 (3), 157-157.

475 Vital, P. G., Velasco, R. N., Demigillo, J. M., & Rivera, W. L. 2010. Antimicrobial activity,
 476 cytotoxicity and phytochemical screening of *Ficus septica* Burm and *Sterculia foetida* L.
 477 leaf extracts. *Journal of medicinal plants research*, 4 (1), 58-63.

478 Weather Atlas, Kolkata, India – Climate data. Online Accessed on: 4 Apr. 2017 -
 479 <http://www.weatheratlas.com/en/india/kolkata-climate>.

480 World Health Organization. 2003. *Diet, nutrition, and the prevention of chronic diseases:*
 481 *report of a joint WHO/FAO expert consultation*.

482

483

Figure and Captions

Figure 1. Antioxidant activities of all the five oils (namely Sterculia, Sunflower, Groundnut, Mustard and Soybean) expressed in IC_{50} ($\mu\text{g/ml}$). All the data are mean \pm SD of triplicate measurements. Bars with '*' are significantly different. IC_{50} , inhibitory concentration; DPPH, DPPH radical scavenging activity; ABTS, ABTS radical scavenging activity; NO, nitrite scavenging activity.

Figure 2. MTT assay of Sterculia seed oil at different concentrations on normal (MCF-10A and HB2) and cancer (AU-565, BT-4T4) cell lines.

Table 1. Proximate and mineral compositions of *Sterculia foetida*, *L.* seeds along with sunflower, groundnut, mustard and soybean seeds.

Table 2. Comparative fatty acids profiling of all the five seed oils namely sterculia, sunflower, groundnut, mustard and soybean based on GCMS analyses of the FAME samples of respective oils.

Table 3. Iodine value and oxidation status of five seed oils namely Sterculia, Sunflower, Groundnut, Soybean and Mustard.


Figure 1


Figure 2

Table 1. Proximate and mineral compositions of *Sterculia foetida* seeds along with Sunflower, Groundnut, Mustard and Soyabean seeds.

	Sterculia	Sunflower	Groundnut	Mustard	Soybean
NUTRITIONAL COMPONENTS (g/100g)					
Oil Content	58.7± 2.37^a	47.5± 2.22^a	49.8± 1.62^a	31.4± 1.45^b	16.3± 1.19^c
Lipid Content	58.4± 4.52^a	51± 3.50^a	50± 6.52^a	36± 3.36^a	18.39± 4.52^b
Protein Content	38.43± 2.81^a	20.91± 3.08^c	38.61± 3.02^a	28.8± 4.05^b	37.69± 2.66^a
Soluble Sugar Content	1.73±1.34^c	2.54± 1.22^c	1.46± 1.26^c	6.74±1.05^b	16.24±2.73^a
Moisture Content	5.28± 3.22^b	7.18± 3.22^b	5.98± 2.01^b	9.12± 4.33^a	13.5± 2.44^a
ALKALINE EARTH METALS (mg/kg)					
Magnesium	3436 ±4.51^b	3251±3.61^a	590.33 ±6.55^a	3040±72.11^a	2756.67±40.5^a
Sodium	350.67±24.44^a	89.33±3.05^b	335±4.35^a	52.67±6.42^b	181±3.60^a
Potassium	19858±345.5^b	6455.67±6.0^b	14500±2.01^a	6882±65.21^b	5293.33±15.6^a
Calcium	590.33±4.51^c	3251±3.61^a	2002.33±5.8^b	5218.33±7.6^a	3124±5.29^a
HEAVY METALS (mg/kg)					
Copper	16.67± 4.16^b	20±2.03^b	52.67±2.08^a	38.67±1.15^a	48±3.0^a
Lead	12.6± 3.05^b	10.33±1.52^b	44±2.35^a	48.67± 8.14^a	38.33±5.50^a
Manganese	10.6± 1.15^b	21±2.01^b	62±2.64^a	41.33±2.5^a	41±1.03^a
Iron	38.67± 2.51^b	53.33±1.52^b	111±2.64^a	146.33±1.52^a	38.67±2.51^b
Zinc	138.33±2.08^a	52.67±3.05^b	75±5.56^a	38.67±1.15^b	52.67±1.52^b

Mean values ± standard deviation for $n = 3$ *Values (means± SD) with different index letters are statistically significantly different ($P < 0.05$).

Table 2. Fatty acid composition of all the five seed oils namely Sterculia, Sunflower, Groundnut, Mustard and Soybean based on MDLC analyses of the FAME samples of respective oils.

Name of the fatty acid	Fatty acid content (mg/ml)				
	Sterculia	Sunflower	Mustard	Groundnut	Soyabean
Sterculic acid (C 19:1) CFA	0.55± 0.01	0	0	0	0
Linolenic acid (C 18:3) PUFA	1.12± 0.01	0.52±0.01	0.76±0.00	1.25±0.05	0.62± 0.02
Linoleic acid (C18:2) PUFA	0.32±0.040	0.04±0.003	0.12±0.039	0	0.1± 0.02
Palmitic acid (C16:0) SFA	1.23± 0.16	0.20± 0.04	0.76±0.032	0.04±0.002	0.76± 0.029
Myristic acid (C14:0) SFA	0.06± 0.001	0.033± 0.001	0.015±0.002	0.04± 0.001	0.012± 0.009
Oleic Acid (C18:1) MUFA	0.016± 0.006	0.62± 0.13	0.18± 0.019	0.024± 0.011	0.142± 0.00
Total saturated fatty acids	1.29± 0.015	0.233± 0.025	0.775± 0.017	0.08± 0.012	0.772± 0.019
Total monounsaturated fatty acids	0.016± 0.006	0.62± 0.013	0.18± 0.019	0.024± 0.011	0.142± 0.00
Total polyunsaturated fatty acids	1.44± 0.020	0.56± 0.016	0.88± 0.019	1.25± 0.005	0.72± 0.015
Total cyclopropenoid fatty acids	0.55± 0.001	0	0	0	0
TOTAL	3.296± 0.013	1.413± 0.019	1.835± 0.018	1.354± 0.072	1.634± 0.023

Table 3. Iodine value and oxidation status of five seed oils namely Sterculia, Sunflower, Ground nut, Soyabean, Mustard.

SEED OIL	Para-Anisidine Value (p-AV)	Peroxide value (PV) (meq./kg oil)	TOTOX (2 PV + p-AV)	Iodine Value
Sterculia	3.73 ± 0.64	0.023 ± 0.0034	2.67 ± 0.72	132-144
Sunflower	2.99 ± 0.45	0.025 ± 0.0052	3.04 ± 0.46	122-140
Ground nut	5.45 ± 0.68	0.018 ± 0.0004	5.48 ± 0.68	87-106
Mustard	5.09 ± 0.87	0.021 ± 0.005	5.13 ± 0.88	94-111
Soyabean	3.77 ± 0.90	0.020 ± 0.0004	3.81 ± 0.90	120-134

The values are mean ± SD of three independent experiments

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

1. Rahul Bose - rahulbosebiotech@gmail.com,
2. Ekta Bhattacharya - ektabhattacharya1990@gmail.com,
3. Arindam Pramanik - A.Pramanik@leeds.ac.uk,
4. Thomas Hughes - T.Hughes@leeds.ac.uk,
5. Suparna Mandal Biswas – suparna@isical.ac.in

On behalf of all the contributing authors, I , Suparna Mandal Biswas (corresponding author of the article), hereby declare that we have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Suparna Mandal Biswas

(Suparna Mandal Biswas)
Corresponding Author
23.02.2021